

June 1976

LRP 103/76

TWO-DIMENSIONAL QUASI-LINEAR EVOLUTION OF THE
ELECTRON-BEAM-PLASMA INSTABILITY

K. Appert, T.M. Tran, and J. Vaclavik

Centre de Recherches en Physique des Plasmas
ECOLE POLYTECHNIQUE FEDERALE DE LAUSANNE

TWO-DIMENSIONAL QUASI-LINEAR EVOLUTION OF THE
ELECTRON-BEAM-PLASMA INSTABILITY

K. Appert, T.M. Tran, and J. Vaclavik

ABSTRACT

We report on the quasi-linear evolution of the electron-beam-plasma instability in a 2-D system. The numerical solutions of the basic equations show that a 2-D system evolves in a different way as compared with a 1-D system. After saturation the wave energy monotonically decreases with time while the width of its spectral distribution in k-space narrows rapidly. In some cases, the system reaches a state which is modulationally unstable, and consequently cannot be described by the weak turbulence theory.

In the problem of plasma heating, the turbulence excited by electron beams is of great importance. In many papers (see the reviews ^{1,2}), the authors have investigated the electron-beam-plasma interaction on the basis of the quasi-linear theory for the case when the turbulent oscillation spectrum is one-dimensional (wave vector k parallel to the motion of the beam). Such a model is appropriate when there is a magnetic field parallel to the beam in the plasma and when this field is strong enough to suppress oscillations that propagate at an angle to the beam axis. If there is a weak or no magnetic field in the plasma, so that $\omega_{pe} \gg \omega_{ce}$ (ω_{pe} and ω_{ce} are the plasma and the cyclotron frequencies of electrons, respectively) the turbulent spectrum becomes essentially three-dimensional. Such situations are met e.g. in astrophysical problems or in the inertial confinement of plasmas. At present not much is yet known about the quasi-linear behaviour of a three-dimensional turbulence excited by electron beams in a plasma. Attempts have been made to find at least certain general features of possible asymptotic states for the three-dimensional system^{3,4}. However, these states seem to be too artificial and they have never been observed in experiments². Thus the relaxation dynamics as well as the final state of the system remain unknown.

In this Letter we report on the two-dimensional quasi-linear evolution of the electron-beam-plasma instability. Since the problem exhibits axial symmetry with respect to beam axis the two-dimensional model is not restrictive⁵ and was only chosen for convenience. The fundamental equations of the quasi-linear theory for the electron-beam-plasma interaction read²:

$$\frac{\partial f}{\partial t} = \frac{\partial}{\partial v} \cdot \pi \sum_{\mathbf{k}} \frac{k_{\parallel} k_{\parallel}}{k^2} I_{\mathbf{k}} \delta(\omega - \mathbf{k} \cdot \mathbf{v}) \cdot \frac{\partial f}{\partial v}, \quad (1)$$

$$\frac{\partial I_{\mathbf{k}}}{\partial t} = \frac{2\pi I_{\mathbf{k}}}{k^2 \partial \epsilon / \partial \omega} \int d^2 v \, k_{\parallel} \cdot \frac{\partial f}{\partial v} \delta(\omega - \mathbf{k} \cdot \mathbf{v}), \quad (2)$$

where $\omega = 1 + 3k^2/2$, $\partial \epsilon / \partial \omega = 2 + 3k^2$, f is the velocity distribution function for electrons and $I_{\mathbf{k}}$ is the spectral distribution of the electrostatic field associated with the oscillations. Equations (1) and (2) are in dimensionless units; the units of time, space, velocity distribution function, and spectral distribution are, respectively, ω_{pe}^{-1} , λ_D , $m_e n / T_e$, and $4\pi n T_e$. Here m_e , T_e and n are the electron mass, temperature, and density respectively, and λ_D is the Debye length.

We have solved Eqs.(1) and (2) by the finite element method^{6,7}. The algorithm consists of the following stages. First, Eq.(1) is put into the Galerkin (weak) form, the natural (Neumann) boundary condition being imposed on the velocity distribution function. The approximate solution for f is then assumed to be a combination $\sum f_i e_i$ of the linear pyramid basis functions $e_i(\mathbf{v})$. In this manner, Eqs.(1) and (2) are transformed into a system of ordinary differential equations in time for the quantities f_i and $I_{\mathbf{k}}$. This system of equations is solved numerically by an implicit, four-level, time-centred, finite difference scheme.

The accuracy of the solutions was verified by checking the variations of the integrals of motion of Eqs.(1) and (2) (particle, momentum and energy densities) and by repeating the computations with different numbers of the basis functions and the wave modes, and with different time

steps. For typical results presented in this Letter we used about 300 basis functions and about the same number of wave modes. The initial distribution function was chosen as

$$f(t=0) = \frac{1}{2\pi(1+\xi)} \left\{ \exp(-v^2/2) + \xi \exp\left(-\frac{1}{2} \left[(v_x - u)^2 + v_y^2 \right] \right) \right\}$$

The values of the parameters ξ and u were varied within certain ranges in such a manner that the condition for the kinetic instability (the condition for Eqs.(1) and (2) to hold), viz. $\xi^{1/3} < 1/u$ was satisfied. Typically, the values of ξ were between 10^{-5} and $5 \cdot 10^{-2}$, and those of u between 4 and 10. The initial spectral distribution $I_k(t=0)$ was taken as a constant the value of which was varied within the range $10^{-2} - 10^{-6}$ times the maximal value of the spectral distribution reached at saturation.

In all sets of computations we have observed roughly two stages in the evolution of the system. They are determined by the time history of the total wave energy density defined as $W = \sum_k W_k = \sum_k \frac{1}{2} I_k \omega \partial \epsilon / \partial \omega$ (in nT_e units). As seen from Fig. 1, the wave energy density first increases from an initial level to its maximum value W_{sat} within a time t_{sat} . After the saturation it monotonically decreases. The other quantities shown in Fig. 1 are the widths of the wave energy density in k -space. They are defined as $\Delta k_\alpha = \left(\sum_k (k_\alpha - \langle k_\alpha \rangle)^2 W_k / W \right)^{1/2}$, where $\langle k_\alpha \rangle = \sum_k k_\alpha W_k / W$. It can be seen that in the first stage they remain more or less constant, the value of Δk_y being about twice the value of Δk_x . During the second stage both widths narrow rapidly. Towards the end of the observation time the spectrum is so narrow that $(\Delta k)^2 \ll W$ in some

cases. This fact indicates that the weak turbulence theory breaks down since the system becomes unstable with respect to the modulational instability of Vedenov and Rudakov⁸. Fig. 2 displays the dependence of the quantity W_{sat} on the parameters ξ and u . Here the solid lines represent the saturation energy computed for the 1-D system, the dots refer to the 2-D system. We notice that these values are very close to each other. Moreover, in all cases it turned out that the saturation time obeys the approximate formula $t_{\text{sat}} \approx 4 \lg(W_{\text{sat}}/W_0) / \gamma_{\text{max}}$, W_0 and γ_{max} being the initial wave energy density and the initial maximum growth rate, respectively. In Fig. 3 we present the spectral distribution of the field in the direction parallel to the beam axis ($k_y = 0$) at a few instants in time. Here the plots (a) and (b) refer to the 2-D and 1-D systems, respectively. We see a distinct difference between these two cases. In the 2-D system the spectrum moves up several half-widths along the k_x -axis whereas in the 1-D system its location remains the same. Furthermore, in the former case the spectrum is always narrower (at least by a factor 2) than in the latter. Fig. 4 shows two cross-sections of the velocity distribution function at several times. It can be observed that in the first stage the distribution function flattens along the v_x -axis in the resonant region of the velocity space. In fact, such a flattening occurs at different heights for all cross-sections along the lines $v_y = \text{const}$ in this region. At the same time the function broadens across these lines, the width being about twice the initial width of the beam. As the wave energy damps down during the second stage, a part of the bulk particles is heated in the direction parallel to the beam axis. Thus, a broad high-energy tail is formed on

the distribution function.

During the course of our investigations⁹ we were made aware¹⁰ of the work of Ivanov, Sobolyeva, and Yushmanov¹¹. With respect to the first stage of the quasi-linear evolution of the electron-beam-plasma instability in multi-dimensions their conclusions are similar to those presented in this Letter. However, the second stage where, in our opinion, the effects of multi-dimensions play the most important role was not treated in their paper. The possibility for the considered system to reach a state which cannot be described by the weak turbulence theory, viz. a modulationally unstable state, may be regarded as one of the main results of our treatment.

This work was supported by the Swiss National Science Foundation.

REFERENCES

- ¹R.C. Davidson, *Methods in Nonlinear Plasma Theory* (Academic, New York, 1972), pp. 174-197.
- ²A.A. Vedenov and D.D. Ryutov, *Reviews of Plasma Physics* (Consultants Bureau, New York, 1975), Vol. 6, pp. 1-76.
- ³V.L. Sizonenko and K.N. Stepanov, *Zh.Eksp.Teor.Fiz.* 49, 1197 (1965) [*Sov.Phys.JETP* 22, 832 (1966)].
- ⁴I.B. Bernstein and F. Engelmann, *Phys.Fluids* 9, 937 (1966).
- ⁵R.L. Morse and C.W. Nielson, *Phys.Rev.Letters* 23, 1087 (1969).
- ⁶G. Strang and G.J. Fix, *An Analysis of the Finite Element Method* (Prentice-Hall, Engelwood Cliffs, 1973).
- ⁷K. Appert, T.M. Tran, and J. Vaclavik, *Computer Phys.Commun.* (to be published).
- ⁸A.A. Vedenov and L.I. Rudakov, *Dokl.Akad.Nauk SSSR* 159, 767 (1964) [*Sov.Phys.Dokl.* 9, 1073 (1965)].
- ⁹K. Appert, *Z.Angew.Math.Phys.* 26, 663 (1975).
- ¹⁰D.F. Smith (private communication)
- ¹¹A.A. Ivanov, T.K. Sobolyeva, and P.N. Yushmanov, *Zh.Eksp.Teor.Fiz.* 69, 2023 (1975).

FIGURE CAPTIONS

Figure 1: Time history of W , Δk_x and Δk_y . The initial conditions are $W_0 = 7 \times 10^{-8}$, $\xi = 10^{-5}$, and $u = 9$.

Figure 2: Saturation wave energy W_{sat} versus parameter ξ for different values of u . The solid lines and the dots refer to the 1-D and 2-D systems, respectively.

Figure 3: Spectral distribution of the field at several instants in time for the 2-D system at $k_y = 0$ (a) and for the 1-D system (b). The initial conditions are $\xi = .01$ and $u = 4.5$.

Figure 4: Two cross-sections of the velocity distribution function at several instants in time. The saturation time $t_{\text{sat}} = 1.5 \times 10^5$, and $W_0 = 7 \times 10^{-8}$.




Figure 1




Figure 2





Figure 3




Figure 4