

Non-perturbative Debye mass in finite T QCD

K. Kajantie^{a,b}, M. Laine^c, J. Peisa^d, A. Rajantie^b, K. Rummukainen^e and M. Shaposhnikov^a

^a*Theory Division, CERN, CH-1211 Geneva 23, Switzerland*

^b*Department of Physics, P.O.Box 9, 00014 University of Helsinki, Finland*

^c*Institut für Theoretische Physik, Philosophenweg 16, D-69120 Heidelberg, Germany*

^d*Department of Mathematical Sciences, University of Liverpool, Liverpool L69 3BX, UK*

^e*Fakultät für Physik, Postfach 100131, D-33501 Bielefeld, Germany*

(February 1, 2008)

Employing a non-perturbative gauge invariant definition of the Debye screening mass m_D in the effective field theory approach to finite T QCD, we use 3d lattice simulations to determine the leading $\mathcal{O}(g^2)$ and to estimate the next-to-leading $\mathcal{O}(g^3)$ corrections to m_D in the high temperature region. The $\mathcal{O}(g^2)$ correction is large and modifies qualitatively the standard power-counting hierarchy picture of correlation lengths in high temperature QCD.

PACS numbers: 11.10.Wx, 11.15.Ha, 12.38.Mh

QCD matter, a spatially and temporally extended system of matter described by the laws of Quantum Chromodynamics, goes at high temperatures into a quark-gluon plasma phase, in which color is no more confined and chiral symmetry is restored. An essential quantity, describing coherent static interactions in the plasma, is the inverse screening length of color electric fields, the Debye mass m_D . The Debye mass enters in many essential characteristics of static properties of the plasma. Its numerical value is important for phenomenological discussions of formation of the quark-gluon plasma, for the analysis of J/Ψ and Υ suppression in heavy ion collisions, for the computation of parton equilibration rates, etc. (see, e.g. [1]).

The definition and computation of the Debye mass for *abelian* QED plasma is well understood [2]. The electromagnetic current j_μ is a gauge-invariant quantity, and the Debye mass can be extracted from the 2-point gauge invariant correlation function of j_0 in the plasma. There are no massless charged particles in QED, which allows an infrared-safe perturbative computation of the Debye mass in powers of the electromagnetic coupling e . This has been done to order e^5 [3]. The situation in QCD is much more complicated. First, the corresponding current in QCD, j_μ^a , is not a gauge invariant quantity. Second, there are massless charged gluons which give rise to infrared divergences and prevent the perturbative determination of the Debye mass beyond leading order.

A non-perturbative gauge invariant definition of the Debye mass in vectorlike theories with zero chemical potential was suggested in [4]. According to it, m_D can be defined from the large distance exponential fall-off of correlators of gauge-invariant time-reflection odd operators O ,

$$\langle O(\tau, \vec{x}) O(\tau, 0) \rangle \sim C |\vec{x}|^\beta \exp(-m_D |\vec{x}|), \quad (1)$$

where C and β are some constants. The simplest choice

for the operator O is $F_{03}^a F_{12}^a$, and other examples can be found in [4]. In principle, 4-dimensional lattice simulations of hot QCD would thus allow a measurement of the Debye mass at any temperature.

The aim of this letter is a non-perturbative determination of the *high temperature limit* of the Debye mass, at $T > \text{a few} \times T_c$. We will see that the effective 3d approach to high temperature gauge theories, developed in [5–7] (for a review, see [8]) allows a simple and transparent gauge invariant definition of the Debye mass [4], while 3d lattice Monte Carlo simulations provide an economical way to determine its value. The corrections to the leading result we shall find are numerically large; thus many computations in the phenomenology of quark-gluon plasma in heavy ion collisions should be re-analysed.

The theory we shall study is QCD with N_f massless quark flavours and with the gauge group $SU(N)$ with $N = 2, 3$. At high temperatures and zero chemical potential the Debye mass can be expanded in a power series in the QCD coupling constant $g = g(\mu)$ (the scale μ will be specified later; the result for $N_f = 0$ is shown explicitly in eq. (5)):

$$m_D = m_D^{\text{LO}} + \frac{Ng^2T}{4\pi} \ln \frac{m_D^{\text{LO}}}{g^2T} + c_N g^2T + d_{N,N_f} g^3T + \mathcal{O}(g^4T). \quad (2)$$

The leading order (LO) perturbative result, $m_D^{\text{LO}} = (N/3 + N_f/6)^{1/2} gT$, has been known for a long time [9]. The logarithmic part of the $\mathcal{O}(g^2)$ correction can be extracted perturbatively [10], but c_N and the higher order corrections are non-perturbative. We are going to evaluate numerically the coefficients c_N and d_{N,N_f} .

Static Green's functions for bosonic fields of high temperature QCD at distances $|x| \gg T^{-1}$ we are interested in can be determined by constructing an effective 3d gauge theory, containing static magnetic gluons and the

zero component of the 4d gauge field, A_0 [5–7]. Moreover, a *super-renormalizable* 3d theory, defined by the Lagrangian

$$L_{\text{eff}}[A_i^a, A_0^a] = \frac{1}{4} F_{ij}^a F_{ij}^a + \text{Tr} [D_i, A_0][D_i, A_0] + m_3^2 \text{Tr} A_0^2 + \lambda_A (\text{Tr} A_0^2)^2, \quad (3)$$

gives the Green's functions to a relative accuracy $\mathcal{O}(g^4)$ [6], which is sufficient for the accuracy of the expansion in eq. (2). The parameters of the effective theory are related to the parameters of 4d QCD ($\Lambda_{\overline{\text{MS}}}, N, N_f$) and the temperature as described in [11]. For brevity, we give here the explicit expressions only for $N_f = 0$:

$$g_3^2 = g^2 (4\pi e^{-\gamma_E - \frac{1}{22}} T) T = \frac{24\pi^2 T}{11N \ln(6.742T/\Lambda_{\overline{\text{MS}}})}, \quad (4)$$

$$m_3^2 = \frac{N}{3} g^2 (4\pi e^{-\gamma_E - \frac{5}{22}} T) T^2, \quad (5)$$

$$\lambda_A = \frac{6+N}{24\pi^2} g^4 (4\pi e^{-\gamma_E - \frac{7}{44}} T) T. \quad (6)$$

Here $g(\mu)$ is the QCD coupling in the $\overline{\text{MS}}$ scheme and all the effective theory couplings have been computed including both the leading and the next-to-leading order contributions. The couplings (4)-(6) are independent of the gauge chosen for the perturbative computation. The expansion parameter is $g^2/16\pi^2$ so that the result should be accurate down to $T \approx$ a few $\times \Lambda_{\overline{\text{MS}}}$.

The dynamics of the 3d effective theory is fully characterised by the two dimensionless ratios

$$y = \frac{m_3^2}{g_3^4}, \quad x = \frac{\lambda_A}{g_3^2}, \quad (7)$$

and by the dimensionful coupling g_3^2 . The value of x is essentially fixed by T ,

$$x = \frac{6+N}{24\pi^2} g^2 (4\pi e^{-\gamma_E - 3/11} T) = \frac{6+N}{11N} \frac{1}{\ln(5.371T/\Lambda_{\overline{\text{MS}}})}, \quad (8)$$

while y and x , corresponding to physical 4d finite T QCD for $N_f = 0$, are related by:

$$y = y_{\text{dr}}(x) = \frac{2}{9\pi^2 x} + \frac{1}{4\pi^2} + \mathcal{O}(x), \quad N = 2 \quad (9)$$

$$= \frac{3}{8\pi^2 x} + \frac{9}{16\pi^2} + \mathcal{O}(x), \quad N = 3. \quad (10)$$

We are now ready to give a gauge-invariant definition of the Debye mass in the 3d language [4]. Physically, we want a local operator which makes A_0^a gauge invariant in the 3d theory and contains A_0^a singly. We can single out this state by a symmetry consideration. Note that the effective Lagrangian (3) has a discrete symmetry $A_0 \leftrightarrow -A_0$. Then the Debye mass can be defined as the mass of

the lightest 3d state which is odd under this symmetry. The operator of lowest dimension of this type is

$$h_i = \epsilon_{ijk} \text{Tr} A_0 F_{jk}. \quad (11)$$

Here the field A_0^a has been made gauge invariant by dressing it with a cloud of magnetic gluons.

At high T , one has $g \ll 1$ and, according to eqs. (4), (5), $m_3 \gg g_3^2$. This is the ‘‘heavy quark limit’’ of the 3d theory, in which the mass m_D of the singlet state is dominated by the bare mass m_3 of the scalar ‘‘quark’’ A_0^a . For dimensional reasons, the exact mass m_D can in this limit be expanded as

$$m_D = m_3 + a_N g_3^2 + \frac{b_N g_3^4}{m_3} + \mathcal{O}(\lambda_A, \frac{g_3^2 \lambda_A}{m_3}, \frac{g_3^6}{m_3^2}, \dots), \quad (12)$$

where a_N and b_N are constants, perhaps involving a logarithm of m_3/g_3^2 . The terms neglected are of higher order using the power counting in eqs. (4)-(6). Comparing eqs. (2) and (12), one sees that

$$a_N = \frac{N}{4\pi} \ln \frac{\sqrt{N/3 + N_f/6}}{g} + c_N, \quad d_{N, N_f} = \frac{b_N}{\sqrt{N/3 + N_f/6}}. \quad (13)$$

Here we used the fact that the scale dependence of the non-perturbative terms in eq. (2) is at least of order $\mathcal{O}(g^4)$. Since the expansion (12) refers only to the 3d theory, the constants a_N and b_N depend on N but clearly not on N_f . Thus c_N is N_f independent, while d_{N, N_f} depends on N_f only through m_D^{LO} .

In terms of our dimensionless variables (7), eq. (12) becomes

$$\frac{m_D}{g_3^2} = \sqrt{y} + \frac{N}{4\pi} \ln \sqrt{y} + c_N + \frac{b_N}{\sqrt{y}} + \dots \quad (14)$$

The mass m_D can now be measured by putting the effective 3d theory on the lattice [11] and by measuring the exponential falloff of the correlator $\langle h_3(x_3) h_3(0) \rangle \sim \exp(-m_D |x_3|)$, where $h_3(x_3)$ is summed over the transverse (x_1, x_2) plane. The correlation function is measured both with zero and finite transverse momentum, and in order to enhance the overlap with the asymptotic state the measurements are performed with several levels of recursive blocking of the operators. We select the blocking level and momentum sector which has the best signal for the asymptotic mass separately for each Monte Carlo run. Since the longest correlation length in each case is less than 1/5 of the linear size of the lattice, we expect the finite volume effects to be negligible in comparison with the statistical errors. This was also explicitly checked by performing simulations with different volumes in isolated cases.




FIG. 1. The gauge invariant Debye mass for SU(3), as a function of x , or $T/\Lambda_{\overline{\text{MS}}}$ through eq. (8). The scale of g on the y -axis is fixed according to eq. (5). The dashed line marks the leading order m_D^{LO} , and the continuous line the 2-parameter fit to eq. (12) with the parameters as in eq. (15).

The mass m_D is defined in the whole y, x parameter space. To have results which are relevant for 4d physics, we perform the measurements along the 2-loop dimensional reduction lines $y_{\text{dr}}(x)$, eqs. (9–10). To measure the coefficients in eq. (14) one should use the part of this curve corresponding to $\sqrt{y} \gg 1$. The results for $N = 3$ are shown in Fig. 1, in units of 4d gT ($= g_3^2 \sqrt{3y/N}$ in 3d units). The Monte Carlo runs are performed with several lattice spacings a , parametrised by $\beta_G \equiv 2N/(g_3^2 a)$. For SU(3) β_G varies by more than an order of magnitude (although not at the same value of x), as shown in Fig. 1; for SU(2), the measurements are done with $\beta_G = 20$ and 32. The top scale of Fig. 1 shows the physical temperature $T/\Lambda_{\overline{\text{MS}}}$ along $y_{\text{dr}}(x)$ -line. Note that the highest temperatures are larger than $10^{100} \times \Lambda_{\overline{\text{MS}}} \sim 10^{100} \times T_c$.

At small x (large y), the fit to the function (12) is very good, as indicated by the continuous line in Fig. 1. In order to see in detail the sensitivity of the fit to the parameters, in Fig. 2 we replot the SU(3) data (restricted to $x < 0.05$) in terms of the quantity $\delta m/g_3^2 = m_D/g_3^2 - \sqrt{y} - \frac{N}{4\pi} \ln \sqrt{y}$ as a function of $1/\sqrt{y}$. The intersection of the curve with the vertical axis gives the value of c_N and the slope gives $b_N = d_{N,N_f} \sqrt{N/3 + N_f/6}$. One can see that the linear fit is rather good even down to small values of \sqrt{y} . The large non-zero value of the intercept is very robustly determined. The slope b_N is small and has a relatively large error. Only the statistical error is given, but the value of b_N also depends on the range of




FIG. 2. The corrections of order $\mathcal{O}(g_3^2)$ and $\mathcal{O}(g_3^2/\sqrt{y})$ to the SU(3) screening mass m_D in 3d units, corresponding to eq. (14).

$y^{-1/2}$ included.

The result of the fits are

$$\begin{aligned} \text{SU(2): } c_N &= 1.58 \pm 0.20 & b_N &= -0.03 \pm 0.25 \\ \text{SU(3): } c_N &= 2.46 \pm 0.15 & b_N &= -0.49 \pm 0.15. \end{aligned} \quad (15)$$

The large number c_N is related to non-perturbative 3d effects, while the smaller d_{N,N_f} can be viewed as being related to the choice of scale in m_D^{LO} . For $N = 2$ we can in practice only verify that d_{2,N_f} is close to zero. Note that writing $c_N = N\tilde{c}_N$, one has $\tilde{c}_N = 0.79 \pm 0.10$ ($N = 2$), 0.82 ± 0.05 ($N = 3$).

One can observe the following:

- The leading term is dominant only at extremely large T . For SU(3), the leading term is larger than the $\mathcal{O}(g^2)$ correction for $g < 1/2.46$ or for $T/\Lambda_{\overline{\text{MS}}} \gtrsim \exp(8\pi^2 2.46^2/11) \gtrsim 10^{19}$. This implies that the leading term only dominates when QCD anyway merges into a unified theory.
- The four terms in eq. (14) fit the data over all the range $T \gtrsim 100T_c$ rather well, and there is no need for further corrections.
- In the range $\Lambda_{\overline{\text{MS}}} \approx T_c \lesssim T \lesssim 100T_c$, m_D is rather constant and $\approx 3.0 m_D^{\text{LO}}$ for SU(2) and $\approx 3.3 m_D^{\text{LO}}$ for SU(3). It should be noted, though, that in this regime $m_D^{\text{LO}} > T$ so that the hierarchy $m_D^2/(2\pi T)^2 \ll 1$ required for an accurate description of 4d physics through a 3d effective theory is getting weaker.

- The mass measured from the $\langle A_0 A_0 \rangle$ -correlator in the Landau gauge in 4d simulations for $N_f = 0$ has also been observed to be clearly larger than the leading term [12].
- If the mass $\sim gT$ of the A_0^a field is “large”, larger than the non-perturbative $\mathcal{O}(g^2)T$ correction, the A_0^a field can be further perturbatively integrated out and a simpler effective theory, containing only A_i^a (and possible scalar fields) can be derived [5–7]. Our results imply that this can be accurately carried out for QCD only at extremely high temperatures, $T \gg T_c$. In the electroweak case the accuracy of the integration is sufficient even for $T \sim T_c$ both since the leading term has a bigger coefficient ($m_D^{\text{LO}} = \sqrt{11/6}gT$) than the $N_f = 0$ QCD considered here and since in the relevant T regime the coupling constant $g = g(m_W) \approx 2/3$ is smaller.
- The usual parametric “power counting” picture of correlation lengths in high temperature QCD says that the longest scale, related to the magnetic sector of the theory, is $m_M^{-1} \sim (\text{const} \times g^2 T)^{-1}$. A shorter scale, $\sim (gT)^{-1}$, is associated with Debye screening. Our results show that this picture can be quantitatively correct only at extremely large temperatures. Indeed, purely magnetic effects, as measured by the 3d glueball (operator $F_{ij}^a F_{ij}^a$) mass ($m_G \approx 2g_3^2$ for pure SU(2) [13,11]) tend to be numerically large, so that $m_M \sim m_D$ in a very wide range of temperatures (This gauge invariant result is in contrast to the small magnetic gluon masses measured in Landau gauge [12]). In this range the longest length scale corresponds to a scalar 0^{++} 3d “bound state” of two A_0 quanta, associated with the operator $A_0^a A_0^a$ (the power counting suggestion that this state is roughly twice as heavy as m_D holds again only at extremely high T).

Summarizing, we have carried out with lattice Monte Carlo techniques a gauge independent measurement of the Debye mass in finite T QCD. The measurement is based on first deriving with 2-loop perturbative computations a 3d effective theory. The expansion parameter is α_s/π , so that the result is accurate down to T close to T_c . The mass is obtained by measuring correlators of the gauge invariant local operator $A_0^a F_{jk}^a$ in the 3d theory. The leading and next-to-leading corrections to m_D were determined and found to be large. In fact, for temperatures from T_c up to $T \sim 1000\Lambda_{\overline{\text{MS}}}$ the non-perturbative Debye screening mass is about a factor 3 larger than the lowest order estimate.

It remains to be seen whether this modification of the standard picture of high-temperature gauge theories has applications in the cosmological discussion of the quark-hadron phase transition or in the phenomenology of heavy ion collisions.

The simulations were carried out with a Cray C94 and Cray T3E at the Finnish Center for Scientific Computing.

-
- [1] F. Karsch, M. Mehr and H. Satz, Z. Phys. C **37**, 617 (1988); X.-N. Wang, Phys. Rep. **280**, 287 (1997).
 - [2] I. Akhiezer and S. Peletminski, Zh. Eksp. Teor. Fiz. **38**, 1829 (1960).
 - [3] J.-P. Blaizot, E. Iancu and R. Parwani, Phys. Rev. D **52**, 2543 (1995).
 - [4] P. Arnold and L.G. Yaffe, Phys. Rev. D **52**, 7208 (1995).
 - [5] P. Ginsparg, Nucl. Phys. B **170**, 388 (1980); T. Appelquist and R. Pisarski, Phys. Rev. D **23**, 2305 (1981); S. Nadkarni, Phys. Rev. D **27**, 917 (1983).
 - [6] K. Kajantie, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. B **407**, 356 (1993); K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. B **458**, 90 (1996).
 - [7] E. Braaten and A. Nieto, Phys. Rev. D **51**, 6990 (1995); Phys. Rev. D **53**, 3421 (1996).
 - [8] M.E. Shaposhnikov, Proceedings of the Summer School on Effective Theories and Fundamental Interactions, Erice, 1996 [CERN-TH-96-280, hep-ph/9610247].
 - [9] E. Shuryak, Zh. Eksp. Teor. Fiz. **74**, 408 (1978); J. Kapusta, Nucl. Phys. B **148**, 461 (1979); D. Gross, R. Pisarski and L. Yaffe, Rev. Mod. Phys. **53**, 43 (1981).
 - [10] A.K. Rebhan, Phys. Rev. D **48**, R3967 (1993); Nucl. Phys. B **430**, 319 (1994).
 - [11] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. B, in press [hep-ph/9704416].
 - [12] U. Heller, F. Karsch and J. Rank, Proceedings of Lattice '97, Nucl. Phys. B (Proc. Suppl.), to be published.
 - [13] O. Philipsen, M. Teper and H. Wittig, Nucl. Phys. B **469**, 445 (1996).