



Effect of fermionic components on trion–electron scattering

M. Combescot^{a,*}, O. Betbeder-Matibet^a, M.A. Dupertuis^b

^a Institut des NanoSciences de Paris, Université Pierre et Marie Curie, CNRS, 140 rue de Lourmel, 75015 Paris, France

^b Laboratoire d'Optoélectronique Quantique, Ecole Polytechnique Fédérale de Lausanne, Station 3, CH-1015 Lausanne, Switzerland

ARTICLE INFO

Article history:

Received 26 June 2008

Accepted 30 June 2008 by C. Tejedor

Available online 3 July 2008

PACS:

71.35.-y

Keywords:

A. Semiconductors

D. Trion

ABSTRACT

To test the validity of replacing a composite fermion by an elementary fermion, we calculate the transition rate from a state made of one free electron and one trion to a similar electron–trion pair, through the time evolution of such a pair induced by Coulomb interaction between elementary fermions. It is convenient to describe trion as one electron interacting with one exciton. This allows us to use the tools we have developed in the new composite-exciton many-body theory. The trion–electron scattering contains a direct channel in which “in” and “out” trions are made with the same fermions, and an exchange channel in which the “in” free electron becomes one of the “out” trion components. As expected, momenta are conserved in these two channels. The direct scattering is found to read as the bare Coulomb potential between elementary particles multiplied by a form factor which depends on the “in” and “out” trion relative motion indices η and η' , this factor reducing to $\delta_{\eta\eta'}$ in the zero momentum transfer limit. In this direct channel, the trion at large distance reacts as an elementary particle, its composite nature showing up at large momentum transfer. In contrast, the fact that the trion is not elementary does affect the exchange channel for all momentum transfers. We thus conclude that a 3-component fermion behaves as an elementary fermion for direct processes in the small momentum transfer limit only.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

The proper description of composite quantum particles has been a long-standing problem for decades. The simplest idea by far is to replace them by elementary particles, these particles being fermionlike if the number of fermions they contain is odd and bosonlike if this number is even.

A few years ago, we have reconsidered the problem of quantum-particle compositeness through the simplest case: just two fermions. Through a new many-body procedure [1,2] which allows to treat Pauli exclusion between fermionic components of these composite bosons exactly, we have shown that, in all physical effects we have studied up to now [2], the replacement of Wannier excitons by elementary bosons with effective interactions dressed by exchanges (as usually done), misses terms which can even be dominant in problems dealing with unabsorbed photons [3]. A way to grasp the difficulty is to note that replacement of a free electron–hole pair by an elementary boson strongly reduces the degrees of freedom of the system. This is beautifully seen through the prefactor change from $(1/N!)$ to $(1/N!)^2$ in the closure relation of elementary and composite bosons [4], making all sum rules for elementary and composite bosons irretrievably different,

whatever the effective scatterings generated by bosonization procedures are.

Composite bosons made of two fermions now are under good control, the subtle many-body physics of these systems resulting from fermion exchanges being nicely visualized through the so-called “Shiva diagrams” [5]. This is why it is now time to start tackling fermionlike composite particles. In this very first paper, we study the simplest problem: one trion made of three different fermions – to avoid complication linked to fermion exchange inside the particle itself. Such a 3-fermion particle can be deuterium atom made of one electron, one proton and one neutron. Other possibilities are H^- ion made of two opposite-spin electrons and one proton, or X^- semiconductor trion [6–13] in which proton is replaced by valence hole. While deuterium is neutral, both H^- ion and X^- semiconductor trion are negatively charged. Consequently, the scattering of such a composite fermion with a free electron is directly related to the way charge compositeness affects Coulomb interaction. We *a priori* expect this scattering to be the bare Coulomb potential V_Q between two elementary charges, with a form factor f_Q which comes from the trion composite nature. Since at large distance, trion should appear as one elementary negative charge, this form factor should reduce to 1 in the small Q limit. Closer, the fact that the trion is made of two electrons and one hole should show up through a form factor which differs from 1 when Q increases.

The purpose of this communication is to study the effect of compositeness of fermion-like particles through the precise

* Corresponding author.

E-mail address: Monique.Combescot@insp.jussieu.fr (M. Combescot).

calculation of the scattering of one electron with one semiconductor trion made of two opposite-spin electrons and one hole. We derive this scattering from the time evolution of electron–trion pair state induced by Coulomb interaction. Due to the quantum nature of trion components, it appears that this scattering contains a direct and an exchange channel. While the direct scattering tends to the scattering of elementary particles for small momentum transfer, the exchange channel leads to a scattering which has a totally different structure. A way to physically grasp this difference, is to say that a trion must behave as an elementary particle in processes in which its three fermions stay far apart from the free electron with which it interacts, as in direct processes with small momentum transfer. In contrast, trion and free electron are, by construction, not far apart when they exchange their fermions. This is why the composite-fermion nature of the trion must show up for all exchange processes.

2. Procedure

We consider a state made of one conduction electron \mathbf{K}_e with momentum \mathbf{k}_e , spin $\sigma = (\pm 1/2)$, and one trion J made of one valence hole and two opposite-spin conduction electrons, with center-of-mass momentum \mathbf{k}_j , relative motion index η_j and total electron spin ($S_j = (0, 1), S_{jz} = 0$). Let $a_{\mathbf{k}_e}^\dagger$ be the electron creation operator and T_j^\dagger the trion creation operator. The time evolution of this electron–trion pair state, due to Coulomb interaction included in the system Hamiltonian H , is given by $|\psi_t(\mathbf{K}_e, J)\rangle = \exp(-iHt)|\psi(\mathbf{K}_e, J)\rangle$, with $|\psi(\mathbf{K}_e, J)\rangle = a_{\mathbf{k}_e}^\dagger T_j^\dagger |v\rangle$. By using the integral representation of the exponential, this state also reads

$$|\psi_t(\mathbf{K}_e, J)\rangle = \int_{-\infty}^{+\infty} \frac{dx}{(-2i\pi)} \frac{e^{-i(x+i0_+)t}}{x+i0_+ - H} a_{\mathbf{k}_e}^\dagger T_j^\dagger |v\rangle, \quad (1)$$

which is valid for $t > 0$, provided that 0_+ is a positive constant.

To calculate this quantity in a convenient way, we introduce the electron creation potential [14] defined as $V_{\mathbf{k}_e}^\dagger = Ha_{\mathbf{k}_e}^\dagger - a_{\mathbf{k}_e}^\dagger(H + \epsilon_{\mathbf{k}_e}^{(e)})$. This operator describes all interactions of electron \mathbf{K}_e with the rest of the system. This allows us to write the key equation [14] for correlation effects with electron \mathbf{K}_e , namely,

$$\frac{1}{z - H} a_{\mathbf{k}_e}^\dagger = \left(a_{\mathbf{k}_e}^\dagger + \frac{1}{z - H} V_{\mathbf{k}_e}^\dagger \right) \frac{1}{z - H - \epsilon_{\mathbf{k}_e}^{(e)}}, \quad (2)$$

valid for any z . By inserting this equation into Eq. (1) and by noting that $(H - \epsilon_{\mathbf{k}_e}^{(e)})T_j^\dagger |v\rangle = 0$, the state $|\psi_t(\mathbf{K}_e, J)\rangle$ splits as

$$|\psi_t(\mathbf{K}_e, J)\rangle = e^{-i(\epsilon_{\mathbf{k}_e}^{(e)} + \epsilon_j^{(T)})t} a_{\mathbf{k}_e}^\dagger T_j^\dagger |v\rangle + |\tilde{\psi}_t(\mathbf{K}_e, J)\rangle, \quad (3)$$

where the state change is given by

$$|\tilde{\psi}_t(\mathbf{K}_e, J)\rangle = \int_{-\infty}^{+\infty} \frac{dx}{(-2i\pi)} \frac{e^{-i(x+i0_+)t}}{(x+i0_+ - H)(x+i0_+ - \epsilon_{\mathbf{k}_e}^{(e)} - \epsilon_j^{(T)})} V_{\mathbf{k}_e}^\dagger T_j^\dagger |v\rangle. \quad (4)$$

The transition rate towards another electron–trion state (\mathbf{K}'_e, J') must be identified with [15,16]

$$\frac{t}{\mathcal{T}_{(\mathbf{K}_e, J) \rightarrow (\mathbf{K}'_e, J')}} = \left| \langle \psi(\mathbf{K}'_e, J') | \tilde{\psi}_t(\mathbf{K}_e, J) \rangle \right|^2, \quad (5)$$

in order for the RHS of this equation to cancel with t , the state change reducing to 0 for $t = 0$, as readily seen from Eq. (3). Since $|\tilde{\psi}_t\rangle$ is first order in the interactions, due to the creation potential $V_{\mathbf{k}_e}^\dagger$ in Eq. (4), we find from $a_{\mathbf{k}_e}^\dagger(z - H)^{-1}$ deduced from

Eq. (2), that the scalar product in Eq. (5) reduces, at first order in the interactions, to

$$\langle \psi(\mathbf{K}'_e, J') | \tilde{\psi}_t(\mathbf{K}_e, J) \rangle \simeq \langle v | T_{J'} a_{\mathbf{k}'_e} V_{\mathbf{k}_e}^\dagger T_J^\dagger | F_t(\mathbf{K}'_e, J'; \mathbf{K}_e, J) \rangle, \\ F_t(\mathbf{K}'_e, J'; \mathbf{K}_e, J) = \int_{-\infty}^{+\infty} \frac{dx}{(-2i\pi)} \frac{e^{-i(x+i0_+)t}}{(x+i0_+ - \epsilon_{\mathbf{k}'_e}^{(e)} - \epsilon_j^{(T)})(x+i0_+ - \epsilon_{\mathbf{k}_e}^{(e)} - \epsilon_j^{(T)})}. \quad (6)$$

The t part $F_t(\mathbf{K}'_e, J'; \mathbf{K}_e, J)$ readily gives $-2i\pi e^{-it\Delta_+/2} \delta_t(\Delta_-)$, where $\Delta_\pm = \epsilon_{\mathbf{k}_e}^{(e)} + \epsilon_j^{(T)} \pm (\epsilon_{\mathbf{k}'_e}^{(e)} + \epsilon_j^{(T)})$, while $\delta_t(\Delta) = \sin(t\Delta/2)/\pi\Delta$ is the usual delta function of width $t/2$. Since $\delta_t(0) = t/2\pi$, the transition rate from state (\mathbf{K}_e, J) to state (\mathbf{K}'_e, J') then takes the physically expected form

$$\frac{1}{\mathcal{T}_{(\mathbf{K}_e, J) \rightarrow (\mathbf{K}'_e, J')}} = 2\pi \delta_t(\epsilon_{\mathbf{k}_e}^{(e)} + \epsilon_j^{(T)} - \epsilon_{\mathbf{k}'_e}^{(e)} - \epsilon_j^{(T)}) \times \left| \langle v | T_{J'} a_{\mathbf{k}'_e} V_{\mathbf{k}_e}^\dagger T_J^\dagger \rangle \right|^2. \quad (7)$$

3. Calculation of the transition rate

To calculate the matrix element appearing in this transition rate, we first need to determine the creation potential $V_{\mathbf{k}_e}^\dagger$. By writing the system Hamiltonian in second quantization as $H = H_e + H_h + V_{ee} + V_{hh} + V_{eh}$, this operator reduces to $[V_{ee} + V_{eh}, a_{\mathbf{k}_e}^\dagger]$. For $\mathbf{K}_e = (\mathbf{k}_e, \sigma)$, it reads

$$V_{\mathbf{k}_e}^\dagger = \sum_{\mathbf{q}} V_{\mathbf{q}} a_{\mathbf{k}_e + \mathbf{q}, \sigma}^\dagger \left(\sum_{\mathbf{p}, s} a_{\mathbf{p} - \mathbf{q}, s}^\dagger a_{\mathbf{p}, s} - \sum_{\mathbf{p}, m} b_{\mathbf{p} - \mathbf{q}, m}^\dagger b_{\mathbf{p}, m} \right), \quad (8)$$

where $b_{\mathbf{p}, m}^\dagger$ creates hole with momentum \mathbf{p} and “spin” $m = (\pm 3/2, \pm 1/2)$ or $m = (\pm 3/2)$ for bulk or quantum well samples.

We have shown [17] that the creation operator for trion made of electrons with spins (s, s') can be written in terms of electron–exciton pairs, as

$$T_j^\dagger = \sum_{\mathbf{v}, \mathbf{p}} \langle v, \mathbf{p} | \eta_j, S_j \rangle a_{\mathbf{p} + \beta_e \mathbf{k}_j, s}^\dagger B_{\mathbf{v}, -\mathbf{p} + \beta_x \mathbf{k}_j, s'}^\dagger, \quad (9)$$

where $\beta_e = 1 - \beta_x = m_e/(2m_e + m_h)$. Operator $B_{\mathbf{v}, \mathbf{Q}, s}^\dagger$ creates an exciton with center-of-mass momentum \mathbf{Q} , relative motion index \mathbf{v} , and electron spin $s = \pm 1/2$. The hole spin m being unimportant here, since we have one hole only, we can forget it to simplify notations. This exciton creation operator reads in terms of electron–hole pairs as

$$B_{\mathbf{v}, \mathbf{Q}, s}^\dagger = \sum_{\mathbf{p}} \langle \mathbf{p} | v \rangle a_{\mathbf{p} + \alpha_e \mathbf{Q}, s}^\dagger b_{-\mathbf{p} + \alpha_h \mathbf{Q}}^\dagger, \quad (10)$$

where $\alpha_e = 1 - \alpha_h = m_e/(m_e + m_h)$ while $\langle \mathbf{p} | v \rangle$ is the Fourier transform of the exciton relative motion wave function $(\mathbf{r} | v)$.

By comparing the two above equations, we note that the trion center-of-mass momentum \mathbf{k}_j splits between electron and exciton according to their masses, just as the exciton does. In the same way, the prefactor $\langle v, \mathbf{p} | \eta, S \rangle$ in the trion expansion (9) is the “Fourier transform in the exciton sense” of the trion relative motion wave function, as shown in previous works [17,18]. Let us briefly recall a few important points for the trion physics we have obtained in these works.

The physically relevant spatial variables for trions, i.e. the variables which fulfill $[\mathbf{r}_n, \mathbf{p}_n] = i\delta_{nn'}$ are $(\mathbf{R}, \mathbf{r}, \mathbf{u})$ or $(\mathbf{R}, \mathbf{r}', \mathbf{u}')$, where $\mathbf{R} = [m_e(\mathbf{r}_e + \mathbf{r}_{e'}) + m_h \mathbf{r}_h]/(2m_e + m_h)$ is the center-of-mass coordinate, $\mathbf{r} = \mathbf{r}_e - \mathbf{r}_h$, and $\mathbf{u} = \mathbf{r}_{e'} - (m_e \mathbf{r}_e + m_h \mathbf{r}_h)/(m_e + m_h)$ is the

distance between electron e' and the center of mass of (e, h) . The other two variables $(\mathbf{r}', \mathbf{u}')$ read as (\mathbf{r}, \mathbf{u}) with $(\mathbf{r}_e, \mathbf{r}_{e'})$ exchanged [19]. Within these variables, the trion Hamiltonian appears as

$$H(\mathbf{r}_e, \mathbf{r}_{e'}, \mathbf{r}_h) = \frac{p_{\mathbf{R}}^2}{2M_T} + h(\mathbf{r}, \mathbf{u}), \quad (11)$$

where the trion relative motion part is such that $h(\mathbf{r}, \mathbf{u}) = h(\mathbf{r}', \mathbf{u}')$ with

$$h(\mathbf{r}, \mathbf{u}) = h_X(\mathbf{r}) + \frac{p_{\mathbf{u}}^2}{2\mu_T} + v(\mathbf{r}, \mathbf{u}). \quad (12)$$

$h_X(\mathbf{r}) = p_{\mathbf{r}}^2/2\mu_X - e^2/r$ is the exciton Hamiltonian with effective mass $\mu_X^{-1} = m_e^{-1} + m_h^{-1}$, while the trion effective mass is $\mu_T^{-1} = m_e^{-1} + (m_e + m_h)^{-1}$. The coupling $v(\mathbf{r}, \mathbf{u})$, which comes from interactions of electron e' with the (e, h) pair is given by

$$v(\mathbf{r}, \mathbf{u}) = \frac{e^2}{|\mathbf{r}_{e'} - \mathbf{r}_e|} - \frac{e^2}{|\mathbf{r}_{e'} - \mathbf{r}_h|} = \frac{e^2}{|\mathbf{u} - \alpha_h \mathbf{r}|} - \frac{e^2}{|\mathbf{u} + \alpha_e \mathbf{r}|}. \quad (13)$$

Since the trion Hamiltonian is such that $H(\mathbf{r}_e, \mathbf{r}_{e'}, \mathbf{r}_h) = H(\mathbf{r}_{e'}, \mathbf{r}_e, \mathbf{r}_h)$, the orbital eigenstates are even or odd with respect to $(\mathbf{r}_e \leftrightarrow \mathbf{r}_{e'})$ exchange; due to Pauli exclusion, the even ones are associated with electron singlet states $S = 0$ while the odd ones are associated with triplets $S = 1$, the molecular ground state having an even orbital wave function as usual. Within these trion variables, the orbital parity reads $\langle \mathbf{r}, \mathbf{u} | \eta, S \rangle = (-1)^S \langle \mathbf{r}', \mathbf{u}' | \eta, S \rangle$. This condition leads, for the Fourier transform in the exciton sense, to [18]

$$\begin{aligned} \langle v, \mathbf{p} | \eta, S \rangle &= \int d\mathbf{r} d\mathbf{u} \langle v | \mathbf{r} \rangle \langle \mathbf{p} | \mathbf{u} \rangle \langle \mathbf{r}, \mathbf{u} | \eta, S \rangle \\ &= (-1)^S \sum_{\mathbf{v}', \mathbf{p}'} \langle v | \mathbf{p}' + \alpha_e \mathbf{p} \rangle \langle \mathbf{p} + \alpha_e \mathbf{p}' | v' \rangle \langle v', \mathbf{p}' | \eta, S \rangle. \end{aligned} \quad (14)$$

It is then possible to show that expression (9) for trion creation operator also reads

$$\begin{aligned} T_J^\dagger &= \frac{1}{2} \sum_{\mathbf{v}, \mathbf{p}} \langle v, \mathbf{p} | \eta_j, S_j \rangle \left[a_{\mathbf{p}+\beta_e \mathbf{k}_j, s}^\dagger B_{v, -\mathbf{p}+\beta_X \mathbf{k}_j, s'} \right. \\ &\quad \left. - (-1)^{S_j} a_{\mathbf{p}+\beta_e \mathbf{k}_j, s'}^\dagger B_{v, -\mathbf{p}+\beta_X \mathbf{k}_j, s} \right]. \end{aligned} \quad (15)$$

This makes this operator readily creation of triplet or singlet state, depending if S_j is equal to 1 or 0. However, calculations performed with T_J^\dagger written as in Eq. (9) with $\langle v, \mathbf{p} | \eta, S \rangle$ fulfilling Eq. (14), turns out to be far simpler than the ones using Eq. (15). (Terms like the sum in Eq. (14) are generated by crossed scalar products when using Eq. (15)).

Eq. (9) allows us to rewrite $V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle$ as

$$\begin{aligned} V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle &= \sum_{\mathbf{v}, \mathbf{p}} \langle v, \mathbf{p} | \eta_j, S_j \rangle \left(\{V_{\mathbf{k}_e}^\dagger, a_{\mathbf{p}+\beta_e \mathbf{k}_j, s}^\dagger\} B_{v, -\mathbf{p}+\beta_X \mathbf{k}_j, s'}^\dagger \right. \\ &\quad \left. - a_{\mathbf{p}+\beta_e \mathbf{k}_j, s}^\dagger [V_{\mathbf{k}_e}^\dagger, B_{v, -\mathbf{p}+\beta_X \mathbf{k}_j, s'}^\dagger] \right) |v\rangle, \end{aligned} \quad (16)$$

where $\{F, G\}$ stands for the anticommutator $(FG + GF)$, while $[F, G]$ stands for the commutator $(FG - GF)$. Due to Eq. (8), the anticommutator reduces to $\sum_{\mathbf{q}} V_{\mathbf{q}} a_{\mathbf{q}+\mathbf{k}_e, \sigma}^\dagger a_{-\mathbf{q}+\mathbf{p}+\beta_e \mathbf{k}_j, s'}$, so that the first part of $V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle$ corresponds to direct Coulomb process between free electron \mathbf{K}_e and the electron of the electron-exciton pair making trion J (see Fig. 1(a)). Similarly, the second part of $V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle$ corresponds to interactions with the exciton of this pair, as seen from the commutator which reduces to $\sum_{\mathbf{q}, v'} V_{\mathbf{q}} \gamma_{-q}(v', v) a_{\mathbf{q}+\mathbf{k}_e, \sigma}^\dagger B_{v', -\mathbf{q}-\mathbf{p}+\beta_X \mathbf{k}_j, s'}^\dagger$ (see Fig. 1(b)). The electron-exciton scattering amplitude, easy to obtain by expanding




Fig. 1. (a), (b) In order to have free electron (\mathbf{k}_e, σ) interacting with trion $(\mathbf{k}_j, \eta_j, S_j)$ in a convenient way, we first write trion in terms of electron-exciton pair, the vertex being the “Fourier transform in the exciton sense” $\langle v, \mathbf{p} | \eta_j, S_j \rangle$ of the trion relative motion wave function $|\eta_j, S_j\rangle$. In (a), the free electron interacts with the electron part of the trion, the coupling being $V_{\mathbf{q}} = V_{-\mathbf{q}}$, while in (b) the free electron interacts with the exciton part of the trion, the coupling being $V_{\mathbf{q}} \gamma_{-q}(v', v)$: In this interaction, the exciton goes from v to v' while its momentum change is $(-\mathbf{q})$. Exciton being neutral, $\gamma_{-q}(v', v)$ goes to zero when q goes to zero (see Eq. (17)). (c): “Out” state made of one free electron (\mathbf{k}_e', σ) and one trion $(\mathbf{k}_j, \eta_j, S_j)$ which results from the time evolution of the electron-trion pair state resulting from diagrams (a) and (b).

the exciton in electron-hole pairs according to Eq. (10), leads to [20]

$$\begin{aligned} \gamma_{\mathbf{q}}(v', v) &= \sum_{\mathbf{p}} (\langle v' | \mathbf{p} + \alpha_h \mathbf{q} \rangle - \langle v' | \mathbf{p} - \alpha_e \mathbf{q} \rangle) \langle \mathbf{p} | v \rangle \\ &= \langle v' | e^{i\alpha_h \mathbf{q} \cdot \mathbf{r}} - e^{-i\alpha_e \mathbf{q} \cdot \mathbf{r}} | v \rangle. \end{aligned} \quad (17)$$

This quantity, which also appears in the direct scattering of two excitons, is calculated in Ref. [20]. We find $\langle v_0 | e^{i\mathbf{q} \cdot \mathbf{r}} | v_0 \rangle = (1 + \tilde{q}^2/4)^{-2}$ or $(1 + \tilde{q}^2/16)^{-3/2}$, with $\tilde{q} = q a_X$, for 3D or 2D ground state excitons, i.e. for $|v_0\rangle$ states such that $\langle \mathbf{r} | v_0 \rangle = e^{-r/a_X} (a_X^{3/2} \sqrt{\pi})^{-1}$ or $\langle \mathbf{r} | v_0 \rangle = e^{-2r/a_X} 2^{3/2} (a_X \sqrt{\pi})^{-1}$.

All this leads to $V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle = \sum_{\mathbf{q}} V_{\mathbf{q}} a_{\mathbf{q}+\mathbf{k}_e, \sigma}^\dagger \mathcal{T}_{J, -\mathbf{q}}^\dagger |v\rangle$, where

$$\begin{aligned} \mathcal{T}_{J, \mathbf{q}}^\dagger &= \sum_{\mathbf{v}, \mathbf{p}} \left[a_{\mathbf{p}+\beta_e \mathbf{k}_j, s}^\dagger B_{v, -\mathbf{p}+\beta_X \mathbf{k}_j, s'}^\dagger \right. \\ &\quad \left. + a_{\mathbf{p}+\beta_e \mathbf{k}_j, s}^\dagger \sum_{v'} \gamma_{\mathbf{q}}(v', v) B_{v', \mathbf{q}-\mathbf{p}+\beta_X \mathbf{k}_j, s'}^\dagger \right] \langle v, \mathbf{p} | \eta_j, S_j \rangle. \end{aligned} \quad (18)$$

We now turn to the scalar product of $\langle v | T_J a_{\mathbf{k}_e}^\dagger$ (see Fig. 1(c)) with the two parts of $V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle$. For “out” electron $\mathbf{K}_e' = (\mathbf{k}_e', \sigma)$ with momentum \mathbf{k}_e' and same spin σ as the “in” electron, this scalar product splits into a direct and an exchange channel (see Fig. 2),

$$\langle v | T_J a_{\mathbf{k}_e'}^\dagger V_{\mathbf{k}_e}^\dagger T_J^\dagger |v\rangle = \xi^{\text{dir}} \begin{pmatrix} \mathbf{K}_e' & \mathbf{K}_e \\ J' & J \end{pmatrix} - \xi^{\text{ex}} \begin{pmatrix} \mathbf{K}_e' & \mathbf{K}_e \\ J' & J \end{pmatrix}. \quad (19)$$




Fig. 2. (a) Diagram for the time evolution of the “in” state made of one free electron (\mathbf{k}_e, σ) interacting with one trion (\mathbf{k}_j, η_j), due to their Coulomb interaction, the “out” state being electron (\mathbf{k}'_e, σ) and similar trion (\mathbf{k}'_j, η_j). (b): Direct Coulomb scattering between free electron and trion as given in Eq. (20). (c): “In” exchange Coulomb scattering between free electron and trion, as given in Eq. (21). In this scattering, interactions take place between “in” particles, “in” and “out” exchange scatterings being equal when energy is conserved (see Eq. (22)). (d): “In” exchange scattering between two free electrons.

In the direct channel, the “in” and “out” trions are made with the same fermions. Its precise value reads

$$\xi^{\text{dir}} \begin{pmatrix} \mathbf{K}'_e & \mathbf{K}_e \\ J' & J \end{pmatrix} = V_{\mathbf{K}'_e - \mathbf{k}_e} \langle v | T_{J'} \mathcal{T}_{J, \mathbf{k}_e - \mathbf{K}'_e}^\dagger | v \rangle. \quad (20)$$

In the exchange channel, the “in” electron becomes one component of the “out” trion. Its precise value reads

$$\xi^{\text{in}} \begin{pmatrix} \mathbf{K}'_e & \mathbf{K}_e \\ J' & J \end{pmatrix} = \sum_{\mathbf{q}} V_{\mathbf{q}} \langle v | T_{J'} a_{\mathbf{q} + \mathbf{k}_e, \sigma}^\dagger a_{\mathbf{K}'_e, \sigma}^\dagger \mathcal{T}_{J, -\mathbf{q}}^\dagger | v \rangle. \quad (21)$$

Note that, in this exchange scattering, Coulomb interactions take place between the “in” particles (\mathbf{K}_e, J). This makes the scattering rate defined in Eq. (7) not symmetrical with respect to “in” and “out” states, as reasonable since the state which evolves is the “in” state. It is, however, important to note that the “in” scattering obtained through $\langle v | T_{J'} a_{\mathbf{K}'_e}^\dagger V_{\mathbf{K}_e}^\dagger T_J^\dagger | v \rangle$ and the “out” scattering possibly obtained from $\langle v | T_{J'} V_{\mathbf{K}'_e} a_{\mathbf{K}_e}^\dagger T_J^\dagger | v \rangle$, are equal for energy conserving processes. Indeed, by calculating $\langle v | T_{J'} a_{\mathbf{K}'_e}^\dagger H a_{\mathbf{K}_e}^\dagger T_J^\dagger | v \rangle$ with H acting on the right side and on the left side, we find

$$\begin{aligned} & \langle v | T_{J'} V_{\mathbf{K}'_e} a_{\mathbf{K}_e}^\dagger T_J^\dagger | v \rangle - \langle v | T_{J'} a_{\mathbf{K}'_e}^\dagger V_{\mathbf{K}_e}^\dagger T_J^\dagger | v \rangle \\ &= \left(\epsilon_{\mathbf{K}_e}^{(e)} + \mathcal{E}_J^{(T)} - \epsilon_{\mathbf{K}'_e}^{(e)} - \mathcal{E}_{J'}^{(T)} \right) \langle v | T_{J'} a_{\mathbf{K}'_e}^\dagger a_{\mathbf{K}_e}^\dagger T_J^\dagger | v \rangle. \end{aligned} \quad (22)$$

Since states (\mathbf{K}_e, J) and (\mathbf{K}'_e, J') in the transition rate have the same energy in the large t limit (see Eq. (7)), the relevant scattering for transition rate can be calculated with Coulomb interactions acting either between “in” or between “out” states.

4. Direct channel

Let $\mathbf{Q} = \mathbf{K}'_e - \mathbf{k}_e$ be the momentum transfer of the electron–trion scattering of interest (see Fig. 2(a)). The scattering associated to

the direct channel given in Eq. (20) appears as the bare Coulomb scattering $V_{\mathbf{Q}}$ multiplied by a form factor $f_{\mathbf{Q}}(J', J)$ which is equal to $\langle v | T_{J'} \mathcal{T}_{J, -\mathbf{Q}}^\dagger | v \rangle$. The operator $\mathcal{T}_{J, -\mathbf{Q}}^\dagger$ contains two terms. The first one comes from free electron \mathbf{K}_e having Coulomb interaction with the electron part of the trion, while, in the second term, this interaction takes place with the exciton part of the trion. In the $Q \rightarrow 0$ limit, the first term of $\mathcal{T}_{J, -\mathbf{Q}}^\dagger$ tends to T_J^\dagger , while in the second term, $\gamma_{\mathbf{Q}}(v', v)$ goes to zero, as seen from Eq. (17): This physically comes from the fact that exciton is neutral, so that at large distance, i.e., at small Q , the effects of its two opposite charges cancel. Consequently, $\lim_{Q \rightarrow 0} \mathcal{T}_{J, -\mathbf{Q}}^\dagger = T_J^\dagger$. This readily shows that the form factor $f_{\mathbf{Q}}(J', J)$ reduces to $\delta_{J'J}$ for $Q \rightarrow 0$: In this limit, the direct scattering of one free electron and one trion thus tends to $V_{\mathbf{Q}} \delta_{J'J}$, as if the trion were an elementary particle J .

The composite nature of the trion shows up with $f_{\mathbf{Q}}(J', J)$ departing from $\delta_{J'J}$ when Q increases, i.e. at small distance, as physically reasonable. Its precise value reads

$$\begin{aligned} f_{\mathbf{Q}}(J', J) &= \langle v | T_{J'} \mathcal{T}_{J, -\mathbf{Q}}^\dagger | v \rangle \\ &= \delta_{\mathbf{k}_{J'}, \mathbf{k}_j - \mathbf{Q}} \sum_{v, v', \mathbf{p}} [\langle S_{J'} | \eta_j | \mathbf{p} - \beta_{\mathbf{x}} \mathbf{Q}, v' \rangle \delta_{v', v} \\ &\quad + \langle S_{J'} | \eta_j | \mathbf{p} + \beta_e \mathbf{Q}, v' \rangle \gamma_{-\mathbf{Q}}(v', v)] \langle v, \mathbf{p} | \eta_j, S_j \rangle. \end{aligned} \quad (23)$$

5. Exchange channel

We now turn to the exchange scattering given in Eq. (21). Using Eq. (9) for T_J and Eq. (18) for $\mathcal{T}_{J, \mathbf{q}}$, we find for trions made of opposite-spin electrons

$$\begin{aligned} \xi^{\text{in}} \begin{pmatrix} \mathbf{K}'_e & \mathbf{K}_e \\ J' & J \end{pmatrix} &= \delta_{\mathbf{K}'_e + \mathbf{k}_{J'}, \mathbf{K}_e + \mathbf{k}_j} \sum_{\mathbf{q}, v'} V_{\mathbf{q}} \langle S_{J'} | \eta_j | v', \mathbf{q} + \mathbf{k}_e - \beta_e \mathbf{k}_{J'} \rangle \\ &\quad \times \left[\langle v', \mathbf{q} + \mathbf{k}'_e - \beta_e \mathbf{k}_j | \eta_j, S_j \rangle \right. \\ &\quad \left. + \sum_v \gamma_{-\mathbf{q}}(v', v) \langle v, \mathbf{k}'_e - \beta_e \mathbf{k}_j | \eta_j, S_j \rangle \right]. \end{aligned} \quad (24)$$

It is of interest to note that, if trion J could be reduced to elementary electron with momentum \mathbf{k}_j , the corresponding exchange scattering shown in Fig. 2(d) would read $\delta_{\mathbf{K}'_e + \mathbf{k}_{J'}, \mathbf{K}_e + \mathbf{k}_j} V_{\mathbf{k}_j - \mathbf{k}_e}$. We see that the above result for the exchange scattering of one electron and one composite trion never reduces to the one for two elementary charges, even for zero momentum transfer, i.e., for $\mathbf{K}'_e = \mathbf{k}_e$, which is the limit in which the direct scatterings for composite and elementary trion are found to be the same.

The precise calculation of the form factor $f_{\mathbf{Q}}(J', J)$ for arbitrary momentum transfer \mathbf{Q} , as well as the “in” exchange scattering requires the knowledge of the trion Fourier transform in the exciton sense $\langle v, \mathbf{p} | \eta, S \rangle$, i.e. the knowledge of the trion relative motion wave function $\langle \mathbf{r}, \mathbf{u} | \eta, S \rangle$ (see Eq. (14)). While the trion ground state energy can be obtained from variational procedures through not too heavy numerical calculations based on minimizing $\langle \psi | H | \psi \rangle / \langle \psi | \psi \rangle$ [21–29], the derivation of trion wave function for bulk or quantum well samples, i.e., the resolution of the Schrödinger equation $(H - E) | \psi \rangle = 0$, with $\psi(\mathbf{r}_e, \mathbf{r}_{e'}, \mathbf{r}_h) = \psi(\mathbf{r}_{e'}, \mathbf{r}_e, \mathbf{r}_h)$ for 2D and 3D systems, is far more tricky: It is known that trial functions giving good energies can in fact be very far from the exact eigenfunctions. This is why it would be necessary to really face a more accurate solution of the Schrödinger equation in order to get reliable form for the trion wave function. Unfortunately, such a reliable form, realistic [30,31] but simple enough to be of possible use for the calculation of $f_{\mathbf{Q}}(J', J)$ and $\xi^{\text{in}} \begin{pmatrix} \mathbf{K}'_e & \mathbf{K}_e \\ J' & J \end{pmatrix}$, is not

yet available in the literature and is beyond the scope of this work. This is why the calculation of these quantities for any momentum transfer cannot be incorporated here. We have in mind, in a near future, to tackle the difficult problem of determining the trion wave function along the ideas we have here developed.

6. Conclusion

This work shows that a composite fermion made of two opposite-spin electrons and one hole behaves as an elementary fermion for direct process in the small momentum transfer limit only. For all other cases, in particular when fermion exchanges take place, the compositeness of the particle affects its scattering substantially. This work also shows that the representation of the trion as an electron interacting with a composite exciton is again quite convenient as it leads to very compact calculations, the “Fourier transform in the exciton sense” of the trion relative motion wave function appearing as the relevant quantity for interacting-trion problems.

References

- [1] For a short review, see M. Combescot, O. Betbeder-Matibet, Solid State Commun. 134 (2005) 11.
- [2] For a longer review, see M. Combescot, O. Betbeder-Matibet, F. Dubin, Phys. Rep. 463 (2008) 214.
- [3] M. Combescot, O. Betbeder-Matibet, K. Cho, H. Ajiki, Europhys. Lett. 72 (2005) 618.
- [4] M. Combescot, O. Betbeder-Matibet, Phys. Rev. B 72 (2005) 193105.
- [5] M. Combescot, O. Betbeder-Matibet, Eur. Phys. J. B 55 (2007) 63.
- [6] M. Lampert, Phys. Rev. Lett. 1 (1958) 450.
- [7] B. Stebé, E. Feddi, G. Munschy, Phys. Rev. B 35 (1987) 4331.
- [8] B. Stebé, A. Ainane, Superlatt. Microstruct. 5 (1989) 545.
- [9] K. Kheng, R.T. Cox, Y. Merle d’Aubigné, F. Bassani, K. Saminadayar, S. Tatarenko, Phys. Rev. Lett. 71 (1993) 1752.
- [10] G. Finkelstein, H. Shtrikman, I. Bar-Joseph, Phys. Rev. Lett. 74 (1995) 976.
- [11] A.J. Shields, J.L. Osborne, M.Y. Simmons, M. Pepper, D.A. Ritchie, Phys. Rev. B 52 (1995) R 5522.
- [12] A. Thiligam, Phys. Rev. B 55 (1997) 7804.
- [13] B. Stebé, E. Feddi, A. Ainane, F. Dujardin, Phys. Rev. B 58 (1998) 9926.
- [14] This creation potential as well as Eq. (2) for correlation effects are similar to the ones for excitons.
- [15] M. Combescot, O. Betbeder-Matibet, Phys. Rev. Lett. 93 (2004) 016403.
- [16] M. Combescot, O. Betbeder-Matibet, R. Combescot, Phys. Rev. B 75 (2007) 174305.
- [17] M. Combescot, Eur. Phys. J. B 33 (2003) 311.
- [18] M. Combescot, O. Betbeder-Matibet, F. Dubin, Eur. Phys. J. B 42 (2004) 63.
- [19] The trion variables commonly used in the literature have \mathbf{u} replaced $\mathbf{r}_{e'} - \mathbf{r}_h$. The corresponding set of variables as well as the variables $|\mathbf{r}_e - \mathbf{r}_{e'}|$, $|\mathbf{r}_e| + |\mathbf{r}_{e'}|$ and $|\mathbf{r}_e| - |\mathbf{r}_{e'}|$ introduced by Hylleraas (Phys. Rev. 71, 491 (1947)) do not fulfill the required commutators $[\mathbf{r}_n, \mathbf{p}_{n'}] = i\delta_{nn'}$. This prevents proper decoupling between relative motions.
- [20] O. Betbeder-Matibet, M. Combescot, Eur. Phys. J. B 27 (2002) 505.
- [21] G. Munschy, B. Stebé, Phys. Status Solidi B 63 (1974) 459; Phys. Status Solidi B 72 (1975) 113.
- [22] B. Stebé, G. Munschy, Solid State Commun. 17 (1975) 1051.
- [23] Z.A. Insepov, G.E. Norman, JETP 42 (1976) 674.
- [24] O.V. Volkov, M. Pohl, I.V. Kukushkin, W. Dietsche, K.v Klitzing, Europhys. Lett. 50 (2000) 409.
- [25] A.B. Dzyubenko, A. Yu Sivachenko, Phys. Rev. Lett. 84 (2000) 4429.
- [26] C. Riva, F.M. Peeters, K. Varga, Phys. Rev. B 61 (2000) 13873.
- [27] A. Esser, E. Runge, R. Zimmermann, W. Langbein, Phys. Rev. B 62 (2000) 8232.
- [28] A. Esser, R. Zimmermann, E. Runge, Phys. Status Solidi. 227 (2001) 317.
- [29] L.C. Dacal, R. Ferreira, G. Bastard, J.A. Brum, Phys. Rev. B 65 (2002) 115325.
- [30] R. Schilling, D.C. Mattis, Phys. Rev. Lett. 49 (1982) 808; Phys. Rev. B 27 (1983) 3318.
- [31] M. Shibata, Y. Kuramoto, T. Watanabe, Phys. Rev. B 42 (1990) 2432.