

Effect of manganese chloride on the neurochemical profile of the rat hypothalamus

Nathalie Just^{1,2}, Cristina Cudalbu¹, Hongxia Lei¹ and Rolf Gruetter^{1,2,3}

¹Laboratory for Functional and Metabolic Imaging, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland; ²Department of Radiology, University of Lausanne, Lausanne, Switzerland; ³Department of Radiology, University of Geneva, Geneva, Switzerland

Manganese (Mn²⁺)-enhanced magnetic resonance imaging studies of the neuronal pathways of the hypothalamus showed that information about the regulation of food intake and energy balance circulate through specific hypothalamic nuclei. The dehydration-induced anorexia (DIA) model demonstrated to be appropriate for studying the hypothalamus with Mn²⁺-enhanced magnetic resonance imaging. Manganese is involved in the normal functioning of a variety of physiological processes and is associated with enzymes contributing to neurotransmitter synthesis and metabolism. It also induces psychiatric and motor disturbances. The molecular mechanisms by which Mn²⁺ produces alterations of the hypothalamic physiological processes are not well understood. ¹H-magnetic resonance spectroscopy measurements of the rodent hypothalamus are challenging due to the distant location of the hypothalamus resulting in limited measurement sensitivity. The present study proposed to investigate the effects of Mn²⁺ on the neurochemical profile of the hypothalamus in normal, DIA, and overnight fasted female rats at 14.1 T. Results provide evidence that γ -aminobutyric acid has an essential role in the maintenance of energy homeostasis in the hypothalamus but is not condition specific. On the contrary, glutamine, glutamate, and taurine appear to respond more accurately to Mn²⁺ exposure. An increase in glutamine levels could also be a characteristic response of the hypothalamus to DIA.

Journal of Cerebral Blood Flow & Metabolism (2011) 31, 2324–2333; doi:10.1038/jcbfm.2011.92; published online 29 June 2011

Keywords: animal models; astrocytes; calcium channels; GABA; MR spectroscopy

Introduction

Manganese (Mn²⁺) is an essential metal necessary for the normal functioning of a variety of physiological processes: it has a role in immune system functioning, regulation of cellular energy, and is cofactor for a variety of enzymes involved in neurotransmitter synthesis and metabolism (Aschner *et al*, 2007; Benedetto *et al*, 2009). However, it is also a neurotoxicant and induces psychiatric and motor disturbances as well as cognitive deficits such as memory and information processing impairments, reduced learning capacity, cognitive slowing, and decreased mental flexibility (Aschner *et al*, 2007;

Benedetto *et al*, 2009). The molecular mechanisms by which Mn²⁺ produces alterations of the hypothalamic physiological processes are not well understood.

Recently, Mn²⁺-enhanced magnetic resonance imaging techniques demonstrated a growing potential for structural imaging (Silva *et al*, 2008) and the identification of neuronal pathways through tract tracing studies (Pautler, 2004). Manganese has the potential to enter excitable cells through voltage-gated calcium channels and acts as a marker of brain activation (Aoki *et al*, 2004). Previous Mn²⁺-enhanced magnetic resonance imaging studies established that several hypothalamic nuclei (the paraventricular nuclei of the hypothalamus (PVN), the lateral hypothalamus (LH), the ventromedial nucleus of the hypothalamus, and the arcuate nucleus) of rats and mice show specific enhancement patterns following systemic administration of MnCl₂ associated with food-related challenges (Parkinson *et al*, 2009; Chaudhri *et al*, 2006). These enhancements were related to neuronal activity in the aforementioned hypothalamic nuclei.

In relation to these novel and promising techniques, dosage and administration routes of Mn²⁺ have been investigated (Silva and Bock, 2008), since acute

Correspondence: Dr N Just, Laboratory for Functional and metabolic Imaging (LIFMET), Centre d'Imagerie Biomédicale (CIBM), CH-1015 Lausanne, Switzerland.

E-mail: nathalie.just@epfl.ch

This work was supported by the Centre d'Imagerie BioMédicale (CIBM) of The University of Lausanne (UNIL), the Ecole Polytechnique Fédérale de Lausanne (EPFL), of the Hôpital Universitaire de Genève (HUG), and the Centre Hospitalier Universitaire Vaudois (CHUV) and Leenards and Jeantet foundations.

Received 9 February 2011; revised 19 April 2011; accepted 3 June 2011; published online 29 June 2011

exposure to high doses of Mn^{2+} can lead to hepatic failure and cardiac toxicity (Silva and Bock, 2008; Silva *et al*, 2004). Only a few NMR (nuclear magnetic resonance) spectroscopic studies were conducted to investigate the influence of Mn^{2+} on brain metabolism (Zwingmann *et al*, 2003, 2004, 2007). In rat models, Mn^{2+} affected both astrocytic and neuronal integrity (Zwingmann *et al*, 2003, 2004, 2007). Moreover, Mn^{2+} neurotoxicity has been associated with changes in brain energy metabolism and stimulation of oxidative glucose metabolism as well as protective astrocytic response (Zwingmann *et al*, 2004). Manganese in brain is preferentially accumulated in astrocytes and is known to stimulate the production of reactive oxygen and nitrogen species resulting in oxidative/nitrosative stress (Zwingmann *et al*, 2004, 2007). Additionally, Mn^{2+} was shown to induce cell swelling (Rama Rao *et al*, 2010). On the contrary, a model of dehydration-induced anorexia (DIA) was shown to induce cell shrinkage due to hyperosmolar challenge in rats (Jaimes-Hoy *et al*, 2008; Salter-Venzon and Watts, 2008, 2009).

Another benefit of using experimental models of Mn^{2+} exposure could therefore be the possibility to investigate neuronal and astrocytic mechanisms of regulation of the metabolism.

In the present work, our aim was to investigate the effects of Mn^{2+} on the proton neurochemical profile of the hypothalamus under disrupted energy balance conditions and following homeostatic changes. The following questions were addressed:

- How does Mn^{2+} affect hypothalamic metabolism? What are the energetic effects of acute Mn^{2+} exposure?
- Can Mn^{2+} modify the metabolic response to a specific challenge? Can this modification be characterized?

The neurochemical profiles of the hypothalamus of healthy, dehydrated-induced anorexic (Jaimes-Hoy *et al*, 2008; Watts, 1999), and overnight food suppressed (OFS) rats were measured and examined with and without the influence of Mn^{2+} .

Materials and methods

Animal Preparation

All studies were performed following the approval of Service de la consommation et des affaires vétérinaires du canton de Vaud (Switzerland). Thirty female Wistar rats (225 to 250 g body weight on day 0 (D0)) were divided into three groups: the control group (CTL, $n=10$), the OFS group ($n=10$), and the DIA group ($n=10$). Magnetic resonance (MR) experiments were conducted on one rat per group per session. Animals were acclimatized in an animal facility under humidity and temperature-controlled conditions on a 12-hour light/dark cycle during 1 week before the beginning of the experiments. Animals were allowed *ad libitum* access to standard chow and drinking water during this period.

On day 0 (D0), animals were placed in individual cages. Animals from the CTL group and the OFS group received water and food *ad libitum*; the dehydrated group (DIA) received 2.5% NaCl (Sigma, Zofingen, Switzerland) as drinking solution (Jaimes-Hoy *et al*, 2008; Watts, 1999). On D3 at 1800 hours, the OFS rats had their chow removed until the MR scan took place, 24 hours later. The amounts of food and drink were measured at 1800 hours on D0, D1, D3, D7, D10, and D14. The rats' body weight was also measured. On D3, rats from each group were anesthetized with 2% isoflurane in a mixture of O_2 and N_2O via a facemask. The tail vein was catheterized for intravenous infusion of $MnCl_2$. Each rat received an intravenous infusion of 100 mg/kg at 100 mmol/L $MnCl_2$ at a rate of 0.5 mL to 1 mL/h via the tail vein to avoid cardiac arrest. Animals were all the time gently manipulated to avoid stress and suffering. During infusion and until animals were completely recovered from anesthesia, their temperature was carefully monitored and kept at $37^\circ C \pm 1^\circ C$.

Five female Wistar rats per group did not undergo $MnCl_2$ infusions and were scanned on D4 after the start of anorexic protocol (D0) for the DIA group of rats.

During the MR experiments, rats were anesthetized with 2% isoflurane in a mixture of O_2 and N_2O . The rats were positioned in a dedicated holder with head fixation. Their body temperature was maintained at $37.5^\circ C \pm 0.5^\circ C$ using a temperature-controlled water circulation and a rectal probe. Their respiration rate was monitored throughout the experiment. Magnetic resonance imaging and magnetic resonance spectroscopy (MRS) were performed on D4 24 hours after $MnCl_2$ infusion.

Proton Magnetic Resonance Spectroscopy in the Hypothalamus and Hippocampus

Localized proton spectroscopy was performed using Spin-Echo full intensity Acquired Localized (SPECIAL) (Mlynárik *et al*, 2006), using echo time/repetition time (TE/TR) = 2.8/4,000 milliseconds in combination with outer volume suppression and vapor water suppression (Tkác *et al*, 1999). A voxel of interest of 50 μL was localized in the hippocampus and the PVN-LH region for all the rats using the previously acquired 3D-gradient echo (GRE) images. All first- and second-order shims were readjusted using Fast Automatic Shimming by Mapping along Projections (FASTMAP) (Gruetter and Tkác, 2000) in the voxel of interest, which resulted in water linewidth of 23 ± 5 Hz. To obtain adequate signal-to-noise ratio, 640 scans were acquired. For absolute quantification, water signal was acquired using identical parameters, except without water suppression and averaging eight scans.

Quantification and Statistics

The *in vivo* 1H -MR spectra were processed using linear combination (LC) model (Provencher, 1993). In this study, 18 metabolite concentrations were quantified using databases of simulated spectra of metabolites (Mlynárik *et al*, 2006). Only metabolites with Cramer–Rao lower bounds under 20% were kept for further analysis.

All data are presented as mean \pm standard error of the mean (s.e.m.) unless otherwise stated. Two-tailed unpaired Student's *t*-tests were applied to compare data acquired in the three groups of rats. A *P* value <0.05 is considered significant.

Results

3D-gradient echo T_1 -weighted images obtained at 14.1 T were used to position carefully the voxels of interest in the hippocampus and the hypothalamus of female rats. 3D images provided an increased spatial resolution in the rat brain and voxels of interest were positioned with reference to the Paxinos atlas (Paxinos and Watson, 1998). Despite challenges due to the depth of the hypothalamus in the rat, good quality spectra were obtained using a surface radiofrequency (RF) coil at 14.1 T and relative to the hippocampus (see Supplementary file). Eleven

metabolites were quantified reliably in the hypothalamus with Cramer–Rao lower bounds under 20%. The comparison of ^1H -MRS spectra of the hypothalamus with spectra of the hippocampus in normal female rats demonstrated important differences such as lower Glu, Tau, *N*-acetyl-aspartate (NAA), PCr, PCr + Cr (tCr) concentrations in agreement with the previous findings in mice (Lei *et al*, 2010). A lower γ -aminobutyric acid (GABA) peak was also found in female rats. A comparison of ^1H -MRS spectra of the hypothalamus with spectra of the hippocampus in female rats subjected to dehydration by drinking a hypertonic saline solution demonstrated higher concentrations of most metabolites in both the hippocampus and the hypothalamus (relative differences in hypothalamus: $\Delta\text{Tau} = +106\%$, $\Delta\text{tCr} = +40\%$, $\Delta\text{myo-Ins} = +92\%$, $\Delta\text{GABA} = +160\%$; in hippocampus: $\Delta\text{Tau} = +30\%$, $\Delta\text{tCr} = +50\%$, $\Delta\text{myo-Ins} = +72\%$). Interestingly, GABA was lower in the hippocampus of DIA rats ($\Delta\text{GABA} = -50\%$).




Figure 1 ^1H -nuclear magnetic resonance (NMR) spectra obtained at 14.1 T of the rat hypothalamus in control (CTL), dehydrated-induced anorexic (DIA), and overnight food-restricted (OFS) rats with metabolite assignments. Peak assignments: Cr, creatine; GABA, γ -aminobutyric acid; Gln, glutamine; Glu, glutamate; Lac, lactate; myo-Ins, myo-inositol; NAA, *N*-acetyl-aspartate; Tau, taurine. The voxels of interest are overlaid on 3D- T_1 -weighted gradient echo images (TR/TE = 20/5 milliseconds, flip angle = 70° , field of view = $25 \times 25 \times 25 \text{ mm}^3$, matrix size = $256 \times 256 \times 128$, coronal slices, BW = 25 KHz, five averages). Spectra were also obtained 24 hours after an intravenous injection of 100 mg/kg MnCl_2 in CTL (MnCl_2), DIA (MnCl_2), and OFS (MnCl_2) rats. Colored arrows indicate increases or decreases of the metabolites of interest relative to the CTL spectrum: red for GABA and lactate; green for myo-Ins; and violet for Tau. The color reproduction of this figure is available at the *Journal of Cerebral Blood Flow and Metabolism* journal online. BW, bandwidth; TE, echo time; TR, repetition time.

^1H -MRS spectra are presented in Figure 1 for each condition CTL, DIA, and OFS with and without MnCl_2 . Twenty-four hours after infusion of MnCl_2 in the rat tail vein, CTL rats demonstrated a significant increase in GABA and Lac concentrations (t -test, $P=0.01$) compared with CTL rats with no MnCl_2 infusion (Figure 2A). MnCl_2 -infused DIA rats demonstrated no significant difference with non- MnCl_2 -infused DIA rats except for Tau (t -test, $P=0.01$) (Figure 2B). Under DIA condition, all metabolite concentrations increased relative to the CTL condition. However, no significant differences were detected between CTL and DIA rats in the presence of MnCl_2 . In particular, the relative difference for GABA between CTL and DIA was only 3.3%. To verify that changes seen between MnCl_2 and non- MnCl_2 -infused DIA rats were solely related to osmotic changes due to hypertonic saline in the rat hypothalamus, a third group of rats was introduced. These rats had their food suppressed overnight (OFS). Five of them received an infusion of MnCl_2 . Their hypothalamic neurochemical pro-

files 24 hours post- MnCl_2 infusion showed lower metabolite concentrations compared with OFS rats without MnCl_2 ($\Delta\text{Glu}=-28\%$, t -test, $P=0.03$; $\Delta\text{Tau}=-55\%$, t -test, $P=0.04$; ΔNAA , $\Delta\text{Cr}\sim-40\%$ but $P>0.05$) (Figure 2C). Glc was undetectable and GABA and myo-Ins levels increased, although not significantly, relative to OFS rats without MnCl_2 . In MnCl_2 -infused OFS rats, metabolite concentrations had a tendency to decrease relative to their CTL ($\Delta\text{GABA}=-6\%$, $\Delta\text{Tau}=-17\%$, $\Delta\text{Glu}=-18\%$, $\Delta\text{Lac}=-40\%$), whereas most metabolite concentrations increased significantly in OFS rats without MnCl_2 relative to their CTL. Figure 3 compares metabolite concentrations for CTL, DIA, and OFS rats (Figure 3A) and for CTL (+ Mn^{2+}), DIA (+ Mn^{2+}), and OFS (+ Mn^{2+}) (Figure 3B). While significant differences were found between the neurochemical profile of CTL and the neurochemical profiles of DIA and OFS, these differences appear to be masked in the presence of Mn^{2+} .

The relative concentration changes (percentage of CTL) are presented in Figure 4. Significant




Figure 2 Effect of manganese (Mn^{2+}) on the neurochemical profile of control (CTL), dehydration-induced anorexia (DIA), and overnight food suppressed (OFS) rats. (A) Comparison between absolute concentrations in CTL and CTL (+ MnCl_2). (B) Comparison between absolute concentrations in DIA and DIA (+ MnCl_2). DIA rats were examined by ^1H -nuclear magnetic resonance (NMR) spectroscopy 4 days after they started drinking a 2.5% NaCl solution (spontaneous drinking) and 24 hours postintravenous infusion of 100 mg/kg MnCl_2 . (C) Comparison between absolute concentrations in OFS and OFS (+ MnCl_2). Rats were fasted 24 hours before ^1H -NMR spectroscopy. * $P < 0.05$.




Figure 3 Comparison between control (CTL), dehydration-induced anorexia (DIA), and overnight food suppressed (OFS) rats. **(A)** Without manganese. † Represents significance level ($P < 0.05$) between CTL and DIA groups. ‡ Represents significance level ($P < 0.05$) between CTL and OFS groups. **(B)** In all, 24 hours postintravenous infusion of manganese. Only Glutamine (Gln) demonstrated a significant difference compared with CTL and OFS ($P < 0.01$). While significant differences exist between the neurochemical profile of CTL and the neurochemical profiles of DIA and OFS, these differences appear to be masked in the presence of manganese.

differences were found between DIA and OFS (two-tailed t -test; Gln, Glu, GABA: $P < 0.0001$; Lac: $P = 0.0048$; NAA: $P = 0.015$; Tau: $P < 0.05$). Significant differences also exist between CTL (+Mn²⁺), OFS (Mn²⁺), and DIA (+Mn²⁺) relative to the CTL group (two-tailed t -test between CTL (+Mn²⁺) and DIA (+Mn²⁺): PCr, Gln: $P < 0.0001$; between OFS (Mn²⁺) and DIA (+Mn²⁺): Gln, Glu, Lac: $P < 0.0001$). Statistically significant differences between groups of rats exposed to Mn²⁺ and rats nonexposed are also depicted in Figure 4. Although averaging groups of rats challenged by different conditions appear arbitrary, the 'gross' effect of Mn²⁺ on the neurochemical profile of the hypothalamus is clearly shown for Gln in DIA, Glu, and NAA in OFS and Tau for all conditions.

Discussion

Manganese is an essential element for normal rodent and human development. It is required for the appropriate functioning of a variety of physiological processes (Aschner *et al*, 2007; Benedetto *et al*, 2009). Recently, magnetic resonance imaging techniques have taken advantage of the paramagnetic properties of this compound identified as a potential indirect noninvasive neuronal marker (Duong *et al*,




Figure 4 Effect of manganese (Mn²⁺) on metabolites of the hypothalamus under normal (control, CTL), dehydration-induced anorexia (DIA), and overnight fasted (OFS) conditions. The calculated mean percentages of the CTL without manganese (\pm s.d.) are represented. Significant differences were found between DIA and OFS (*two-tailed t -test; Gln, Glu, γ -amino-butyric acid (GABA): $P < 0.0001$; Lac: $P = 0.0048$; NAA: $P = 0.015$; Tau: $P < 0.05$). Significant differences also exist between CTL (+Mn²⁺), OFS (Mn²⁺), and DIA (+Mn²⁺) relative to the CTL group (two-tailed t -test † between CTL (+Mn²⁺) and DIA (+Mn²⁺): PCr, Gln: $P < 0.0001$; ‡ between OFS (Mn²⁺) and DIA (+Mn²⁺): Gln, Glu, Lac: $P < 0.0001$). Statistically significant differences between groups of rats exposed to Mn²⁺ and rats nonexposed are also depicted (++ $P < 0.05$).

2000) and allowing the identification of various neuronal pathways within the rodent brain in tract tracing studies (Pautler *et al*, 1998, 2004). However, excessive levels of Mn²⁺ are toxic for the central nervous system (Stanwood *et al*, 2009). Silva *et al* (2004) have reported behavioral disorders such as somnolence, convulsions, tremors, and respiratory effects in both rats and mice at high Mn²⁺ doses (> 90 mg/kg) independent of the route of administration. Symptoms similar to Parkinson's disease have also been reported (Benedetto *et al*, 2009; Olanow, 2004; Pal *et al*, 1999). Moreover, Mn²⁺ was also reported to disrupt the astrocytes glutamine transporter expression and function in astrocytic cell cultures (Sidoryk-Wegrznowicz *et al*, 2009) and to alter dopaminergic and GABAergic neurons within the basal ganglia (Stanwood *et al*, 2009). Although the mechanisms underlying Mn²⁺ neurotoxicity remain unclear and chronic exposure to Mn²⁺ is a topic of interest since neuropathologic changes in humans lead to neuronal loss, gliosis and Alzheimer disease, the present study examined the effects of acute exposure to Mn²⁺ in relation to previous Mn²⁺-enhanced magnetic resonance imaging studies in the hypothalamus.

The hypothalamus is responsible for metabolic processes of the autonomic nervous system. This extremely complex brain region coordinates many

behavioral and physiological mechanisms as well as complex homeostatic processes. It must respond to many different signals internally and externally generated. Moreover, the hypothalamus is richly connected with the central nervous system. For an appropriate understanding of the function of hypothalamus, we challenged it with dehydration and fasting and investigated its metabolic response to Mn^{2+} exposure.

Comparison of the Neurochemical Profiles of the Hypothalamus and the Hippocampus of the Rat

The neurochemical profile of the normal rat hypothalamus was characterized by low PCr, Glu, Tau, and NAA concentrations relative to those of the hippocampus in agreement with a previous study in mice at 14.1 T (Lei *et al*, 2010). However, GABA and myo-Ins concentrations were also lower relative to the rat hippocampus, whereas these metabolites had significantly higher levels in the mouse hypothalamus relative to the hippocampus (Lei *et al*, 2010). γ -Aminobutyric acid is known to have an important inhibitory role in the regulation of the hypothalamic–pituitary–adrenocortical axis (Cullinan *et al*, 2008) and especially at the level of the paraventricular nucleus (Cullinan *et al*, 2008). No GABAergic neurons are located within the PVN itself (Cullinan *et al*, 2008), which may explain the lower GABA levels measured compared with the hippocampus in healthy rats.

In response to changes in brain homeostasis, metabolite concentrations in the hippocampus and the hypothalamus increased. A significant increase of GABA in DIA rats relative to CTL rats but also relative to the hippocampus was noticed.

In response to changes in osmolality, a cell shrinks or swells and finally exhibits a regulatory volume increase or decrease. On hypernatremic challenge, the cell loses water until intracellular and extracellular osmolality are equal. The recovery of water is mediated through accumulation of inorganic (Na^+ , K^+ , Cl^-) and organic (glutamate, glutamine, taurine, myo-Inositol) osmolytes. Gln, Glu, Tau, GABA, myo-Ins rise in the brain of chronic hypernatremic rats (Sterns and Silver, 2007) and return to their baseline levels as soon as normal osmolar conditions are recovered.

A reduction in brain water due to dehydration would be expected to decrease brain metabolite concentrations, 4 days after initiation of DIA. Dehydration-induced anorexia rats lost ~15% of their initial weight (Just and Gruetter, 2011). If it is assumed that the entire weight loss can be attributed to dehydration then a 15% total body water loss can be assumed and metabolite concentrations can be downscaled to 15%, which might be overestimated in the brain, since cellular dehydration is compensated by the continuous ingestion of water loaded with sodium and volume regulation occurs.

Strikingly, levels of NAA also increased in DIA ($P > 0.05$) and OFS ($P < 0.05$) rats. *N*-acetyl-aspartate

is believed to be a marker of neuronal density, integrity, and function. *N*-acetyl-aspartate reductions have been interpreted as reflecting neurotoxicity, neurodegeneration, or impaired mitochondrial energy metabolism. Consistent with this hypothesis, levels of NAA are reduced in a range of pathologic states with dramatic reductions in patients with tumor or stroke (Zimmerman and Wang, 1997). However, NAA has been identified as an osmolyte (Baslow, 2003). *N*-acetyl-aspartate is liberated into extracellular space in response to changing osmotic conditions but NAA was also hypothesized to be a molecular pump in which translocation creates an osmotic environment that passively removes water from the intracellular milieu of neurons (Baslow, 2003; Steen and Ogg, 2005).

Studies of psychiatric disorders have also shown that NAA levels can increase, suggesting a restoration or repair of neuronal viability (Moore *et al*, 2000; Mickael *et al*, 2003). The administration of lithium chloride to humans increased whole brain NAA levels, suggesting an increase of neuronal viability. However, lithium chloride also induces anorectic effects (Parkinson *et al*, 2009; Ishii *et al*, 2004). There were no significant difference in Mn^{2+} uptakes in hypothalamic regions involved in the control of food intake (arcuate nucleus, PVN, and supraoptic nucleus) between overnight fasted mice and mice receiving lithium chloride (Parkinson *et al*, 2009). The upregulation of NAA in OFS rats could be attributed to a transient neuroprotective effect of NAA in neurons in response to a short fasting period. This hypothesis is consistent with studies, demonstrating that fasting induces neuroprotective effects (Davis *et al*, 2008; Chiba *et al*, 2007).

γ -Aminobutyric Acid Responses to Hypertonicity and Manganese

Without $MnCl_2$, the hypothalamic inhibitory neurotransmitter GABA increased significantly in both DIA and OFS rats relative to CTL rats. γ -Aminobutyric acid is known to suppress feeding in rats and impact on body weight loss (Turenius *et al*, 2009). The majority of stress-responsive corticotropin-releasing hormone (CRH) neuronal inputs to the PVN contain the inhibitory neurotransmitter GABA, highlighting the importance of GABAergic neurotransmission in the regulation of the neuroendocrine stress response (Cullinan *et al*, 2008). Neurons of the PVN synthesize thyrotropin-releasing hormone and CRH hormones having an important role in energy homeostasis (Jaimes-Hoy *et al*, 2008). Under restricted food regime, rats show decreased thyrotropin-releasing hormone expression in the PVN while anorexic animals show increased thyrotropin-releasing hormone expression in the PVN. However, a reduction of CRH expression compared with CTL animals (Jaimes-Hoy *et al*, 2008) in the PVN was found in both food-restricted and DIA animals but to

a greater extent in DIA rats. This finding parallels the increase of GABA seen in DIA and OFS rats but to a greater extent in DIA rats. Yet, CRH-R2, one of the receptors of CRH signal involved in anorexigenic effects (de Gortari *et al*, 2009) was downregulated in the PVN of DIA rats but not in food-restricted animals (de Gortari *et al*, 2009). GABAergic projections from regions surrounding the PVN (anterior hypothalamic area, dorsomedial hypothalamic nucleus, LH, medial preoptic area) convey inhibitory signals to CRH neurons in response to a stressor such as DIA and food deficiency and limit hypothalamic activity by an important braking mechanism that includes increase of GABA release at the level of the PVN (Cullinan *et al*, 2008). It was suggested (Watts, 1999) that CRH neurons of the LH are specifically activated by the dehydration-induced stimulus and project to neurons in the PVN so that food inhibition due to low energy intake is overridden by dehydration mechanisms.

It is known that infertility can result when food intake fails to compensate for increased energy demands such as in anorexia nervosa. MnCl_2 stimulates the release of luteinizing hormone releasing hormone from the hypothalamus but simultaneously increases GABA release that in turn inhibits luteinizing hormone releasing hormone release (Prestifilippo *et al*, 2008). These processes also involve the activity of nitric oxide synthase (Prestifilippo *et al*, 2008) and may ultimately contribute to suppressed reproductive function in both male and female rats (Prestifilippo *et al*, 2008). As a consequence, further increases of GABA levels would have been expected in rats under both anorexic and Mn^{2+} condition. Nonetheless, GABA levels were similar in CTL (+ Mn^{2+}), DIA (+ Mn^{2+}), and OFS (+ Mn^{2+}) rats. Moreover, GABA concentrations were not significantly increased in DIA (+ Mn^{2+}) and OFS (+ Mn^{2+}) rats relative to their corresponding DIA and OFS rats. This suggests that the effects of dehydration and food restriction and MnCl_2 are not summed. MnCl_2 appears to blur the effects of anorexia and food restriction on GABA. Dehydration-induced anorexia animals demonstrated similar metabolite levels with and without MnCl_2 except for Tau. As pointed out earlier, both challenges should induce GABA release in the PVN but GABA levels were similar in DIA and DIA (+ Mn^{2+}) rats, suggesting that the effects of one condition may override the effects of the other. Salt loading and Mn^{2+} induce opposing effects at the cellular level. When exposed to hypertonicity, a cell shrinks until intracellular and extracellular osmolalities are equal. In contrast, Mn^{2+} induces cell swelling (Rama Rao *et al*, 2010). Cell swelling and cell shrinkage have opposing effects on Ca^{2+} concentration and hormonal secretion (Ben-Tabou De-Leon *et al*, 2003, 2006). Due to dehydration, morphological plasticity of astrocytes in the PVN occurred. Thus, synaptic and extrasynaptic transmission in magnocellular neurons of the PVN were modified with increased synaptic release of gluta-

mate and GABA but reduced ability of Tau to bind to glycinergic receptors and modulate neuronal excitability (Oliet and Bonfardin, 2010).

Previous investigations also suggest that signaling of astrocytes to neurons is performed through Ca^{2+} -dependent release of glutamate. We suggest that under hypertonic conditions, the Mn^{2+} enhancement seen in the PVN (and corresponding to neuronal activation) is mediated through astrocytic signaling. This hypothesis appears consistent with *in vitro* findings, showing that astrocytes increased their oxidative and anaerobic glucose metabolism to protect against energy failure and oxidative stress induced by Mn^{2+} . Simultaneously, Mn^{2+} induced neuronal energy failure (Zwingmann *et al*, 2003). For the maintenance of neuronal energy and neurotransmitter metabolism, the hypothesis of an astrocyte–neuron ‘lactate shuttle’ (Pellerin *et al*, 2007; Zwingmann *et al*, 2003) might hold.

γ -Aminobutyric acid levels increased significantly for all the conditions explored in the present study including with Mn^{2+} . Hence, GABA appears to be a marker of homeostatic changes within the rat hypothalamus but it cannot be considered as a specific marker.

Spectroscopic Analysis of the Hypothalamus Energy Metabolism in Manganese Neurotoxicity

The characteristic and significant GABA increases in fasted animals (Violante *et al*, 2009) and MnCl_2 -infused animals (Zwingmann *et al*, 2004, 2007) relative to normal animals were already described following the measurements of brain extracts using Carbon-13 (^{13}C)-NMR spectroscopy. These metabolic changes in the glutamine–glutamate–GABA cycle were attributed to impaired cerebral energy metabolism. This hypothesis was further confirmed since both fasted and MnCl_2 -infused animals demonstrated significantly increased lactate levels measured *ex vivo* relative to normal animals. Lactate synthesis is known to counteract decreased mitochondrial energy production (Zwingmann *et al*, 2004, 2007). Lactate also increased in our *in vivo* study for CTL (+ Mn^{2+}) rats but not for OFS (+ Mn^{2+}) rats. Violante *et al* (2009) attributed the increased GABA levels of fasted mice to a reduction in the GABA degradation pathway since no significant increases in Glu and Gln were detected. Without MnCl_2 , Glu levels increased significantly in our OFS rats relative to CTL. This would point toward increased synthesis of GABA through Glu and Gln precursors while in OFS (+ Mn^{2+}) rats, no significant difference was detected relative to CTL, suggesting a decreased degradation of GABA (Violante *et al*, 2009). However, as pointed out earlier, increasing levels of GABA did not appear to be condition specific including in the presence of MnCl_2 (Figure 4). On the contrary, Mn^{2+} clearly influenced Tau, NAA, Cr, and PCr as well Glc levels relative to the CTL group of rats. Moreover, Gln and Glu levels appeared more

specifically affected by Mn^{2+} for DIA (+ Mn^{2+}) and OFS (+ Mn^{2+}) rats. Zwingmann *et al* (2004, 2007) showed that Mn^{2+} disturbs glucose metabolism, which is supported by the results of the present study. Additionally, increases in glutamine levels following exposure of cells to Mn^{2+} were attributed to an impaired efflux of Gln from astrocytes or an impaired uptake of Gln from neurons (Zwingmann *et al*, 2003). These results were consistent with a increased glutamine synthetase activity in Mn^{2+} -intoxicated rats (Zheng *et al*, 1999) and match as well with findings that simultaneous increases in Gln levels with decreases in Tau levels are consistent with alterations in astrocyte morphology (Oliet and Bonfardin, 2010). In DIA (+ Mn^{2+}), Gln increases were further exacerbated compared with all the other groups of rats and relative to CTL, but we also noted that a significant difference between Gln levels in DIA and OFS rats remains with and without Mn^{2+} . In future studies, the role of glutamine in hypothalamic astrocytes and neurons of anorexic rats should be further investigated.

Significantly lower Tau levels were noticed for both OFS (+ Mn^{2+}) and DIA (+ Mn^{2+}) rats relative to non- $MnCl_2$ -infused rats. This change was not noticed in CTL rats. Relative to CTL rats, Tau levels did not change significantly for the OFS (+ Mn^{2+}) and DIA (+ Mn^{2+}) conditions, whereas higher Tau levels were measured in OFS and DIA rats without $MnCl_2$.

Taurine is an amino acid that has neuroprotective properties and is involved as an osmoregulator (Song and Hatton, 2003). Taurine uptake was increased in astrocyte cultures exposed to a hyperosmotic condition (Aschner *et al*, 2001), while Tau variations have already been observed in short-term food-deprived animals (Benuck *et al*, 1995). As in the present study, Tau concentrations in the globus pallidus of rats measured *in vivo* by NMR spectroscopy (Zwingmann *et al*, 2007) after $MnCl_2$ infusion demonstrated significant decreases compared with control rats. Taurine is also an important neurotransmitter known to modulate GABA neurochemistry (Namima *et al*, 1982, 1983) and share GABA transporters in the brain. Here, due to Mn^{2+} exposure taurine responded inversely to GABA. Previous data suggested that Mn^{2+} increases GABA by inhibition of GABA transporter function. In turn, the osmoregulatory and neuromodulatory functions of taurine are altered and perturb neural function (Fordahl *et al*, 2010).

Myo-Inositol concentrations were also increased with $MnCl_2$ in the globus pallidus compared with controls without $MnCl_2$ whereas NAA levels were decreased (Zwingmann *et al*, 2007). Moreover, these changes appear to be regional specific (Zwingmann *et al*, 2007).

Conclusion

In the present work, the metabolism of the hypothalamus in rats was characterized both under normal

conditions and following changes in brain energy homeostasis using a model of DIA and overnight fasting in rats. The data presented in this study show that it is necessary to perform region-specific 1H -MRS for an accurate characterization of the neurochemical profile of the hypothalamus. In addition, compared with previous studies (Lei *et al*, 2010), it shows that an improved characterization is performed when challenges such as dehydration, fasting, and Mn^{2+} infusion are applied: the data of the present study provide evidence that GABA has an essential role in the maintenance of energy homeostasis in the hypothalamus. However, results also reveal that GABA is a nonspecific marker of Mn^{2+} intoxication in the hypothalamus. On the contrary, glutamine, glutamate, and taurine appear to respond more accurately to Mn^{2+} exposure. An increase in glutamine levels could be a characteristic response of the hypothalamus to DIA.

Disclosure/conflict of interest

The authors declare no conflict of interest.

References

- Aoki I, Naruse S, Tanaka C (2004) Manganese-enhanced magnetic resonance imaging (MEMRI) of brain activity and applications to early detection of brain ischemia. *NMR Biomed* 17:569–80
- Aschner M, Guilarte TR, Schneider JS, Zheng W (2007) Manganese: recent advances in understanding its transport and neurotoxicity. *Toxicol Appl Pharmacol* 221: 131–47
- Aschner M, Mutkus L, Allen JW (2001) Aspartate and glutamate transport in acutely and chronically ethanol exposed neonatal rat primary astrocyte cultures. *Neurotoxicology* 22:601–5
- Baslow MH (2003) N-acetylaspartate in the vertebrate brain: metabolism and function. *Neurochem Res* 28: 941–53
- Benedetto A, Au C, Aschner M (2009) Manganese-induced dopaminergic neurodegeneration: insights into mechanisms and genetics shared with Parkinson's disease. *Chem Rev* 109:4862–84
- Benuck M, Banay-Schwartz M, DeGuzman T, Lajtha A (1995) Effect of food deprivation on glutathione and amino acid levels in brain and liver of young and aged rats. *Brain Res* 678:259–64
- Ben-Tabou De-Leon S, Ben-Zeev G, Nussinovitch I (2006) Effects of osmotic shrinkage on voltage-gated Ca^{2+} channel currents in rat anterior pituitary cells. *Am J Physiol Cell Physiol* 290:C222–32
- Ben-Tabou De-Leon S, Blotnick E, Nussinovitch I (2003) Effects of osmotic swelling on voltage-gated calcium channel currents in rat anterior pituitary cells. *Am J Physiol Cell Physiol* 285:C840–52
- Chaudhri OB, Parkinson JR, Kuo YT, Druce MR, Herlihy AH, Bell JD, Dhillo WS, Stanley SA, Ghatei MA, Bloom SR (2006) Differential hypothalamic neuronal activation following peripheral injection of GLP-1 and oxyntomodulin in mice detected by manganese-enhanced

- magnetic resonance imaging. *Biochem Biophys Res Commun* 350:298–306
- Chiba T, Fujita S, Kubota H, Inoue D, Mizuno A, Komatsu T, Yamaza H, Higami Y, Shimokawa I (2007) Identification of fasting-induced genes in the rat hypothalamus: relationship with neuroprotection. *Ann NY Acad Sci* 1119:216–26
- Cullinan WE, Ziegler DR, Herman JP (2008) Functional role of local GABAergic influences on the HPA axis. *Brain Struct Funct* 213:63–72
- de Gortari P, Mancera K, Cote-Vélez A, Amaya MI, Martínez A, Jaimes-Hoy L, Joseph-Bravo P (2009) Involvement of CRH-R2 receptor in eating behavior and in the response of the HPT axis in rats subjected to dehydration-induced anorexia. *Psychoneuroendocrinology* 34:259–72
- Davis LM, Pauly JR, Readnower RD, Rho JM, Sullivan PG (2008) Fasting is neuroprotective following traumatic brain injury. *J Neurosci Res* 86:1812–22
- Duong TQ, Silva AC, Lee SP, Kim SG (2000) Functional MRI of calcium-dependent synaptic activity: cross correlation with CBF and BOLD measurements. *Magn Reson Med* 43:383–92
- Fordahl SC, Anderson JG, Cooney PT, Weaver TL, Colyer CL, Erikson KM (2010) Manganese exposure inhibits the clearance of extracellular GABA and influences taurine homeostasis in the striatum of developing rats. *Neurotoxicology* 31:639–46
- Gruetter R, Tkáč I (2000) Field mapping without reference scan using asymmetric echo-planar techniques. *Magn Reson Med* 43:319–23
- Ishii Y, Blundell JE, Halford JC, Upton N, Porter R, Johns A, Rodgers RJ (2004) Differential effects of the selective orexin-1 receptor antagonist SB-334867 and lithium chloride on the behavioural satiety sequence in rats. *Physiol Behav* 81:129–40
- Jaimes-Hoy L, Joseph-Bravo P, de Gortari P (2008) Differential response of TRHergic neurons of the hypothalamic paraventricular nucleus (PVN) in female animals submitted to food-restriction or dehydration-induced anorexia and cold exposure. *Horm Behav* 53:366–77
- Just N, Gruetter R (2011) Detection of neuronal activity and metabolism in a model of dehydration-induced anorexia in rats at 14.1T using manganese-enhanced MRI and proton MR spectroscopy. *NMR Biomed*; published online, 4 May 2011; doi:10.1002/nbm.1694
- Lei H, Poitry-Yamate C, Preitner F, Thorens B, Gruetter R (2010) Neurochemical profile of the mouse hypothalamus using *in vivo* (1) H MRS at 14.1T. *NMR Biomed* 23:578–83
- Mickael N, Erfurth A, Ohrmann P, Arolt V, Heindel W, Pfeiderer B (2003) Neurotrophic effects of electroconvulsive therapy: a proton magnetic resonance study of the left amygdala region in patients with treatment-resistant depression. *Neuropsychopharmacology* 28:720–5
- Mlynárik V, Gambarota G, Frenkel H, Gruetter R (2006) Localized short-echo-time proton MR spectroscopy with full signal-intensity acquisition. *Magn Reson Med* 56: s965–70
- Moore GJ, Bebchuk JM, Hasanat K, Chen G, Seraji-Bozorgzad N, Wilds IB, Faulk MW, Koch S, Glitz DA, Jolkovsky L, Manji HK (2000) Lithium increases N-acetyl-aspartate in the human brain: *in vivo* evidence in support of bcl-2's neurotrophic effects? *Biol Psychiatry* 48:1–8
- Namima M, Okamoto K, Sakai Y (1982) Taurine acts on presynaptic autoreceptors for GABA in the cerebellum: effects on Ca²⁺ influx and GABA release. *Jpn J Pharmacol* 32:746–9
- Namima M, Okamoto K, Sakai Y (1983) Modulatory action of taurine on the release of GABA in cerebellar slices of the guinea pig. *J Neurochem* 40:1–9
- Olanow CW (2004) Manganese-induced Parkinsonism and Parkinson's disease. *Ann NY Acad Sci* 1012:209–23
- Oliet S, Bonfardin V (2010) Morphological plasticity of the rat supraoptic nucleus-cellular consequences. *Eur J Neurosci* 32:1989–94
- Pal PK, Samii A, Calne DB (1999) Manganese neurotoxicity: a review of clinical features, imaging and pathology. *Neurotoxicology* 20:227–38
- Parkinson JR, Chaudhri OB, Kuo YT, Field BCT, Herlihy AH, Dhillon WS, Ghatei MA, Bloom SR, Bell JD (2009) Differential patterns of neuronal activation in the brainstem and hypothalamus following peripheral injection of GLP-1, oxyntomodulin and lithium chloride in mice detected by manganese-enhanced magnetic resonance imaging (MEMRI). *Neuroimage* 44:1022–31
- Pautler RG (2004) *In vivo*, trans-synaptic tract-tracing utilizing manganese-enhanced magnetic resonance imaging (MEMRI). *NMR Biomed* 17:595–601
- Pautler RG, Silva AC, Koretsky AP (1998) *In vivo* neuronal tract tracing using manganese-enhanced magnetic resonance imaging. *Magn Reson Med* 40:740–8
- Paxinos G, Watson C (1998) *The Rat Brain in Stereotaxic Coordinates*. San Diego: Academic Press
- Pellerin L, Bouzier-Sore AK, Aubert A, Serres S, Merle M, Costalat R, Magistretti PJ (2007) Activity-dependent regulation of energy metabolism by astrocytes: an update. *Glia* 55:1251–62
- Prestifilippo JP, Fernández-Solari J, De Laurentiis A, Mohn CE, de la Cal C, Reynoso R, Dees WL, Rettori V (2008) Acute effect of manganese on hypothalamic luteinizing hormone releasing hormone secretion in adult male rats: involvement of specific neurotransmitter systems. *Toxicol Sci* 105:295–302
- Provencher SW (1993) Estimation of metabolite concentrations from localized *in-vivo* proton MR spectra. *Magn Reson Med* 30:672–9
- Rama Rao K, Jayakumar AR, Reddy PVB, Tong X, Curtis KM, Noremborg MD (2010) Aquaporin-4 in manganese-treated cultured astrocytes. *Glia* 58:1490–9
- Salter-venzon D, Watts AG (2008) The role of hypothalamic ingestive behavior controllers in generating dehydration anorexia: a Fos mapping study. *Am J Physiol Regul Integr Comp Physiol* 295:R1009–19
- Salter-Venzon D, Watts AG (2009) Site-specific attenuation of food intake but not the latency to eat after hypothalamic injections of neuropeptide Y in dehydrated-anorexic rats. *Am J Physiol Regul Integr Comp Physiol* 297:R1813–21
- Sidoryk-Wegrzynowicz M, Lee E, Albrecht J, Aschner M (2009) Manganese disrupts astrocyte glutamine transporter expression and function. *J Neurochem* 110: 822–30
- Silva AC, Bock NA (2008) Manganese-enhanced MRI: an exceptional tool in translational neuroimaging. *Schizophr Bull* 34:595–604
- Silva AC, Lee JH, Aoki I, Koretsky AP (2004) Manganese-enhanced magnetic resonance imaging (MEMRI): methodological and practical considerations. *NMR Biomed* 17:532–43
- Silva AC, Lee JH, Wu CW, Tucciarone J, Pelled G, Aoki I, Koretsky AP (2008) Detection of cortical laminar

- architecture using manganese-enhanced MRI. *J Neurosci Methods* 167:246–57
- Song Z, Hatton GI (2003) Taurine and the control of basal hormone release from rat neurohypophysis. *Exp Neurol* 183:330–7
- Stanwood GD, Leitch DB, Savchenko V, Wu J, Fitsanakis VA, Anderson DJ, Stankowski JN, Aschner M, McLaughlin B (2009) Manganese exposure is cytotoxic and alters dopaminergic and GABAergic neurons within the basal ganglia. *J Neurochem* 110:378–89
- Steen RG, Ogg RJ (2005) Abnormally high levels of brain N-acetylaspartate in children with sickle cell disease. *AJNR Am J Neuroradiol* 26:463–8
- Sterns RH, Silver SM (2007) Physiology and pathophysiology: hyponatremic states. In *Seldin and Giebisch's The Kidney* (Alpern RJ, Hebert SC, eds). Academic Press: London, pp. 1203–1224
- Tkác I, Starcuk Z, Choi IY, Gruetter R (1999) *In vivo* 1H NMR spectroscopy of rat brain at 1 ms echo time. *Magn Reson Med* 41:649–56
- Turenius CI, Htut MM, Prodon DA, Ebersole PL, Ngo PT, Lara RN, Wilczynski JL, Stanley BG (2009) GABA(A) receptors in the lateral hypothalamus as mediators of satiety and body weight regulation. *Brain Res* 1262:16–24
- Violante IR, Anastasovska J, Sanchez-Canon GJ, Rodrigues TB, Righi V, Nieto-Charques L, Parkinson JR, Bloom SR, Bell JD, Cerdán S (2009) Cerebral activation by fasting induces lactate accumulation in the hypothalamus. *Magn Reson Med* 62:279–83
- Watts AG (1999) Dehydration-associated anorexia: development and rapid reversal. *Physiol Behav* 65:871–8
- Zheng W, Zhao Q, Slavkovich V, Aschner M, Graziano JH (1999) Alterations in ion homeostasis following chronic exposure to manganese in rats. *Brain Res* 833:125–32
- Zimmerman RA, Wang ZJ (1997) The value of proton MR spectroscopy in pediatric metabolic brain disease. *AJNR Am J Neuroradiol* 18:1872–9
- Zwingmann C, Leibfritz D, Hazell AS (2003) Energy metabolism in astrocytes and neurons treated with manganese: relation among cell-specific energy failure, glucose metabolism, and intercellular trafficking using multinuclear NMR-spectroscopic analysis. *J Cereb Blood Flow Metab* 23:756–71
- Zwingmann C, Leibfritz D, Hazell AS (2004) Brain energy metabolism in a sub-acute rat model of manganese neurotoxicity: an *ex vivo* nuclear magnetic resonance study using [1-¹³C] glucose. *Neurotoxicology* 25:573–87
- Zwingmann C, Leibfritz D, Hazell AS (2007) NMR spectroscopic analysis of regional brain energy metabolism in manganese neurotoxicity. *Glia*: 55:1610–7

Supplementary Information accompanies the paper on the Journal of Cerebral Blood Flow & Metabolism website (<http://www.nature.com/jcbfm>)