



Visual consciousness and bodily self-consciousness

Nathan Faivre^{a,b,*}, Roy Salomon^{a,b,*}, and Olaf Blanke^{a,b,c}

Purpose of review

In recent years, consciousness has become a central topic in cognitive neuroscience. This review focuses on the relation between bodily self-consciousness – the feeling of being a subject in a body – and visual consciousness – the subjective experience associated with the perception of visual signals.

Recent findings

Findings from clinical and experimental work have shown that bodily self-consciousness depends on specific brain networks and is related to the integration of signals from multiple sensory modalities including vision. In addition, recent experiments have shown that visual consciousness is shaped by the body, including vestibular, tactile, proprioceptive, and motor signals.

Summary

Several lines of evidence suggest reciprocal relationships between vision and bodily signals, indicating that a comprehensive understanding of visual and bodily self-consciousness requires studying them in unison.

Keywords

bodily illusions, bodily self-consciousness, body ownership, embodiment, visual consciousness

INTRODUCTION

Brain activity causes subjective experience. It ‘feels like something’ to see a rose, and this feeling is qualitatively different from touching or smelling it and seeing an orchid [2]. For the past 20 years, there have been considerable efforts to describe the so-called ‘neural correlates of consciousness’ (NCC), which is the minimal set of neuronal events and mechanisms sufficient for a specific conscious percept [3]. The NCC have traditionally been sketched using a contrastive approach, whereby neural and cognitive responses elicited by perceived vs. unperceived stimuli are compared [4]. Theories that emerged from this approach state that consciousness requires, for example, information integration throughout the brain, involving mechanisms such as short-range [5] or long-range [6] feedback connections, or neural synchrony [7]. Most data and greatest detail regarding the neural correlates of perceptual consciousness arguably comes from visual psychophysics and brain imaging, where an arsenal of techniques allows the display of subliminal, invisible images in various manners [8^a,9]. Although these studies led to a better understanding of visual consciousness, the interaction of conscious and unconscious vision with other senses remains to be explored, which is crucial considering the multi-sensory nature of perception [10^a]. One can note that the sense of touch constitutes a promising

model to study perceptual consciousness, considering its rather ‘primitive’ attributes, with both phylogenetic and ontogenetic roots preceding those of vision [1]. Accordingly, its functional organization and the computational steps underlying tactile consciousness are likely to be simpler compared to vision, and therefore more accessible to empirical investigations and theoretical interpretations.

Moreover, subjective aspects of perceptual consciousness (or qualia) have remained elusive in models of visual consciousness or continue to be associated with most disagreement among consciousness researchers and philosophers (for a recent example, see [11] and the related commentary [12]). Notably, there is no consensus as to whether a theory of visual consciousness should explain how phenomenal experience arises from physical events in the brain (the hard problem or

^aLaboratory of Cognitive Neuroscience, Brain Mind Institute, ^bCenter for Neuroprosthetics, School of Life Sciences, École Polytechnique Fédérale de Lausanne, Lausanne and ^cDepartment of Neurology, University Hospital, Geneva, Switzerland

Correspondence to Olaf Blanke, Faculty of Life Sciences, École Polytechnique Fédérale de Lausanne, Station 19, Lausanne CH-1015, Switzerland. Tel: +41 21 693 9621; e-mail: olaf.blanke@epfl.ch

*Nathan Faivre and Roy Salomon contributed equally to the writing of this article.

Curr Opin Neurol 2015, 28:000–000

DOI:10.1097/WCO.000000000000160

KEY POINTS

- BSC stems from the integration of visual, tactile, proprioceptive, and vestibular signals.
- Visual perception reflects the product of integration with other sensory signals, including bodily signals.
- Multisensory integration occurs unconsciously, prior to conscious access, potentially resulting in phenomenal unity.

phenomenal consciousness) or rather characterize the mechanisms supporting cognitive access to consciousness (the easy problem or access consciousness) [13,14].

Furthermore, the observer who is the subject of conscious experience is generally not accounted for by these models of visual consciousness. Yet, a fundamental property of consciousness is its link with the self: the subject of conscious experience. The 'rose' that is seen, felt, or smelled is bounded to the 'I' of subjective experience, is felt by somebody, the self or subject of experience. This sense of self and its close link to neural body representation has been termed bodily self-consciousness (BSC) and has been a target of recent research [15]. BSC is commonly thought to involve self-identification (the experience of owning 'my' body), self-location (the experience of where 'I' am in space), and first-person perspective (the experience from where 'I' perceive the world) [16]. The scientific study of BSC has expanded over the last years, benefiting from the examination of neurological patients with altered states of BSC and virtual reality coupled with robotics to induce well controlled states in the research laboratory. For instance, altered states of BSC of neurological origin may be experienced from a location and perspective that are not centered on the physical body of the observer, but are nevertheless characterized by an experience of the visual world that is complete, differentiated, and integrated. In these so-called out-of-body experiences, the subject or center of consciousness is experienced as being displaced to an elevated and down-looking first-person perspective and location [17]. The 'rose' on a table in the field of view of the observer would thus still be perceived consciously during an out-of-body experience, but seen from a distanced and elevated location and first-person perspective (see below for recent studies in healthy subjects on BSC).

Do visual consciousness and BSC relate to each other? It has been argued that (visual) perception is by essence embodied within an egocentric, body-centered framework ([18,19]; but see [20] for a critical discussion about embodiment and consciousness).

Much less is known about whether and how bodily processing impacts visual consciousness and how visual processing relates to BSC, as each aspect of consciousness has mostly been studied in relative separation. However, consciousness at the phenomenal level is an integrated and unitary experience: integrated as multiple sensory streams are seamlessly joined, and unitary as consciousness is experienced by a singular self [21]. Interestingly, integrated visual consciousness may break down in neurological patients with simultagnosia reporting the inability to perceive more than one object at once [22], while unitary consciousness may break down in neurological patients with heautoscopy reporting two simultaneous, but distinct, centers of consciousness [23[¶]]. Here, we review empirical results documenting the interplay between visual consciousness and BSC, pointing toward the importance of their combined investigation for a fuller understanding of consciousness.

INFLUENCES OF VISION ON BODILY SELF-CONSCIOUSNESS

There is now compelling evidence that BSC involves the cortical integration of different signals from the external environment (exteroception: visual, auditory, and tactile senses, see [24] for a review) and the body (vestibular and proprioceptive senses; interoceptive cardiac and respiratory signals, see [25]). Data from neurological patients with altered states of BSC were associated with deficits in multisensory integration and involved the right temporoparietal cortex with abnormal self-location and first-person perspective, but unitary consciousness [26], and the left insula with abnormal self-location, first-person perspective, and the loss of unitary consciousness [23[¶]]. Recent experimental paradigms in healthy participants employed multisensory conflicts, reporting modulations of the three aforementioned main aspects of BSC. For example, in the well known rubber-hand illusion, participants viewing an artificial hand being stroked in synchrony with their own occluded hand (placed at a different position) experience that the artificial hand belongs to their own body, and perceive their hand as shifted toward the artificial hand's position [27]. This illusion is thought to stem from visual capture of the multisensory conflict between visual, proprioceptive, and tactile stimuli that are temporally synchronous (visual, tactile) but spatially incongruent (visual, proprioceptive): what is seen (an artificial hand being stroked) does not match with what is felt (one's hand being stroked). With temporal delay between the visual and tactile stimulation, the illusion and

shift in perceived hand position vanish (but see [28]). In recent years, the development of video, virtual reality, and robotic devices have allowed researchers to extend the study of limb ownership to full-body ownership and other aspects of BSC [15]. In the so-called full-body (or ‘out-of-body’) illusion (FBI), multisensory conflicts modulate not only limb ownership but also ownership/self-identification as well as self-location for the full body, mimicking neurological conditions such as out-of-body experiences, heautoscopy, and related altered states of BSC [29,30]. Typically, participants view their filmed own body (a ‘virtual body’) from a posterior third-person perspective while perceiving repeated tactile stimulations on their back or chest. As in the rubber-hand illusion, conflicting multisensory stimulation exists between what is seen (the avatar being stroked) and what is felt (the participant’s back being stroked). Compared to asynchronous visuotactile stroking, participants in synchronous visuotactile stroking self-identify more with the seen virtual body, judge their positions as closer to it, and mislocalize the tactile stimulus to the virtual body. Additional studies have shown autonomic bodily responses, such as skin cooling stemming from such bodily illusions [31,32^{*}]. Brain imaging studies using the FBI have revealed that the premotor cortex, intraparietal sulcus, and extrastriate visual regions are involved in the sense of body ownership, whereas the temporoparietal cortex and insula are involved in the sense of self-location and first-person perspective [26,33^{*},34].

These results support the idea that BSC stems from the integration of visual, tactile, proprioceptive [35], and vestibular [36] signals. Both in the rubber-hand and the FBIs, subjective changes in BSC are held to arise from a resolution of multisensory conflicts by means of minimization of error signals [37^{**},38^{**}]. Visual dominance is prevalent in that respect, as it dominates proprioceptive inputs [39] when the apparent seen location of a body part does not match its actual location. Such visual modulation of bodily signals is also reflected in the tactile domain, as seeing one’s hand improves tactile acuity on the hand, compared to seeing a nonhand object (visual enhancement of touch [40]). This latter effect is further boosted by hand ownership manipulations [41], as revealed by the crossmodal congruency effect for hands [42] and avatars [43]. Taken together, these results highlight the role of visual signals in experimentally and neurologically induced altered states of BSC. In the next section, we evaluate the complementary idea by assessing whether vision is affected by bodily signals.

INFLUENCES OF BODILY SIGNALS ON VISION

On a continuous basis, coalitions of interoceptive and exteroceptive sensors convey information about various aspects of the body and the environment. Despite this tremendous amount of information, percepts are not experienced as sums of independent features, but as integrated multimodal scenes [21]. That is, colors, smells, and textures are all experienced as integrated into a common perceptual object. The same holds for bodily experience, as subjects typically experience themselves as a unified whole (but see heautoscopy; [16,23^{*}]). As argued above, consciousness is a unified and essentially multisensory experience, rather than a sum of unimodal features [10^{*}]. Multisensory integration – the function whereby information from different sensory modalities is combined together – is thus central to both visual consciousness and BSC. Building upon the notion of phenomenal unity (i.e., the perception of multiple sensory streams as a multimodal), most theories of visual consciousness postulate strong interdependencies between multisensory integration and consciousness (see [44^{**}] for a review). Yet, these interdependencies are unlikely to imply necessity, as sensory signals from different modalities are integrated at early stages or processing already in subcortical structures, arguably unconsciously [45]. In addition, there is also compelling evidence for unconscious multisensory integration of more complex stimuli, as the perception of invisible images can be modulated by the simultaneous presence of another supraliminal stimulus presented in the auditory [46^{*},47^{*}], tactile [48,49^{*}], or proprioceptive modalities [50^{*}]. Beyond the integration of a supraliminal and a subliminal stimulus, a recent study showed that an invisible stimulus can also be integrated with an inaudible one (up to semantic levels), therefore arguing for high-level multisensory integration in the complete absence of perceptual consciousness [51^{*}]. Taken together, these results suggest that multisensory integration that is of relevance for visual consciousness and BSC most likely is an automatic brain process, governed by bottom-up mechanisms that precede conscious access.

If perception and visual consciousness are embodied, changes in visual consciousness depending on bodily signals are expected. Yet, the mere idea of all-or-none modulations of visual perception depending on BSC seems hardly testable empirically, as bodily signals of interoceptive, vestibular, or proprioceptive origin are never completely disrupted, even in the most severe neurological conditions (e.g., locked-in syndrome, deafferentation, and tetraplegia) or experimental manipulations (e.g., sensory deprivation).

However, several studies have investigated directly the impact of bodily signals on thresholds for visual consciousness. One focused on proprioception and showed that images of hands emerged faster into visual consciousness when their orientation was congruent with the actual position of the participants' hands when seeing the display (palm up vs. palm down) [50[□]]. This suggests that proprioceptive signals can modulate the threshold for visual consciousness for an object in a quantitative manner. Another study applied a similar experimental logic to touch–vision interactions showing facilitatory effects of congruent tactile stimuli on visual consciousness [52[□]]. More recently, it was also shown that neural events locked to heartbeats correlated with the detection of a faint visual grating, revealing the impact of interoceptive bodily cues (cardiac signal) on visual consciousness [53[□]] as also revealed for BSC [25,54[□]]. Taken together these studies suggest that visual consciousness is facilitated when visual signals are congruent with bodily signals at the spatial or temporal level, and inhibited in the presence of multimodal conflicts. Further studies are required to investigate these effects and the possible influences of other sensory modalities such as vestibular signals on visual consciousness, and the involved neural mechanism responsible for such multisensory effects.

Beyond the threshold for visual consciousness, there is long-standing compelling evidence that bodily signals can alter the content of visual perception. Indeed, numerous studies reported changes in visual percepts depending on vestibular inputs (e.g., oculogravic illusion [55,56]; bistable perception [57[□]]), proprioceptive inputs (e.g., Taylor illusion [58]; kinesthetic visual motion [59[□]]), tactile inputs (e.g., bistable perception [52[□]]), and motor inputs (e.g., biological motion [60–62]). An open question is also whether certain signals impacting BSC, play a particular role in shaping visual consciousness.

Taken together, the reviewed data suggest that the quality of visual perception is not a simple function of retinal input, but also the product of integration with other sensory signals, including bodily signals. This bidirectional interaction in which bodily and visual information modulate one another indicate that to comprehensively understand visual consciousness, or BSC, or both, they must be studied together, including their influence on self-consciousness.

OPEN QUESTIONS

We reviewed clinical and experimental evidence suggesting a bidirectional relationship between visual consciousness and BSC. Many questions remain

regarding this relationship, with three intriguing examples below.

Does bodily self-consciousness require perceptual consciousness?

Is BSC based on the (perceptually) conscious brain or are unconscious multisensory processes sufficient to induce BSC? The latter seems likely considering that vestibular and proprioceptive signals are both primordial for BSC, while remaining most of the time outside the stream of perceptual consciousness under normal conditions. Progress on this issue would require assessment of whether changes of self-identification and self-location can be obtained in cases when multisensory conflicts are not detected explicitly (i.e., participants feel the visuotactile stimuli but cannot discriminate whether they are synchronous or asynchronous), or in case the visuotactile stimuli themselves remain under the threshold for perceptual consciousness (i.e., due to psychophysical manipulations or pathological conditions such as blindsight).

Does bodily self-consciousness involve attention?

Although the necessity and sufficiency of attention for visual consciousness is subject to a heated debate [63–65], it has only rarely been explicitly explored in the field of BSC, with the exception of neurological research revealing close links between attention and altered states of BSC (e.g., somatoparaphrenia). We recently showed in a visual search paradigm with self-motion and nonself-motion stimuli that self-movement 'pops-out', suggesting that visual attention is integrated with efferent information at early stages of processing [66[□]]. Experiments manipulating attention and investigating its influence on BSC (and visual consciousness) would be important and may range from investigations of preattentive to explicit top–down aspects.

Does bodily self-consciousness influence cognitive processes beyond perception?

Recently, metacognitive vision, that is the second-order knowledge about a visual process, or 'knowing about seeing', has received much attention [67]. So far, studies about metacognition have focused on unisensory perception, and the role of multisensory processes in metacognitive processes remains unknown: is what we feel we know about a bimodal stimulus substantially better than what we know about a unimodal stimulus? Although, distinct from the notion of the minimal sense of self reviewed

here (BSC), it seems interesting to investigate whether metacognition relates differentially to BSC ('knowing about feeling') vs. more classical cognitive aspects of the self (such as narrative, remembered, and conceptual aspects of the self) ('knowing about knowing').

CONCLUSION

Picturing the interplay between visual consciousness and BSC presents both methodological and theoretical challenges. We believe that deciphering the interactions between perception of the environment, perception of one's body, and their respective necessity and sufficiency for visual consciousness and BSC require an integrated, multisensory study of consciousness.

Acknowledgements

The authors thank Steven Gale for his helpful comments on the manuscript.

Financial support and sponsorship

This work was supported by the National Center of Competence in Research - Synapsy (The Synaptic Bases of Mental Diseases) and financed by the Swiss National Science Foundation (#51AU40_125759) and by the Bertarelli foundation. N.F. was supported by the EU Human Brain Project. R.S. was supported by the National Center of Competence in Research (NCCR) "SYNAPSY—The Synaptic Bases of Mental Diseases" financed by the Swiss National Science Foundation (no. 51AU40_125759).

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Gregory R. Origin of eyes and brains. *Nature* 1967; 213:369–372.
 2. Nagel T. What is it like to be a bat? *Philos Rev* 1974; 83:435–450.
 3. Koch C. *The quest for consciousness: a neurobiological approach*. Englewood, CO: Roberts; 2004.
 4. Baars B. Contrastive phenomenology: a thoroughly empirical approach to consciousness. In: *The nature of consciousness: philosophical debates*. Cambridge, MA: MIT Press; 1997.
 5. Lamme V, Roelfsema PR. The distinct modes of vision offered by feedforward and recurrent processing. *Trends Neurosci* 2000; 23:571–579.
 6. Dehaene S, Changeux JP. Experimental and theoretical approaches to conscious processing. *Neuron* 2011; 70:200–227.
 7. Engel AK, Singer W. Temporal binding and the neural correlates of sensory awareness. *Trends Cognit Sci* 2001; 5:10749–10753.
 8. Dubois J, Faivre N. Invisible, but how? The depth of unconscious processing as inferred from different suppression techniques. *Front Psychol* 2014; 5:1117.
 9. Kim CY, Blake R. Psychophysical magic: rendering the visible 'invisible'. *Trends Cognit Sci* 2005; 9:381–388.
 10. Derooy O, Chen Y, Spence C. Multisensory constraints on awareness. *Philos Trans Royal Soc B: Biol Sci* 2014; 369:20130207.
- This review argues in favor of a multisensory study of perceptual consciousness and highlights the methodological and theoretical challenges associated with such a perspective.
11. Cohen M, Dennett DC. Consciousness cannot be separated from function. *Trends Cognit Sci* 2011; 15:358–364.
 12. Fahrenfort JJ, Lamme VAF. A true science of consciousness explains phenomenology: comment on Cohen and Dennett. *Trends Cognit Sci* 2012; 16:138–139.
 13. Chalmers D. Facing up to the problem of consciousness. *J Conscious Stud* 1995; 2:1–27.
 14. Block N. How many concepts of consciousness? *Behav Brain Sci* 1995; 18:272–287.
 15. Blanke O. Multisensory brain mechanisms of bodily self-consciousness. *Nat Rev Neurosci* 2012; 13:556–571.
 16. Blanke O, Metzinger T. Full-body illusions and minimal phenomenal selfhood. *Trends Cognit Sci* 2009; 13:7–13.
 17. Blanke O, Landis T, Spinelli L, Seeck M. Out-of-body experience and autoscopia of neurological origin. *Brain* 2004; 127:243–258.
 18. Damasio A. *The feeling of what happens: body, emotion and the making of consciousness*. New York: Mariner Books; 2000.
 19. Gallagher S. Philosophical conceptions of the self: implications for cognitive science. *Trends Cognit Sci* 2000; 4:14–21.
 20. Prinz J. *Is consciousness embodied. The Cambridge handbook of situated cognition*. New York: Cambridge University Press 2009; Cambridge University Press.
 21. Bayne T, Chalmers D. What is the unity of consciousness? In: Cleeremans, AE, editor. *The unity of consciousness: binding, integration, and dissociation*. Oxford University Press; 2003.
 22. Farah MJ. *Visual agnosia: disorders of object recognition and what they tell us about normal vision*. Cambridge, MA: MIT Press; 1990.
 23. Heydrich L, Blanke O. Distinct illusory own-body perceptions caused by damage to posterior insula and extrastriate cortex. *Brain* 2013; 136:790–803.
- Based on a quantitative lesion analysis, this article suggests that heautoscopy is caused by a breakdown of self-other discrimination associated with damages in the left posterior insula.
24. Driver J, Noesselt T. Multisensory interplay reveals crossmodal influences on 'sensory-specific' brain regions, neural responses, and judgments. *Neuron* 2008; 57:11–23.
 25. Aspell JE, Heydrich L, Marillier G, et al. Turning body and self inside out: visualized heartbeats alter bodily self-consciousness and tactile perception. *Psychol Sci* 2013; 24:2445–2453.
 26. Ionta S, Heydrich L, Lenggenhager B, et al. Multisensory mechanisms in temporo-parietal cortex support self-location and first-person perspective. *Neuron* 2011; 70:363–374.
 27. Botvinick M, Cohen J. Rubber hands 'feel' touch that eyes see. *Nature* 1998; 391:756.
 28. Rohde M, Di Luca M, Ernst MO. The rubber hand illusion: feeling of ownership and proprioceptive drift do not go hand in hand. *PLoS One* 2011; 6:e21659.
 29. Lenggenhager B, Tadi T, Metzinger T, Blanke O. Video ergo sum: manipulating bodily self-consciousness. *Science* 2007; 317:1096–1099.
 30. Ehrsson HH. The experimental induction of out-of-body experiences. *Science* 2007; 317:1048.
 31. Moseley GL, Olthof N, Venema A, et al. Psychologically induced cooling of a specific body part caused by the illusory ownership of an artificial counterpart. *Proc Natl Acad Sci USA* 2008; 105:13169–13173.
 32. Salomon R, Lim M, Pfeiffer C, et al. Full body illusion is associated with widespread skin temperature reduction. *Front Behav Neurosci* 2013; 7:65.
- This study reveals that body temperature can be used as an objective signature of changes in bodily self-consciousness during the full body illusion.
33. Ionta S, Martuzzi R, Salomon R, Blanke O. The brain network reflecting bodily self-consciousness: a functional connectivity study. *Soc Cognit Affect Neurosci* 2014; 9:1904–1913.
- Extending previous brain imaging results [26], this study shows functional connectivity between the temporoparietal junction and the supplementary motor area, ventral premotor cortex, insula, intraparietal sulcus, and occipitotemporal cortex.
34. Petkova VI, Björnsdóttir M, Gentile G, et al. From part- to whole-body ownership in the multisensory brain. *Curr Biol* 2011; 21:1118–1122.
 35. Palluel E, Aspell JE, Blanke O. Leg muscle vibration modulates bodily self-consciousness: integration of proprioceptive, visual, and tactile signals. *J Neurophysiol* 2011; 105:2239–2247.
 36. Pfeiffer C, Serino A, Blanke O. The vestibular system: a spatial reference for bodily self-consciousness. *Front Integr Neurosci* 2014; 8:31.
 37. Apps M, Tsakiris M. The free-energy self: a predictive coding account of self-recognition. *Neurosci Biobehav Rev* 2014; 41:85–97.
- This article presents a Bayesian account for self-recognition, in which the integration of unimodal signals into multimodal areas explains away the surprising incoming sensory information from unimodal areas.

38. Seth A. Interoceptive inference, emotion, and the embodied self. *Trends Cognit Sci* 2013; 17:565–573.
 This opinion extends the predictive coding framework to interoception. It identifies emotional states with top-down inference of the causes of interoceptive signals, and link predictive models of the self with statistical correlation between interoceptive and exteroceptive signals.
39. Hay J, Ikeda K. Visual capture produced by prism spectacles. *Psychonomic Sci* 1965; 2:215–216.
40. Kennett S, Taylor-Clarke M, Haggard P. Noninformative vision improves the spatial resolution of touch in humans. *Curr Biol* 2001; 11:1188–1191.
41. Longo M, Cardozo S, Haggard P. Visual enhancement of touch and the bodily self. *Conscious Cognit* 2008; 17:1181–1191.
42. Pavani F, Spence C, Driver J. Visual capture of touch: out-of-the-body experiences with rubber gloves. *Psychol Sci* 2000; 11:353–359.
43. Aspell J, Lenggenhager B, Blanke O. Keeping in touch with one's self: multisensory mechanisms of self-consciousness. *PLoS One* 2009; 4:e6488.
44. Mudrik L, Faviere N, Koch C. Information integration without awareness. *Trends Cognit Sci* 2014; 18:488–496.
 This opinion article summarizes different views about the relationship between consciousness and information integration. By showing compelling evidence for the existence of unconscious integration across space, time, semantic domains, and sensory modalities, the authors conclude that a relation of both necessity and sufficiency between consciousness and integration is unlikely.
45. Meredith MA, Stein BE. Visual, auditory, and somatosensory convergence on cells in superior colliculus results in multisensory integration. *J Neurophysiol* 1986; 56:640–662.
46. Alsius A, Munhall K. Detection of audiovisual speech correspondences without visual awareness. *Psychol Sci* 2013; 24:423–431.
 This study shows that the detection of a face rendered invisible by continuous flash suppression is facilitated by the presentation of a matching vs. mismatching auditory sentence, hereby arguing for unconscious audiovisual integration.
47. Plass J, Guzman-Martinez E. Lip reading without awareness. *Psychol Sci* 2014; 25:1835–1837.
 Complementing the results from [46], this study shows that invisible lip movements facilitate the categorization of simultaneous congruent speech sounds, supporting further the existence of unconscious audiovisual integration.
48. Lunghi C, Binda P, Morrone M. Touch disambiguates rivalrous perception at early stages of visual analysis. *Curr Biol* 2010; 20:R143–R144.
49. Lunghi C, Morrone MC, Alais D. Auditory and tactile signals combine to influence vision during binocular rivalry. *J Neurosci* 2014; 34:784–792.
 In this study, rhythmic auditory and tactile stimuli are both shown to modulate the dominance of rivalrous visual stimuli, suggesting that visual activity can be influenced by congruent cross-modal signals with frequency selectivity through supramodal mechanisms.
50. Salomon R, Lim M, Herbelin B, et al. Posing for awareness: proprioception modulates access to visual consciousness in a continuous flash suppression task. *J Vision* 2013; 13:2.
 This study shows that the image of one's hands breaks continuous flash suppression faster when it matches the position of one's real hands, suggesting that the threshold for visual consciousness is modulated by proprioceptive signals.
51. Faviere N, Mudrik L, Schwartz N, Koch C. Multisensory integration in complete unawareness: evidence from audiovisual congruency priming. *Psychol Sci* 2014; 25:2006–2016.
 This study argues for the existence of audiovisual integration between two unconscious stimuli, by showing that two numbers conveyed by signals rendered invisible and inaudible by masking can nevertheless be compared up to a semantic level.
52. Lunghi C, Alais D. Touch interacts with vision during binocular rivalry with a tight orientation tuning. *PLoS One* 2013; 8:e58754.
 This study shows that the competition between two rival-oriented images presented to each eye is biased by simultaneous tactile signals with similar orientation.
53. Park H, Correia S, Ducorps A, Tallon-Baudry C. Spontaneous fluctuations in neural responses to heartbeats predict visual detection. *Nat Neurosci* 2014; 17:612–618.
 In this article, the neural signals evoked by heartbeats is shown to modulate the threshold for visual consciousness, suggesting the role of interoception in shaping visual perception.
54. Suzuki K, Garfinkel SN, Critchley HD, Seth AK. Multisensory integration across exteroceptive and interoceptive domains modulates self-experience in the rubber-hand illusion. *Neuropsychologia* 2013; 51:2909–2917.
 This study extends the visuotactile rubber-hand illusion to the interoceptive domain, by showing changes in limb ownership when one's hand glows in synchrony with heartbeat in augmented reality.
55. Clark B, Graybiel A. Factors contributing to the delay in the perception of the oculogravic illusion. *Am J Psychol* 1966; 79:377–388.
56. Lackner JR, Dizio P. Audiogravic and oculogravic illusions represent a unified spatial remapping. *Exp Brain Res* 2010; 202:513–518.
57. van Elk M, Blanke O. Imagined own-body transformations during passive self-motion. *Psychol Res* 2014; 78:18–27.
 This study demonstrates the role of the vestibular system in spatial perspective taking, by showing that imagined own-body rotations are facilitated when preceded by a rotation of the actual body with a congruent direction.
58. Gregory RL, Wallace JG, Campbell FW. Changes in the size and shape of visual after-images observed in complete darkness during changes of position in space. *Quar J Exp Psychol* 1959; 11:54–55.
59. Dieter K, Hu B, Knill D. Kinesthesia can make an invisible hand visible. *Psychol Sci* 2013; 25:66–75.
 This study shows that self-generated body movements in total darkness evoke visual sensations of motion, suggesting that the proprioceptive signals and/or motor commands that are highly predictive of a perceptual response are sufficient to evoke a conscious percept.
60. Christensen A, Ilg W, Giese M. Spatiotemporal tuning of the facilitation of biological motion perception by concurrent motor execution. *J Neurosci* 2011; 31:3493–3499.
61. Maruya K, Yang E, Blake R. Voluntary action influences visual competition. *Psychol Sci* 2007; 18:1090–1098.
62. Salomon R, Szpiro-Grinberg S, Lamy D. Self-motion holds a special status in visual processing. *PLoS One* 2011; 6:e24347.
63. Cohen MA, Cavanagh P, Chun MM, Nakayama K. The attentional requirements of consciousness. *Trends Cognit Sci* 2012; 16:411–417.
64. Koch C, Tsuchiya N. Attention and consciousness: two distinct brain processes. *Trends Cognit Sci* 2007; 11:16–22.
65. Lamme VAF. Why visual attention and awareness are different. *Trends Cognit Sci* 2003; 7:12–18.
66. Salomon R, Lim M, Kannape O, et al. 'Self pop-out': agency enhances self-recognition in visual search. *Exp Brain Res* 2013; 228:173–181.
 This study shows that when moving actively, a participant discriminate her own movements among virtual avatars independently of their number, and faster compared to a passive condition, revealing the role of agency in self-recognition.
67. Fleming S, Dolan R, Frith C. Metacognition: computation, biology and function. *Philos Trans Royal Soc B: Biol Sci* 2012; 367:1280–1286.