

HIV-1 Vpr and p21 restrict LINE-1 mobility

Koudai Kawano^{1,†}, Aurélien J. Doucet^{2,†}, Mikinori Ueno¹, Ryusho Kariya¹, Wenfeng An³, Flavia Marzetta⁴, Misao Kuroki¹, Priscilla Turelli⁴, Sayaka Sukegawa^{5,6}, Seiji Okada¹, Klaus Strebel⁵, Didier Trono⁴ and Yasuo Ariumi^{1,*}

¹Center for AIDS Research, Kumamoto University, Kumamoto 860-0811, Japan, ²Institute for Research on Cancer and Aging, Nice (IRCAN), INSERM U1081, CNRS UMR 7284, Université de Nice-Sophia-Antipolis, Faculté de Médecine, 06107 Nice Cedex 2, France, ³Department of Pharmaceutical Sciences, South Dakota State University, Brookings, SD 57007, USA, ⁴School of Life Sciences, Ecole Polytechnique Fédérale de Lausanne, Lausanne CH-1015, Switzerland, ⁵National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20892-0460, USA and ⁶Clinical Research Center, Nagoya Medical Center, Nagoya 460-0001, Japan

Received June 12, 2018; Revised July 13, 2018; Editorial Decision July 16, 2018; Accepted July 18, 2018

ABSTRACT

Long interspersed element-1 (LINE-1, L1) composes ~17% of the human genome. However, genetic interactions between L1 and human immunodeficiency virus type 1 (HIV-1) remain poorly understood. In this study, we found that HIV-1 suppresses L1 retrotransposition. Notably, HIV-1 Vpr strongly inhibited retrotransposition without inhibiting L1 promoter activity. Since Vpr is known to regulate host cell cycle, we examined the possibility whether Vpr suppresses L1 retrotransposition in a cell cycle dependent manner. We showed that the inhibitory effect of a mutant Vpr (H71R), which is unable to arrest the cell cycle, was significantly relieved compared with that of wild-type Vpr, suggesting that Vpr suppresses L1 mobility in a cell cycle dependent manner. Furthermore, a host cell cycle regulator p21^{Waf1} strongly suppressed L1 retrotransposition. The N-terminal kinase inhibitory domain (KID) of p21 was required for this inhibitory effect. Another KID-containing host cell cycle regulator p27^{Kip1} also strongly suppressed L1 retrotransposition. We showed that Vpr and p21 coimmunoprecipitated with L1 ORF2p and they suppressed the L1 reverse transcriptase activity in LEAP assay, suggesting that Vpr and p21 inhibit ORF2p-mediated reverse transcription. Altogether, our results suggest that viral and host cell cycle regulatory machinery limit L1 mobility in cultured cells.

INTRODUCTION

Long interspersed element-1 (LINE-1, L1) is an active and autonomous non-long terminal repeat (LTR) retrotranspo-

son composing ~17% of the human genome (1–3). L1 encodes two open reading frames (ORFs), ORF1p with RNA binding domain and nucleic acid chaperone activity, and ORF2p with endonuclease and reverse transcriptase activities required for its retrotransposition (1,2,4,5). L1 transcription occurs through promoter activity located in its 5'UTR (6). Several transcription factors including p53 (7), RUNX3 (8), SOX11 (9) and YY1 (10,11) positively regulate the L1 transcription (12). On the other hand, SRY (9) and SOX2 (13) negatively regulate the L1 transcription. L1 RNA assembles with ORF1p and ORF2p to form a ribonucleoprotein (RNP) complex in the cytoplasm (14). Then, L1-RNP complex enters the nucleus in which genomic integration occurs by a mechanism termed target-primed reverse transcription (TPRT). During TPRT, the L1 endonuclease creates a nicked DNA that serves as a primer for reverse transcription of L1 RNA, leading to integration of L1 cDNA into the human genome (15). Although L1 expression and retrotransposition can occur during early embryogenesis (16–18) and gametogenesis (18,19), L1 transcription is largely repressed by DNA methylation in somatic cells (19,20). In addition to the epigenetic control of L1 expression, L1 retrotransposition is controlled by several host restriction factors such as APOBEC3G (A3G), APOBEC3F (A3F) and MOV10 (12,21–27). A3G was first identified as anti-human immunodeficiency virus type 1 (HIV-1) restriction factor (28) and HIV-1 restriction requires A3G cytidine deaminase activity (29,30). A3G restricts exogenous retroviruses, hepatitis B virus (HBV), and endogenous retroelements, such as L1, Alu, SVA and HERVs (21,29,31–34). However, the A3G cytidine deaminase activity is dispensable for L1 restriction. Escape of these control pathways can lead to *de novo* L1 retrotransposition in somatic cells that could contribute to mutagenesis and genomic instability leading to cancer (35–38). L1 retrotransposition can also generate mutations of genes in the germ line or during devel-

*To whom correspondence should be addressed. Tel: +81 96 373 6526; Fax: +81 96 373 6526; Email: ariumi@kumamoto-u.ac.jp

†The authors wish it to be known that, in their opinion, the first two authors should be regarded as Joint First Authors.

opment that could contribute to diseases (39,40). Therefore, L1 must be regulated during normal development.

HIV-1 is a retrovirus, which encodes three structural proteins, group-specific antigen (Gag), polymerase (Pol), and envelope (Env), two regulatory proteins, Tat and Rev, and four accessory proteins, Vif, Vpr, Vpr and Nef. The gene expression of HIV-1 is transcriptionally regulated by Tat through its binding to a nascent HIV-1 *trans*-activation responsive (TAR) RNA (41,42), and post-transcriptionally by Rev through its interaction with Rev-responsive element (RRE) RNA in the *env* gene (43–45). Rev forms a complex with CRM1-Ran-GTP and enhances the nuclear export of HIV-1 mRNA (43–45). In addition, several host DEAD-box RNA helicases cooperate to modulate HIV-1 Rev function (46–50). HIV-1 Vpr is a virion-associated nuclear protein with multiple functions (51,52). Vpr facilitates HIV-1 infection of nondividing cells by regulating the nuclear export of the HIV-1 pre-integration complex (PIC). Vpr also induces cell cycle arrest at the G₂ phase in proliferating infected cells and stimulates the LTR-directed gene expression (53). Following HIV-1 entry, its own reverse transcriptase synthesizes a DNA copy of the HIV-1 genomic RNA. Integration of a DNA copy of the viral RNA genome is a crucial step in the life cycle of HIV-1. Therefore, both HIV-1 and L1 might mutually influence their mobility. However, interactions between HIV-1 and L1 are not well understood. Therefore, we investigated a cross talk of HIV-1 with L1 in this study.

MATERIALS AND METHODS

Cell culture

293T, TET293T, P4.2 and TZM-bl cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Life Technology, Carlsbad, CA, USA) with high glucose (4.5 g/l) supplemented with 10% fetal bovine serum (FBS) and 100 U/ml penicillin/streptomycin. Details of specific transfection conditions for each experiment are provided in the figure legends.

Plasmid construction

To construct pcDNA3-HA-ORF1 or pcDNA3-ORF1-HA, a DNA fragment encoding ORF1p was amplified from pEGFP-L1_{RP} wt (54) by PCR using KOD-Plus DNA polymerase (TOYOBO, Osaka, Japan) and the following pairs of primers: HA-ORF1, 5'-CGGGATC CAAGATGGGGAAAAACAGAACA-3' (Forward), 5'-CCG GCGGCCGCTTACATTTTGGCATGATTT-3' (Reverse); ORF1-HA, 5'-CG GGATCCAAGATG GGGAAAAACAGAACA-3' (Forward), 5'-CCG GCGGCCGCTTAAAGAAGTCTCCAGGCTGGC ATAGTCAGGCACGTCATAAGGATAGCTAGAAG CCATCATTTTGGCATGATTTT-3' (Reverse). The obtained DNA fragments were subcloned into either the BamHI-XhoI sites of the pcDNA3-HA vector or the BamHI-NotI sites of the pcDNA3 vector (Invitrogen), and the nucleotide sequences were determined by Sanger sequencing. To construct pcDNA3-FLAG-ORF1, the DNA fragments encoding ORF1 obtained by digestion with BamHI and XhoI were subcloned

into the BamHI-XhoI sites of the pcDNA3-FLAG vector. To construct pcDNA3-HA-Vpr WT, pcDNA3-HA-Vpr H71R, or pEGFP-Vpr, a DNA fragment encoding HIV-1 Vpr was amplified from pCMX Vpr96 by PCR using KOD-Plus DNA polymerase and the following pairs of primers: HA-Vpr WT, 5'-CGGGATC CAAGATGGAACAAGCCCCAGAAG-3' (Forward), 5'-CCGCGCGCCGCTAGGATCTACTGGCTCCA-3' (Reverse); HA-Vpr H71K, 5'-CGGGATCCAAGAT GGAACAAGCCCCAGAAG-3' (Forward), 5'-CCG GCGGCCGCTAGGATCTACTGGCTCCATTCT TGCTCTCCTCTGTGCGAGTAACGCCTATTCTGCTA TGTCGACACCCAATTCTGAAACGGATAAACA-3' (Reverse); EGFP-Vpr, 5'-CGCTCGAGAAG ATGGAACAAGCCCCAGAAG-3' (Forward), 5'-CGGATCCCTAGGATCTACTGGCTCCATT-3' (Reverse). The obtained DNA fragments were subcloned into the BamHI-NotI sites of the pcDNA3-HA vector or the XhoI-BamHI sites of the pEGFP-C3 (Promega). To construct pAIP-Vpr WT or pAIP-Vpr H71R, a DNA fragment encoding Vpr WT or H71R was amplified from pcDNA3-HA-Vpr WT or pcDNA3-HA-Vpr H71R by TKs Gflex DNA polymerase (TAKARA) and the following pair of primers: 5'-TCTCTCCCCATCTAGATGG AACAAGCCCCAGAAGAC-3' (Forward), 5'-GCGTT TAAACGGATCCTAGGATCTACTGGCTCCATT-3' (Reverse). The obtained DNA fragments were subcloned into the BamHI-XbaI sites of pAIP (55) by using In-Fusion Cloning kit (Clontech). pcDNA3p21 and p21N were kindly provided by Dr. Dutta (56). To construct pcDNA3p21C, a DNA fragment encoding p21C was amplified from pcDNA3p21 by PCR using KOD-Plus DNA polymerase and the following pairs of primers: 5'-CGGGA TCCAAGATGCGAGGCCGGGATGAGT-3' (Forward), 5'-CCGCTCGAGTTAGGGCTTCTCTTGGAGAA-3' (Reverse). The obtained DNA fragments were subcloned into the BamHI-XhoI sites of the pcDNA3-HA vector.

RNA interference

Oligonucleotides with the following sense and antisense sequences were used for the cloning of short hairpin RNA (shRNA)-encoding sequences targeted to p21 in a lentiviral vector: 5'-GATCCCCGACCATGTGGACCTGTCCT TCAAGAGAGTGACAGGTCCACATGGTCTTTTT GGAAA-3' (sense), 5'-AGCTTTTCCAAAAGACCA TGTGGACCTGTCCTCTTTGAAGTGACAGGT CCACATGGTTCGGG-3' (antisense). The oligonucleotides above were annealed and subcloned into the BglII-HindIII sites, downstream from an RNA polymerase III promoter of pSUPER (57), to generate pSUPER-p21i. To construct pLV-p21i, the BamHI-SalI fragment of the pSUPER-p21i was subcloned into the BamHI-SalI sites of pRDI292, an HIV-1-derived self-inactivating lentiviral vector containing a puromycin resistance marker allowing for the selection of transduced cells (58).

Lentiviral vector production

The vesicular stomatitis virus (VSV)-G-pseudotyped HIV-1-based vector system has been described previously

(59,60). The lentiviral vector particles were produced by transient transfection of the second-generation packaging construct pCMV Δ R8.74 (59,60) and the VSV-G-envelope-expressing plasmid pMD2G as well as lentiviral vector pLV-p21i or pAIP (55) into 293T cells with FuGENE6 (Promega, Madison, WI, USA) or TransIT-LT1 (Mirus Bio LLC, Madison, WI 53711, USA) transfection reagents.

HIV-1 infection

Different full-length HIV-1 molecular clones used for transfection were as follows: pR9 (59,60), pNL4-3 (61), pJR-FL (62) and pAD8 (63). Recombinant HIV-1 was prepared as described previously (64). Briefly, 293T cells were used as viral producer cells. The cells were seeded at 2×10^5 cells per well into six-well tissue culture plates and transfected the next day using TransIT-LT1 reagent with one of the following HIV-1 molecular clones: pR9, pNL4-3, pJR-FL or pAD8. Culture media were replaced with fresh media 24 h after transfection, and the cells were cultured for an additional 48 h. Then, the supernatants containing recombinant viruses were clarified by centrifugation and filtration with 0.45 μ m filter (Kurabo, Osaka, Japan), and stored at -80°C before use. Released HIV-1 virions were collected by centrifugation of the culture supernatants from transfected 293T cells at $20\,000 \times g$ for 2 h at 4°C . The pellets containing virions were dissolved in the lysis buffer and then subjected to western blot. HIV-1 infection was performed as described previously (64). P4.2, CD4⁺ HeLa cells were incubated with the supernatants of 293T cells containing HIV-1 at the indicated multiplicity of infection (MOI), and the cells were cultured for an additional 48h.

Western blot analysis

Cells were lysed in buffer containing 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 4 mM EDTA, 1% Nonidet (N) P-40, 0.1% sodium dodecyl sulfate (SDS), 1 mM dithiothreitol (DTT) and 1 mM phenylmethylsulfonyl fluoride (PMSF). Supernatants from these lysates were subjected to SDS-polyacrylamide gel electrophoresis, followed by immunoblot analysis using anti-HIV-1 p24 (65-005; Bioacademia, Osaka, Japan), anti-p21^{Waf1/Cip1} (12D1) (#2947; Cell Signaling Technology, Inc., Danvers, MA 01923, USA), anti-p21 [EPR362] (ab 109520; Abcam), anti- β -actin (A5441; Sigma, Saint Louis, MI, USA), anti-HA (HA-7; Sigma), anti-HA (12CA5; Santa Cruz), anti-FLAG antibody (M2; Sigma), anti-S6 ribosomal protein (5G10; Cell Signaling Technology), anti-GAPDH (GA1R, Pierce Biotechnology), anti-hrGFP (#240141, Agilent Technologies), or anti-hORF1p (SE-6798) (65) as primary antibodies. We used peroxidase-conjugated Donkey anti-Rabbit IgG (H+L) (Jackson ImmunoResearch), Amersham ECL peroxidase-linked Sheep anti-Mouse IgG (GE Healthcare), IRDye 800CW Goat anti-Rabbit IgG, IRDye 800CW Goat anti-Mouse IgG, IRDye 680RD Goat anti-Rabbit IgG or IRDye 680RD Goat anti-Mouse IgG (LI-COR Biosciences) as secondary antibodies.

Luciferase assay

293T cells were seeded in 24-well plates at 2×10^4 cells per plate and then transfected the next day with plasmids using FuGENE6 (Promega, Madison, WI, USA) or TransIT-LT1 transfection reagent. Luciferase assays were performed 72 h after transfection using luciferase assay reagent according to the manufacturer's instructions (Promega). All transfections were performed using equal number of tested plasmid DNA molecules, with the addition of empty vector into the transfection mixtures to compensate for plasmid size differences and reach equal amounts of DNA quantities per condition. Results were obtained through three independent transfections. A Lumat LB9507 luminometer (Berthold, Bad Wildbad, Germany) was used to measure luciferase activities.

Cell cycle analysis

For cell cycle analysis, cells were fixed in 70% ethanol at 4°C overnight and stained with propidium iodide (PI, 10 μ g/ml) (66). The DNA content in each cell was analyzed on LSRII flow cytometer (BD Bioscience, San Jose, CA). Data were analyzed on Flowjo software (Tree Star, San Carlos, CA)

Immunoprecipitation

Cells were lysed in buffer containing 10 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM EDTA, 0.1% NP-40, 10 mM NaF, 1 mM DTT and 1 mM PMSF. Lysates were pre-cleared with 30 μ l of protein-G-Sepharose (GE Healthcare Bio-Sciences, Uppsala, Sweden). Pre-cleared supernatants were incubated with 5 μ g of either anti-HA antibody (3F10; Roche Diagnostics, Mannheim, Germany) or anti-FLAG antibody (M2, Sigma) at 4°C for 1 h. Following absorption of the precipitates on 30 μ l of protein-G-Sepharose resin for 1 h, the resin was washed four times with 700 μ l of lysis buffer. Proteins were eluted by boiling the resin for 5 min in $2 \times$ Laemmli sample buffer. The proteins were then subjected to SDS-PAGE, followed by immunoblot analysis.

Immunofluorescence and confocal microscopy analysis

293T cells were grown on Lab-Tek 2 well chamber slide (Nunc, Thermo) at 2×10^4 cells per well and transfected the next days using TransIT-LT1 transfection reagent. Two days after transfection, the cells were fixed in 3.6% formaldehyde in $1 \times$ phosphate-buffered saline (PBS), permeabilized in 0.1% NP-40 in $1 \times$ PBS at room temperature, and incubated with anti-HA antibody (HA-7) at a 1:300 dilution in $1 \times$ PBS containing 3% bovine serum albumin (BSA) at 37°C for 30 min. Cells were then stained with Cy3-conjugated anti-mouse antibody (Jackson ImmunoResearch, West Grove, PA, USA) at a 1:300 dilution in $1 \times$ PBS containing 3% BSA at 37°C for 30 min. Nuclei were stained with DAPI (4',6'-diamidino-2-phenylindole). Following washing three times in $1 \times$ PBS, the coverslips were mounted on slides using SlowFade Gold antifade reagent (Life Technology). Samples were analyzed under a confocal laser-scanning microscope (FV1200; Olympus, Tokyo, Japan).

Real-time RT-PCR

Total RNA was isolated using RNeasy Mini kit (Qiagen) and treated with Turbo DNA-free kit (Life Technology). The total RNA was reverse transcribed with oligo(dT)12-18 primer (Life Technology). The quantitative RT-PCR analysis for LINE-1 mRNA was performed by real-time Light-Cycler PCR (Roche). We used the following forward and reverse primer sets:

5UTR L1 Hs, GGGAGGAGGAGCCAAGATG (Forward), ACATGGCCTGCGCCACTG (Reverse); ORF1 L1 Hs, AAAACGCAGAGCGCCTCTCC (Forward), GTTTGAATGTCCTCCCGTAG (Reverse); ORF1 L1PA, GAACGCCACAAAGATACTCC (Forward), CTCTTCTGGCTTGTTAGGGTTTCTG (Reverse); *β-actin*, TGACGGGGTCACCCACACTG (forward), AAGCTGTAGCCGCGCTCGGT (reverse).

Purification of L1 RNPs

L1 RNPs were prepared as previously described with minor modifications (65,67). Briefly, 293T cells were seeded in 10 cm plates at 3×10^6 cells per plate and then transfected the next days with 24 μg of plasmid using the calcium phosphate-mediated method (68). Culture medium was changed 5h later. Cells were collected 5 days post-transfection by trypsinization and were then lysed in CHAPS extract buffer containing 10 mM Tris-HCl pH7.5, 0.5% CHAPS (w/v), 1 mM MgCl_2 , 1 mM EGTA, 10% glycerol, protease inhibitors cocktail (Roche) and 1 mM DTT. After incubation at 4°C for 15 min, cell debris was removed by centrifugation at $20\,000 \times g$ for 15 min. A sucrose cushion was prepared with 8.5% and 17% (w/v) sucrose in 20 mM Tris-HCl pH 7.5, 80 mM NaCl, 5 mM MgCl_2 , 1 mM EGTA, protease inhibitors cocktail (Roche) and 1 mM DTT. For each condition, 1 ml of cell lysate was loaded on a sucrose cushion. Cellular RNPs were pelleted by ultracentrifugation at $178\,000 \times g$ at 4°C for 2 h. The pellets were resuspended in 50 μl of H_2O . Quantification was performed using the Bradford assay (Bio-rad). RNP fractions were diluted at 1 $\mu\text{g}/\mu\text{l}$ in H_2O and aliquoted before use or storage at -80°C .

L1 element amplification protocol (LEAP) assay

LEAP assay was performed as previously described (65,67,69). Briefly, L1 reverse transcription was carried out at 37°C for 1 h in 50 μl reaction mixture containing 0.75 μg of RNPs, 50 mM Tris-HCl pH 7.5, 50 mM KCl, 5 mM MgCl_2 , 0.05% (v/v) Tween-20, 10 mM DTT, 20 U RNasin (Promega), 0.2 mM dNTPs and 0.4 μM RACE primer (5'-GCGAGCACAGAATTAATACGACTCACTATAGGTTTTTTTTTTTTTTVN-3').

1 μl of the LEAP reaction were PCR-amplified in 50 μl reactions containing 0.2 mM dNTPs, 3 mM MgCl_2 , 10 pmol of primers LOU312 (5'-GCGAGCACAGAATTAATACGACT-3'), LOU851 (5'-GGGTTTCGAAATCGATAAGCTTGGATCCAGAC-3') and 2 U of Platinum Taq Polymerase (Life technologies). A first step at 94°C for 2 min was followed by 35

cycles at 94°C for 30 s, at 60°C for 30 s and at 72°C for 30 s and by a final step at 72°C for 5 min. The PCR products were analyzed by 2% agarose gel electrophoresis in $0.5 \times \text{TBE}$ buffer.

RNA extraction and RT-PCR

Total RNA was extracted from 15 μg of L1 RNP fraction using TRI reagent (Molecular Research Center Inc). DNase treatment was performed at 37°C for 30 min in a 10 μl reaction mixture containing 1 μg of extracted RNA, 1 U of RQ1 RNase-free DNase (Promega), 40 mM Tris-HCl pH 8.0, 10 mM MgSO_4 and 1 mM CaCl_2 . Then, cDNA synthesis was performed at 50°C for 1 h in 20 μl reaction mixture containing 6 μl of DNase-treated RNA, 50 pmol of RACE primer, 10 mM dNTPs, 40 U RNaseOUT recombinant ribonuclease inhibitor (Life Technologies), 200 U of SuperScript III Reverse Transcriptase (Life technologies), 50 mM Tris-HCl pH8.0, 75 mM KCl, 3 mM MgCl_2 and 5 mM DTT. After the first-strand cDNA synthesis, PCR amplification was performed as described above but with only 30 cycles of amplification and using primers LOU312 and LOU851 for L1, and LOU1071 (5'-GTTTTCCATTGGCTTGTA GATC-3') and LOU1072 (5'-GAAGATGGTGATGGG ATTTC-3') for GAPDH.

Retrotransposition assays

LINE-1 retrotransposition assays consist of the expression in cells of a retrotransposition-competent L1 element equipped with an indicator cassette for which the reporter gene is expressed only after a cycle of retrotransposition (54,70–72). The indicator cassette consists of a reporter gene (e.g. EGFP, firefly luciferase) with its own promoter and polyadenylation signal. It is introduced in the 3'UTR of an active L1, in the opposite transcriptional orientation compared to L1 (Figure 1A and B). The reporter gene is interrupted by an intron with splice donor and splice acceptor sites in the transcriptional orientation of the L1. Therefore, the only way to express the reporter gene is when L1 RNA is expressed from its promoter, splicing occurs to remove the intron, and then retrotransposition takes place, leading to the insertion at a genomic locus of the reporter gene. In absence of the original intron, the reporter gene can now be expressed from the integration site.

The EGFP-based retrotransposition assay was performed as followed: 293T cells were seeded in 12-well plates at 5×10^4 cells per plate and transfected the next day using FuGENE6 with 500 ng of either an EGFP-based retrotransposition reporter pL1_{RP}-EGFP (54) or its corresponding negative control pL1_{RP}(JM111)-EGFP which contains two missense mutations (ARR at residues 260–262) in the ORF1 coding region known to abolish retrotransposition (54). Three days after transfection, retrotransposition efficiency was determined by GFP fluorescence-activated cell sorter (FACS) analysis with Guava easyCyte flow cytometer (Merk Millipore, Billerica, MA, USA).

The luciferase-based retrotransposition assay was performed using the firefly luciferase-based retrotransposition reporter pYX014 plasmid (72). pYX014 is a dual-luciferase




Figure 1. Suppression of L1 retrotransposition by HIV-1. (A) Schematic representation of the EGFP-based retrotransposition reporter cassette, pL1_{RP}-EGFP (54). The EGFP reporter is introduced in the 3'UTR of L1, in opposite orientation. It is also interrupted by an intron with splice donor (SD) and splice acceptor (SA) in the transcriptional orientation of the L1. 293T cells (5×10^4 cells/well) were co-transfected with HIV-1 molecular clone (R9, NL4-3 or JR-FL) (500 ng) and either pL1_{RP}-EGFP or its mutant pL1_{RP}(JM111)-EGFP lacking the ability of retrotransposition (500 ng). Three days after transfection, retrotransposition efficiency was determined by GFP fluorescence-activated cell sorter (FACS) analysis with Guava easyCyte flow cytometer (Merk Millipore, Billerica, MA, USA). Experiments were done in triplicate and graph shows the mean (\pm SEM) percentage of GFP positive cells ($*P < 0.05$). (B) Schematic representation of the firefly luciferase-based retrotransposition reporter cassette, pYX014 (72). The pYX014 plasmid is a dual-luciferase reporter, in which firefly luciferase is used as the retrotransposition indicator and *Renilla* luciferase is encoded on the same plasmid for normalization. 293T cells (2×10^4 cells/well) were co-transfected with HIV-1 molecular clone (R9, NL4-3, or JR-FL) (100 ng) and pYX014 (100 ng) in triplicate. Three days after transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity normalized with *Renilla* luciferase activity with the condition without HIV-1 set to 100% ($*P < 0.05$). (C, D) Suppression of L1 retrotransposition by HIV-1 infection. P4.2 cells (2×10^4 cells/well) were infected with either HIV-1 strain of R9 (C) or NL4-3 (D) at the indicated multiplicity of infection (MOI). 24h post-infection, the HIV-1-infected cells were transfected with pYX014 (200 ng) in triplicate. Two days post-transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity normalized with *Renilla* luciferase activity in Relative Luminometer Units ($*P < 0.05$).

reporter in which firefly luciferase is used as the retrotransposition indicator and *Renilla* luciferase is encoded on the same plasmid backbone for normalization. 293T cells were seeded in 24-well plates at 2×10^4 cells per plate and then transfected the next day with pYX014 (100 ng) using TransIT-LT1 transfection reagent. Dual luciferase assays were performed 72 h after transfection using luciferase assay reagent according to the manufacturer's instructions (Promega). A Lumat LB9507 luminometer (Berthold, Bad Wildbad, Germany) was used to measure both firefly and *Renilla* luciferase activities. Firefly luciferase activity was normalized with *Renilla* luciferase activity.

Statistical analysis

The statistical significance of the inter-sample differences was determined using the paired Student's *t* test. *P* values of <0.05 were considered statistically significant.

RESULTS

HIV-1 Vpr suppresses L1 retrotransposition

To investigate the potential role of HIV-1 expression in L1 retrotransposition, three different HIV-1 molecular clones (R9, NL4-3 and JR-FL, respectively) (59–62) were co-transfected in 293T cells with pL1_{RP}-EGFP plasmid (Figure 1A), which contains an EGFP-based retrotransposition reporter cassette in the antisense orientation (54). The presence of the intron in the opposite transcriptional orientation of the reporter prevents EGFP expression from the plasmid. GFP positive cells can only arise after expression, splicing and retrotransposition of the labeled L1_{RP} copy into new genomic location. The use of a retrotransposition-deficient L1 expressing plasmid, pL1_{RP}(JM111)-EGFP, serves as negative control of the assay (54). We analyzed by flow cytometry the level of GFP expression and showed that

all HIV-1 clones could strongly suppress the retrotransposition of L1 (Figure 1A). To confirm these findings, we also assayed retrotransposition using the pYX014 plasmid, which has a firefly luciferase-based retrotransposition reporter cassette (72). The pYX014 plasmid is a dual-luciferase reporter in which firefly luciferase is used as the retrotransposition indicator and *Renilla* luciferase is encoded on the same plasmid backbone for normalization (Figure 1B). Similar to the EGFP reporter, the firefly luciferase activity can only be detected once L1 retrotransposition occurred. Again, all HIV-1 clones could strongly suppress the luciferase-based L1 retrotransposition activity (Figure 1B). We next examined whether HIV-1 infection affects L1 retrotransposition activity. CD4⁺HeLa P4.2 cells were infected with HIV-1 (R9 or NL4-3) at the indicated multiplicity of infection (MOI) (Figure 1C and D). The next day, the luciferase-based retrotransposition reporter plasmid pYX014 was transfected into the HIV-1-infected cells. Two days post-transfection, we measured luciferase activities and observed a three-fold reduction of L1 retrotransposition at the higher MOI for both R9 and NL4-3, indicating that HIV-1 infection also modulates L1 mobility (Figure 1C and D). Since we did not infect equalized viral input (p24 amount) of HIV-1 between R9 and NL4-3 strains, the suppression of L1 retrotransposition seems to vary based on the MOI (Figure 1C and D). Subsequently, we used individual HIV-1 protein-expressing plasmids to determine which of the following HIV-1 protein suppresses L1 retrotransposition: Vpr, Tat, Rev, integrase (IN), Env and Nef. Using the EGFP-based retrotransposition reporter, we showed that only Vpr could strongly suppress L1 retrotransposition (Figure 2A). We also noticed that Tat weakly suppress L1 retrotransposition (Figure 2A). Moreover, Vpr suppressed L1 retrotransposition activity in a dose-dependent manner (Figure 2B). Likewise, Vpr could strongly suppress the luciferase-based L1 retrotransposition activity (Figure 2C). To test if Vpr is affecting the L1 promoter activity, we used pYX017 plasmid in which the L1_{RP} 5'UTR (present in pYX014) is replaced by a strong CAG promoter resulting in an increased retrotransposition activity (72) (Figure 2D). Despite the change of promoter, Vpr can also suppress L1 retrotransposition when expressed from pYX017 (Figure 2D), suggesting that Vpr does not affect the endogenous L1 promoter activity. Since HIV-1 Vpr is known to regulate the host cell cycle by arresting cells at G₂ phase (53), we examined whether Vpr suppresses L1 retrotransposition in a cell cycle dependent manner. To this end, we used a mutant Vpr (H71R) unable to arrest cell cycle (53). Consistent with the previous report, we confirmed that wild-type Vpr (WT) arrested cell cycle at G₂ (Figure 3A and B), while the H71R mutant failed to arrest cell cycle (Figure 3C). As a control, we observed similar levels of protein expression for both Vpr (WT) and the mutant Vpr (H71R) in transfected cells (Figure 3D). Although Vpr WT suppresses L1 retrotransposition activities from both pYX014 and pYX017, the inhibitory effect of Vpr (H71R) was significantly relieved compared with that of Vpr (WT) (Figure 3E and F), suggesting that Vpr suppresses L1 mobility in a cell cycle dependent manner.




Figure 2. Suppression of LINE-1 retrotransposition by HIV-1 Vpr. (A) 293T cells (5×10^4 cells/well) were co-transfected with each HIV-1 protein-expressing plasmid, Vpr, Tat, Rev, IN, Env or Nef (500 ng), and either pL1_{RP}-EGFP or pL1_{RP}(JM111)-EGFP (500 ng). Three days after transfection, retrotransposition efficiency was determined by GFP FACS analysis. Experiments were done in triplicate and graph shows the mean (\pm SEM) percentage of GFP positive cells ($*P < 0.05$). (B) Suppression of LINE-1 retrotransposition by HIV-1 Vpr in a dose-dependent manner. 293T cells (5×10^4 cells/well) were co-transfected with the indicated amounts (ng) of pCMX-Vpr96 and either pL1_{RP}-EGFP or pL1_{RP}(JM111)-EGFP (500 ng). Relative retrotransposition efficiency (fold) among three replicates (\pm SEM) is shown ($*P < 0.05$). (C, D) 293T cells (2×10^4 cells/well) were co-transfected with pCMX-Vpr96 (100 ng) and either pYX014 (C) or pYX017 (100 ng) (D) in triplicate. Three days after transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity normalized with *Renilla* luciferase activity with the condition without Vpr set to 100% ($*P < 0.05$).

p21^{Waf1} and p27^{Kip1}, host cell cycle regulators, suppress L1 retrotransposition

Since HIV-1 Vpr restricted L1 retrotransposition in a cell cycle dependent manner (Figure 3), we examined whether host cell cycle regulators, such as p21^{Waf1/Cip1} (p21) and p27




Figure 3. Suppression of L1 retrotransposition by Vpr in a cell cycle dependent manner. (A–C) Cell cycle analysis of 293T cells expressing Vpr WT (B) or Vpr H71R. (C). Cells were fixed in 70% ethanol three days post-transduction and stained with propidium iodide (PI) to perform cell cycle analysis by FACS. (D) Protein expression levels of wild-type Vpr (WT) and a mutant Vpr (H71R). 293T cells (2×10^5 cells/well) were transfected with 2 μ g of pcDNA3-HA-Vpr WT or pcDNA3-HA-Vpr H71R. Western blotting of cell lysates with anti-HA and anti- β -actin antibodies is shown. (E, F) 293T cells (2×10^4 cells/well) were co-transfected with pYX014 (E) or pYX017 (100 ng) (F) and either pcDNA3-HA-Vpr WT or pcDNA3-HA-Vpr H71R (100 ng) in triplicate. Three days after transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity normalized with *Renilla* luciferase activity in Relative Luminometer Units (* $P < 0.05$).

Kip1 (p27) (73–77) might suppress L1 mobility. To test this possibility, several plasmids expressing tumor suppressor-related proteins were independently co-transfected with pL1_{RP}-EGFP plasmid into 293T cells. GFP fluorescence was then analyzed by flow cytometry to monitor the impact on L1 retrotransposition. Notably, p21 strongly suppressed L1 retrotransposition (Figure 4A). p53 and p73, which transcriptionally induce p21, also significantly suppressed L1 retrotransposition (Figure 4A). p21 has an N-terminal kinase inhibitory domain (KID) and a C-terminal PCNA-binding domain (Figure 4B) (56). Therefore, we used the N-terminal half (p21N) or the C-terminal half of p21 (p21C) to determine which domain is important for an inhibitory effect on L1 retrotransposition (Figure 4B). Our results showed that p21N but not p21C strongly suppressed L1 retrotransposition indicating that the kinase inhibitory domain is important for the inhibitory effect (Figure 4C and D). In this regard, p21N is known to inhibit cell growth whereas p21C does not (56). Consistent with this result, p27 harboring the kinase inhibitory domain also strongly suppressed L1 retrotransposition (Figure 4C and D). To further confirm this observation, we established a p21 knock-down 293T cell line using lentiviral vector stably expressing shRNA targeted to p21 (p21KD in Figure 4E). As expected, the L1 retrotransposition efficiency was elevated in p21 knockdown 293T cells (Figure 4F and G), even though we observed no significant effect of p21 knockdown on the 293T cell growth. Thus, host cell cycle inhibitors p21 and p27 restrict L1 mobility.

Incorporation of L1 ORF1p into HIV-1 virions

Since it has been indicated that several P-body and stress granule components, such as APOBEC3G, APOBEC3F and MOV10, are incorporated into HIV-1 virions and affect HIV-1 infectivity or production (78–82), we examined whether ORF1p is incorporated into HIV-1 virions and modifies the HIV-1 infectivity. Notably, we noticed that HA-tagged ORF1p (either in N- or C-terminus) are incorporated into the HIV-1 virions (NL4-3 and JR-FL), respectively (Figure 5A). We also found that the incorporation of ORF1p into HIV-1 virions was independent of HIV-1 accessory proteins including Nef, Vpr, Vpu and Vif, because the mutant viruses lacking the expression of each viral gene still retained the ability to incorporate ORF1p (Figure 5B). However, we failed to detect ORF2p and L1 RNA in HIV-1 virion (data not shown). On the other hand, ORF1p affected neither the intracellular nor extracellular HIV-1 p24 protein levels, suggesting that ORF1p does not negatively impact HIV-1 replication (Figure 5A). Furthermore, the incorporated ORF1p-HA did not drastically affect the HIV-1 infectivity, as measured by luciferase assay with two independent infectious HIV-1 molecular clones (NL4-3 and JR-FL) (61,62) (Figure 5C and D). However, we noticed a modest suppressive effect of HA-ORF1 on the infectivity of NL4-3 but not JR-FL. Since ORF1p is an RNA-binding and nucleic acid chaperone protein, we also examined whether ORF1p modulates the HIV-1 Tat-mediated HIV-1 transcription using the HIV-1 LTR Luc reporter assay (83–85) or HIV-1 Rev-dependent nuclear ex-




Figure 4. p21 and p27 suppress L1 retrotransposition. (A) Potential effect of tumor suppressor proteins on L1 retrotransposition. 293T cells (5×10^4 cells/well) were co-transfected with each plasmid expressing tumor suppressor-related protein, p53 (115), p73 α (116,117), p73 β (116,117), p21 (56,118), Chk2 (119,120), Rb, ATM (121,122), or ATR (500 ng), and either pL1_{RP}-EGFP or pL1_{RP}(JM111)-EGFP (500 ng). Three days after transfection, retrotransposition efficiency was determined by GFP FACS analysis. Experiments were done in triplicate and graph shows the mean (\pm SEM) percentage of GFP positive cells ($*P < 0.05$). (B) Schematic representation of p21, p21 mutants (p21N and p21C), and p27. KID: kinase inhibitory domain, PCNA: PCNA-binding domain, QT: glutamine threonine-rich domain. Protein expression levels of p21, p21N, p21C, and p27. 293T cells (2×10^5 cells/well) were transfected with pcDNA3, pcDNA3p21 (56), pcDNA3p21N (56), pcDNA3p21C or pCMXp27. Western blotting of the cell lysates with anti-p21, anti-p27, or anti- β -actin antibodies is shown. (C, D) KID is required for inhibitory effect on L1 retrotransposition. 293T cells (2×10^4 cells/well) were co-transfected with pcDNA3p21, pcDNA3p21N, pcDNA3p21C or pCMXp27 (100 ng) and either pYX014 (C) or pYX017 (D) (100 ng) in triplicate. Three days after transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity normalized with *Renilla* luciferase activity in Relative Luminometer Units ($*P < 0.05$). (E) Inhibition of p21 expression by shRNA-producing lentiviral vector. Western blotting of cell lysates with anti-p21 or anti- β -actin antibodies is shown. (F, G) Enhancement of L1 retrotransposition efficiency in p21 knockdown cells. Control (ctrl) or p21 knockdown (p21KD) 293T cells (2×10^4 cells/well) were transfected with either pYX014 (F) or pYX017 (G) (100 ng) in triplicate. 24h after transfection, luciferase assays were performed, and results were plotted as described above ($*P < 0.05$).




Figure 5. Incorporation of L1 ORF1p into HIV-1 virions. (A) 293T cells (2×10^5 cells/well) were co-transfected with 2 μ g of pcDNA3-HA-ORF1 or pcDNA3-ORF1-HA and 2 μ g of pNL4-3 or pJR-FL. Western blotting of cell lysates or virion in the concentrated culture supernatants with anti-HA, anti-HIV-1 p24, or anti- β -actin antibodies is shown. (B) HIV-1 accessory protein-independent incorporation of ORF1p into HIV-1 virions. 293T cells (2×10^5 cells/well) were co-transfected with 2 μ g of HIV-1 AD8 wild-type molecular clone (WT) (63) or HIV-1 mutants (Δ Vif, Δ Vpr, Δ Vpu, or Δ Nef) and 2 μ g of pcDNA3-ORF1-HA. HIV-1 virions were then produced in the culture supernatants. Western blotting of HIV-1 in the concentrated culture supernatants with either anti-HA or anti-HIV-1 p24 antibody is shown. (C, D) Incorporation of ORF1p does not affect the HIV-1 infectivity. TZM-bl cells (1×10^5 cells/well) were inoculated with 100 μ l of the culture supernatants of 293T cells co-transfected with pNL4-3 (C) or pJR-FL (D), and pcDNA3-HA-ORF1 or pcDNA3-ORF1-HA as shown in (A). 24 h post-infection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity in Relative Luminometer Units ($*P < 0.05$). (E) L1 ORF1p does not affect the Tat-mediated HIV-1 transcription. 293T cells (2×10^4 cells/well) were co-transfected with HIV-1 LTR-Luc (100 ng), pcDNA3-HA-ORF1 (100 ng), pcDNA3-ORF1-HA, and/or pcDNA3-Tat101-FLAG (123) (100 ng) in triplicate. 24 h after transfection, luciferase assays were performed, and results were plotted as described above ($*P < 0.05$). (F) L1 ORF1p does not affect the Rev-mediated nuclear export function. 293T cells (2×10^4 cells/well) were co-transfected with pDM628 (47) (100 ng), pcDNA3-HA-ORF1 (100 ng), pcDNA3-ORF1-HA, and/or pcRev (47) (100 ng) in triplicate. 24 h after transfection, luciferase assays were performed, and results were plotted as described above ($*P < 0.05$).

port function using the Rev-dependent luciferase-based reporter plasmid pDM628 (47,49). Our results showed that ORF1p did not affect either Tat or Rev function (Figure 5E and F).

Vpr and p21 do not suppress L1 transcription and ORF1p expression

To determine whether Vpr and p21 suppress L1 mobility by inhibiting the L1 promoter activity, we examined L1 mRNA

expression by real-time RT-PCR. Both p21 and Vpr did not affect the level of L1 mRNA from the L1 5'UTR promoter (pYX014) (Figure 6A). HIV-1 clones (R9 and NL4-3) enhanced L1 transcription two days post-transfection (Figure 6A). Similarly, Vpr did not suppress the level of endogenous L1 mRNA (Figure 6B). Thus, Vpr and p21 may suppress L1 mobility post-transcriptionally. Since Vpr is known to enhance p21 promoter activity (86,87), we wished to confirm this regulation in 293T cells. Indeed, we observed that




Figure 6. Vpr and p21 do not suppress L1 transcription. (A) The level of L1 ORF1 mRNA in 293T cells co-transfected with pYX014 and pcDNA3p21, pCMX-Vpr96, pR9, or pNL4-3 was monitored by real-time LightCycler PCR two days post-transfection, respectively. Experiments were done in triplicate, and graph shows the mean percentages (\pm SEM) of L1 ORF1 mRNA normalized with β -actin mRNA, with the pYX014 condition set at 100% (* $P < 0.05$). (B) The level of endogenous L1 mRNA in 293T cells transfected with pCMX-Vpr96 was monitored by real-time LightCycler PCR. Primer pairs targeting primate (L1PA - ORF1) or human (L1Hs - ORF1 and 5'UTR) L1 subfamilies were used. Experiments were done in triplicate, and graph shows the mean percentages (\pm SEM) of L1 mRNA in presence of Vpr normalized with both β -actin mRNA and the 'no Vpr' condition for each primer pair (* $P < 0.05$). (C) Effect of Vpr on p21 promoter activity. 293T cells (2×10^4 cells/well) were co-transfected with pWAF1-Luc (100 ng) and pCMX-Vpr96 (100 ng) in triplicate. 24h after transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity in Relative Luminometer Units (* $P < 0.05$). (D, E) Vpr suppresses L1 retrotransposition efficiency in a p21-independent manner. p21 (p21KD) knockdown 293T cells (2×10^4 cells/well) were co-transfected with pCMX-Vpr96 (100 ng) and either pYX014 (D) or pYX017 (E) (100 ng) in triplicate. 24h after transfection, luciferase assays were performed. Graph shows the mean (\pm SEM) firefly luciferase activity normalized with *Renilla* luciferase activity with the condition without Vpr set to 100% (* $P < 0.05$).

Vpr weakly enhanced p21 promoter activity (Figure 6C). To examine whether Vpr suppress L1 retrotransposition in a p21-dependent manner, we used p21 knockdown 293T cells. Our results showed that Vpr could strongly suppress L1 retrotransposition even in p21 knockdown cells (Figure 6D and E), suggesting that Vpr suppresses L1 mobility in a p21-independent manner. To test whether Vpr and p21 affect L1 transcription and L1-encoded protein expression, we used an anti-human ORF1 antibody (SE-6798) (65). Vpr and p21 did not alter endogenous ORF1p expression (Figure 7A). Similarly, Vpr and p21 did not suppress ORF1p when expressed from plasmids pJM101/L1.3 and pJM105/L1.3, which respectively contain WT and ORF2p RT-mutant (D702A) human L1.3 element in a pCEP4 backbone vector (88) (Figure 7A and B). Moreover, Vpr did not enhance endogenous p21 expression (Figure 7A). To test the impact of Vpr on pCEP4 expression unit, we used the control backbone plasmid pCEP-GFP that expresses hrGFP (89). Vpr did not affect hrGFP expression from pCEP4 when both proteins were co-expressed in 293T cells (Fig-

ure 7B). Thus, Vpr and p21 did not affect L1 transcription and L1 ORF1p expression.

Vpr and p21 suppress L1 ORF2p-mediated reverse transcription

To examine whether HIV-1 Vpr interacts with L1 ORF1p and/or ORF2p, we performed immunoprecipitation and immunofluorescence studies. We did not observe a co-immunoprecipitation of Vpr with ORF1p (Figure 8A). ORF1p was found in cytoplasmic foci (Figure 8B), consistent with previous reports that ORF1p accumulates in cytoplasmic foci associated with stress granules (14,90). We did not observe co-localization of Vpr with ORF1p. In contrast, we noticed that Vpr and p21 co-immunoprecipitated with ORF2p by using TET-induced FLAG-tagged ORF2p (pLD401) (91) in TET293T cells (Figure 8C). In addition, we confirmed that p21 co-immunoprecipitated with FLAG-tagged ORF2p (pTMO2F3) (89) in 293T cells (Figure 8D). Furthermore, we observed a partial colocalization of ORF2p with either Vpr or p21 (Figure 8E). Vpr alone




Figure 7. Vpr and p21 do not suppress L1 ORF1p expression. (A) The level of L1 ORF1p in presence of Vpr WT and H71R mutant. 293T cells (2×10^5 cells/well) were cotransfected with 2 μ g of pCEP-GFP, pJM105/L1.3 reverse transcriptase-deficient mutant (88), or pJM101/L1.3 wild-type L1 (88), and 2 μ g of pcDNA3-HA, pcDNA3-HA-Vpr WT, pcDNA3-HA-Vpr H71R, or pcDNA3p21. Cells were cultured for 4 days, lysed and subjected to western blot to analyze the expression of ORF1p using anti-hORF1p antibody (SE-6798) (65). Western blotting of the cell lysates with anti-Vpr, anti-p21, and anti- β -actin antibodies is also shown, respectively. (B) Protein expression level of L1 ORF1p and hrGFP in presence of Vpr WT and H71R mutant. 293T cells were transfected with pJM105/L1.3 or pCEP-GFP, together with either pcDNA3-HA or pcDNA3-HA-Vpr WT and cultured for the indicated time. Western blotting of the cell lysates with anti-hORF1p, anti-Vpr, anti-GAPDH and anti-hrGFP (240141; Agilent) antibodies is shown, respectively.

predominantly localized to nucleus and nuclear membrane (Figure 8B), whereas Vpr partially colocalized with ORF2p into cytoplasmic foci (Figure 8E). p21 partially colocalized with ORF2p in the nucleus (Figure 8E). Thus, Vpr and p21 seem to interact with L1 ORF2p but not ORF1p, suggesting a possibility that Vpr and p21 inhibit L1 retrotransposition through ORF2p-mediated activities. To test this possibility, we examined whether Vpr and p21 suppress the L1 reverse transcriptase (RT) activity by LEAP assay. For this approach, we first enriched for L1 RNP from transfected 293T cell extracts, since ORF2p is characterized by a low level of

expression due to its unconventional translation mechanism (14,92). L1 RNA assembles with ORF1p and ORF2p to form RNP complexes in the cytoplasm from which ORF2p RT activity can be detected. Indeed, we detected L1 ORF1p and ribosomal S6 proteins by western blot in the enriched RNP fractions (Figure 9A). S6 protein is abundant in cellular RNP preparations. Interestingly, we noticed that low but detectable amounts of Vpr and p21 were present in the RNP fractions when co-expressed with L1 (Figure 9A), suggesting that Vpr and p21 may be part of L1 RNPs. However, at this stage, the data presented are not sufficient to determine if these proteins actually form part of the L1 RNP. In this context, ORF2p RT activity measured by LEAP assay was significantly reduced with Vpr WT or p21 (Figure 9B), indicating that Vpr and p21 strongly inhibit the L1 RT activity. Consistent with the result of L1 retrotransposition activity with Vpr H71R (Figure 3E and F), Vpr H71R had a marginal effect on the ORF2 RT activity compared with that of Vpr WT (Figure 9B). Altogether, Vpr and p21 seem to restrict L1 retrotransposition through an inhibition of ORF2p RT activity.

DISCUSSION

Several host cellular factors have been involved in the regulation of L1 mobility (12,21–27,91,93). In this study, we have demonstrated that viral (Vpr) and cellular (p21 and p27) cell cycle regulators restrict L1 mobility. Both p21 and p27 belong to the Cip/Kip family of CDK kinase inhibitors (CKIs) (73–77). p21 is the first reported cell cycle inhibitor, blocking cell cycle at G₁/S phase (73–75). p27 arrests the cell cycle at G₁/S phase (76,77). Vpr arrests the host cell cycle at G₂/M phase (53). Therefore, cell cycle control might be important for L1 retrotransposition. In this regard, Shi *et al.* previously reported that L1 retrotransposition was strongly inhibited in cells arrested at G₁, S, G₂ or M phase, indicating that cell division is required for L1 retrotransposition (94). However, the same study found that the levels of L1 transcripts were strongly reduced in arrested cells, suggesting that transcriptional inhibition is largely responsible for reduced retrotransposition observed (94). In contrast, both Vpr and p21 did not suppress L1 transcription (Figure 6A and B), indicating that Vpr and p21 inhibit L1 mobility post-transcriptionally. This result is consistent with studies reporting limited retrotransposition levels in cell cycle arrested cells that could not be explained by reduced L1 transcription (95,96). It suggests that even if retrotransposition can occur in arrested cells, cell division can promote L1 retrotransposition efficiency (95,96). This observation is not in disagreement with other studies reporting events of retrotransposition in non-dividing cells (96,97). On the other hand, Mita *et al.* reported that ORF1p together with ORF2p and L1 mRNA enters the nucleus during mitosis and retrotransposition appears to occur mainly during S phase, indicating a cell cycle bias for L1 retrotransposition (98). Furthermore, recent studies reported that ATM and p53, a master regulator of cell cycle, also regulate L1 mobility (99–101), even though we observed no effect of overexpression of ATM in L1 retrotransposition (Figure 4A). Interestingly, PCNA, the polymerase δ -associated sliding DNA clamp, interacts with L1 ORF2p through a PCNA in-




Figure 8. HIV-1 Vpr and p21 bind to L1 ORF2p. (A) Vpr does not bind to L1 ORF1p. 293T cells (2×10^5 cells/well) were co-transfected with 2 μ g of pcDNA3-HA-Vpr and 2 μ g of pcDNA3-FLAG-ORF1. The cell lysates were immunoprecipitated with anti-FLAG (M2) antibody, followed by immunoblot analysis using anti-FLAG (M2) and anti-HA (HA-7) antibodies. (B) Subcellular localization of HA-ORF1p and EGFP-Vpr. 293T cells were co-transfected with pcDNA3-HA-ORF1 (100 ng) and pEGFP-Vpr (100 ng). Cells were stained with anti-HA (HA-7) antibody and then visualized with Cy3-conjugated anti-mouse antibody. Nucleus was stained with DAPI. Images were obtained using confocal laser scanning microscopy (FV1200, Olympus). (C) Vpr and p21 bind to ORF2p. TET293T cells (2×10^5 cells/well) were transfected with pLD401 (91). Cells were incubated with 1 μ g/ml puromycin for 3 days and then treated with 500 ng/ml doxycycline for 2 days. Then, the cells were transfected with either pcDNA3-HA-Vpr or pcDNA3p21. 2 days post-transfection, the cell lysates were immunoprecipitated with anti-FLAG antibody, followed by immunoblot analysis using anti-FLAG, anti-HA, or p21 antibody. (D) p21 binds to ORF2p. 293T cells (2×10^5 cells/well) were transfected with pTMO2F3 (89). Three days post-transfection, the cell lysates were immunoprecipitated with anti-FLAG antibody, followed by immunoblot analysis using anti-FLAG, and p21 antibodies. (E) Subcellular localization of Vpr, p21, and/or ORF2p-FLAG. 293T cells were co-transfected with pTM02F3 (100 ng) and either pcDNA3-HA-Vpr (100 ng) or pcDNA3 p21 (100ng). Cells were stained with anti-FLAG (M2) and either anti-HA (3F10) or anti-p21 antibodies and then visualized with Alexa596-conjugated anti-mouse and either Alexa488-conjugated anti-rat or anti-rabbit antibodies. Nucleus was stained with DAPI. Images were obtained using confocal laser scanning microscopy.

teracting protein (PIP) box (91). The interaction of ORF2p with PCNA is critical for L1 retrotransposition (91). In this regard, PCNA is known to bind to the C-terminal domain of p21 (56), suggesting a possibility that p21 competes with L1 ORF2p for the PCNA-binding, resulting in the repression of L1 retrotransposition. However, the C-terminal half of p21, p21C, did not reduced L1 retrotransposition (Figure 3C and D), indicating that p21 restricts L1 retrotransposition in a PCNA independent manner. In this study, we

showed that p21 interacts with L1 ORF2p and inhibits the ORF2p reverse transcriptase activity in LEAP assay (Figures 8 and 9).

We showed that the inhibitory effect of a mutant Vpr (H71R), which is unable to arrest the cell cycle, was significantly reduced compared with that of wild-type Vpr (Figure 3E and F), suggesting that Vpr suppresses L1 mobility in a cell cycle dependent manner. Independent of cell cycle control, we have demonstrated that Vpr restricts L1




Figure 9. Vpr and p21 suppresses L1 reverse transcriptase activity. (A) Western blotting of human ORF1p, HA-Vpr, p21 or S6 protein as a loading control in RNP prepared from 293T cells transfected with the indicated plasmids: pCEP4 and pcDNA3-HA (HA) empty vectors (-), pJM105/L1.3 and pcDNA3-HA (-) or pJM101/L1.3 together with pcDNA3-HA (-), pcDNA3-HA-Vpr WT, pcDNA3-HA-Vpr H71R, or pcDNA3 p21. (-) Detection of L1 RT activity by LEAP assay (top panel) and of L1 RNA (middle panel) or GAPDH RNA (bottom panel) by RT-PCR. LEAP assay was performed with RNP prepared from cells transfected with indicated plasmids. PCR control denotes a control for the PCR step without cDNA. LEAP/RT control denotes a control for RT step without RNP. RNP samples used in panel A and B are from the same replicate. The experiment was replicated three times with similar results.

retrotransposition by inhibiting ORF2p-mediated reverse transcriptase activity (Figure 9B). Moreover, despite limited levels of ORF2p detection, possibly due to unconventional translational mechanism (14,92) or protein stability, we found that Vpr coimmunoprecipitates with L1 ORF2p but not ORF1p (Figure 8A, C and D). Vpr can also be detected in cellular RNP fractions alongside with L1 (Figure 9A), providing additional circumstantial evidence that Vpr associates with ORF2p and inhibits RT activity (Figure 9B). On the other hand, p21 has been involved in restriction of HIV-1 during reverse transcription (102–105). Ribonucleotide reductase (RNR) increases the dNTP pool and HIV-1 relies on cellular dNTPs for its reverse transcription. p21 represses dNTP biosynthesis by down-regulating the expression of the RNR2 subunit of ribonucleotide reductase, resulting in the restriction of HIV-1 (104). Furthermore, Pauls *et al.* recently reported that p21 regulates a sterile alpha motif (SAM) and histidine/aspartic acid (HD) domain-containing protein 1 (SAMHD1) (105). The deoxynucleoside triphosphohydrolase SAMHD1 has been identified as a novel myeloid cell specific HIV-1 restriction factor (106–111). Indeed, SAMHD1 inhibits HIV-1 reverse transcription by depleting the intracellular dNTP pool (111) and restricts L1 retrotransposition by inhibiting the ORF2p-mediated L1 reverse transcription (112). Similarly, we have demonstrated that p21 interacts with ORF2p (Figure 8C and D) and can also be detected in cellular RNP fractions alongside with L1 (Figure 9A), resulting in inhibition of ORF2p RT activity (Figure 9B). Therefore, in HIV-1-infected patient cells or tumor cells with overexpression of p21, or a mutation in the N-terminus of p21 or p27, these cell cycle regulators may potentially impact L1 retrotransposition. However, if p21 is overexpressed in HIV-1-infected cells, p21 may inhibit both HIV-1 and L1 mobility. As Vpr is known to enhance the p21 promoter activity (86,87), it raised the possibility that Vpr transcriptionally induces the p21, resulting in the suppression of L1 mobility. However,

we showed that Vpr did not accumulate endogenous p21 expression in 293T cells (Figure 7A) and Vpr could suppress L1 retrotransposition even in p21 knockdown cells (Figure 6D and E), suggesting that Vpr suppresses L1 mobility in a p21-independent manner.

Since endogenous retroelements such as L1, Alu, SVA, and HERVs are abundant and constitute ~45% of the human genome, they may influence the integrated exogenous HIV-1 retroviruses, or vice versa. In this study, we demonstrated that L1 ORF1p is incorporated into the HIV-1 virions (Figure 5), suggesting a possibility that infectious HIV-1 particle might horizontally transmit L1 into target cells as a vector. Another possibility is that the incorporated ORF1p affects the HIV-1 infectivity. However, the incorporation of ORF1-HA into HIV-1 virion did not affect the HIV-1 infectivity, except only HA-ORF1 moderately suppressed the infectivity of NL4-3 but not JR-FL (Figure 5C and D), even though similar amounts of ORF1p were incorporated into both NL4-3 and JR-FL virion (Figure 5A). Since ORF1p is an RNA-binding and nucleic acids chaperone protein, we also examined whether ORF1p modulates the HIV-1 Tat-mediated HIV-1 transcription or HIV-1 Rev-dependent nuclear export function of HIV-1 RNA. However, ORF1p did not affect either Tat or Rev function (Figure 5E and F), suggesting that ORF1p might not bind and affect HIV-1 RNA. In this regard, we failed to observe the interaction of ORF1p with HIV-1 RNA (data not shown). Furthermore, we found that the incorporation of ORF1p into HIV-1 virions was independent of HIV-1 accessory proteins including Nef, Vpr, Vpu and Vif, because the mutant viruses lacking the expression of each viral gene still retained the ability to incorporate ORF1p (Figure 5B). Since it has been indicated that several P-body and stress granule components, such as APOBEC3G, APOBEC3F and MOV10, are incorporated into HIV-1 virions through an interaction with nucleocapsid protein (NC) (78–82), ORF1p, a component associated to stress granule

(14,90), also might be incorporated into virion through an interaction with NC. To fully explain these phenomena, further mechanistic studies are required. Jones *et al.* demonstrated that the L1 DNA accumulates in the HIV-1 infected cells (113). In this regard, we also observed that HIV-1 infection slightly enhances the L1 transcription (Figure 6A). In a previous study, Iijima *et al.* reported that extracellular Vpr protein in the culture medium induces L1 retrotransposition (114). In contrast, we found that Vpr strongly suppresses L1 retrotransposition when co-expressed in the cell (Figure 2). The cause for this discrepancy may be due to technical differences and experimental strategies. Iijima *et al.* only used recombinant Vpr protein in their experiments. In contrast, we used Vpr-expressing plasmids to obtain intracellular expression of Vpr in dividing cells. All tested plasmids (pcDNA3-HA-Vpr, pCMX Vpr96, pAIP-Vpr and pEGFP-Vpr) could strongly suppress L1 retrotransposition and LEAP assay showed that Vpr inhibits L1 RT activity (Figure 9B). Except for the demonstration that Vpr inhibits reverse transcription by ORF2p, the other proteins may be inhibiting L1 indirectly or act on cell cycle to limit the expression of the reporter cassette.

Finally, if L1 actively jumps and inserts a new copy of L1 in the integrated HIV-1 proviral locus or in genes essential for the infected cell survival, it will result in the destruction of HIV-1 or the infected cell. Therefore, we can speculate that HIV-1 may restrict L1 mobility to protect its own genome and the genome of the infected cell from *de novo* L1 insertional mutagenesis. Altogether, we suggested that viral and cellular cell cycle regulators can negatively impact L1 mobility through an inhibition of ORF2p reverse transcriptase.

ACKNOWLEDGEMENTS

We thank Drs Gaël Cristofari, Haig H. Kazazian Jr, Jef D. Boeke, John V. Moran, Tomoichiro Miyoshi, Michael B. Kastan, Stephen J. Elledge, Anindya Dutta, Reuven Agami, Richard Iggo, Yoichi Taya, Akio Adachi, Hiroyuki Sakai, Roger Pomerantz, Monsef Benkirane, and Jeremy Luban for reagents. We also thank Drs Castro-Diaz Nathaly, Casano Marco, and members of the Gene Technology Center in Kumamoto University for their important contributions to the experiments. We thank Ms Kazumi Tsuruhara for secretary assistance of a JSPS Lateral Research.

FUNDING

JSPS Lateral Research [14544610]; Grant-in-Aid for Research on Hepatitis from the Ministry of Health, Labor, and Welfare of Japan [17929672; 16768555]; Takeda Foundation; AIDS International Collaborative Research Grant from the Ministry of Education, Science, Sports, and Culture; Kumamoto University. Funding for open access charge: AIDS International Collaborative Research Grant from the Ministry of Education, Science, Sports, and Culture; Kumamoto University.

Conflict of interest statement. None declared.

REFERENCES

- Ostertag, E.M. and Kazazian, H.H. Jr. (2001) Biology of mammalian L1 retrotransposons. *Annu. Rev. Genet.*, **35**, 501–538.
- Hancks, D.C. and Kazazian, H.H. Jr. (2012) Active human retrotransposons: variation and disease. *Curr. Opin. Genet. Dev.*, **22**, 191–203.
- Mills, R.E., Bennett, E.A., Iskow, R.C. and Devine, S.E. (2007) Which transposable elements are active in the human genome? *Trends Genet.*, **23**, 183–191.
- Martin, S.L. and Bushman, F.D. (2001) Nucleic acid chaperon activity of the ORF1 protein from the mouse LINE-1 retrotransposon. *Mol. Cell Biol.*, **21**, 467–475.
- Mathias, S.L., Scott, A.F., Kazazian, H.H. Jr, Boeke, J.D. and Gabriel, A. (1991) Reverse transcriptase encoded by a human transposable element. *Science*, **254**, 1808–1810.
- Swergold, G.D. (1990) Identification, characterization, and cell specificity of a human LINE-1 promoter. *Mol. Cell Biol.*, **10**, 6718–6729.
- Harris, C.R., Dewan, A., Zupnick, A., Normart, R., Gabriel, A., Prives, C., Levine, A.J. and Hoh, J. (2009) p53 responsive elements in human retrotransposons. *Oncogene*, **28**, 3857–3865.
- Yang, N., Zhang, L., Zhang, Y. and Kazazian, H.H. Jr. (2003) An important role for RUNX3 in human L1 transcription and retrotransposition. *Nucleic Acids Res.*, **31**, 4929–4940.
- Tchenio, T., Casella, J.F. and Heidmann, T. (2000) Members of the SRY family regulate the human LINE retrotransposons. *Nucleic Acids Res.*, **28**, 411–415.
- Becker, K.G., Swergold, G.D., Ozato, K. and Thayer, R.E. (1993) Binding of the ubiquitous nuclear transcription factor YY1 to a cis regulatory sequence in the human LINE-1 transposable element. *Hum. Mol. Genet.*, **2**, 1697–1702.
- Athanikar, J.N., Badge, R.M. and Moran, J.V. (2004) A YY1-binding site is required for accurate human LINE-1 transcription initiation. *Nucleic Acids Res.*, **32**, 3846–3855.
- Ariumi, Y. (2016) Guardian of the human genome: host defense mechanisms against LINE-1 retrotransposition. *Front. Chem.*, **4**, 28.
- Muotri, A.R., Chu, V.T., Marchetto, M.C., Deng, W., Moran, J.V. and Gage, F.H. (2005) Somatic mosaicism in neuronal precursor cells mediated by L1 retrotransposition. *Nature*, **435**, 903–910.
- Doucet, A.J., Hulme, A.E., Sahinovic, E., Kulpa, D.A., Moldovan, J.B., Kopera, H.C., Athanikar, J.N., Hasnaoui, M., Bucheton, A., Moran, J.V. *et al.* (2010) Characterization of LINE-1 ribonucleoprotein particles. *PLoS Genet.*, **6**, e1001150.
- Feng, Q., Moran, J.V., Kazazian, H.H. Jr and Boeke, J.D. (1996) Human L1 retrotransposon encodes a conserved endonuclease required for retrotransposition. *Cell*, **87**, 905–916.
- van den Hurk, J.A., Meij, I.C., Seleme, M.C., Kano, H., Nikopoulos, K., Hoefsloot, L.H., Sistermans, E.A., de Wijs, I.J., Mukhopadhyay, A., Plomp, A.S. *et al.* (2007) L1 retrotransposition can occur early in human embryonic development. *Hum. Mol. Genet.*, **16**, 1587–1592.
- Kano, H., Godoy, I., Courtney, C., Vetter, M.R., Gerton, G.L., Ostertag, E.M. and Kazazian, H.H. Jr. (2009) L1 retrotransposition occurs mainly in embryogenesis and creates somatic mosaicism. *Genes Dev.*, **23**, 1303–1312.
- Richardson, S.R., Gerdes, P., Gerhardt, D.J., Sanchez-Luque, F.J., Bodea, G.O., Muñoz-Lopez, M., Jesuadian, J.S., Kempen, M.H.C., Carreira, P.E., Jeddloh, J.A. *et al.* (2017) Heritable L1 retrotransposition in the mouse primordial germline and early embryo. *Genome Res.*, **27**, 1395–1405.
- Newkirk, S.J., Lee, S., Grandi, F.C., Gaysinskaya, V., Rosser, J.M., Vanden Berg, N., Hogarth, C.A., Marchetto, M.C.N., Muotri, A.R., Griswold, M.D. *et al.* (2017) Intact piRNA pathway prevents L1 mobilization in male meiosis. *Proc. Natl. Acad. Sci. U.S.A.*, **114**, E5635–E5644.
- Catro-Diaz, N., Ecco, G., Coluccio, A., Kapopoulou, A., Yazdanpanah, B., Friedli, M., Duc, J., Jang, S.M., Turelli, P. and Trono, D. (2014) Evolutionarily dynamic L1 regulation in embryonic stem cells. *Genes Dev.*, **28**, 1397–1409.
- Bogerd, H.P., Wiegrand, H.L., Hulme, A.E., Garcua-Perez, J.L., O’Shea, K.S., Moran, J.V. and Cullen, B.R. (2006) Cellular inhibitors of long interspersed element 1 and Alu retrotransposition. *Proc. Natl. Acad. Sci. U.S.A.*, **103**, 8780–8785.

22. Muckenfuss, H., Hamdorf, M., Held, U., Perkovic, M., Löwer, J., Cichutek, K., Flory, E., Schumann, G.G. and Münk, C. (2006) APOBEC3 proteins inhibit human LINE-1 retrotransposition. *J. Biol. Chem.*, **281**, 22161–22172.
23. Stenglein, M.D. and Harris, R.S. (2006) APOBEC3B and APOBEC3F inhibit L1 retrotransposition by a DNA deamination-independent mechanism. *J. Biol. Chem.*, **281**, 16837–16841.
24. Kinomoto, M., Kanno, T., Shimura, M., Ishizaka, Y., Kojima, A., Kurata, T., Sata, T. and Tokunaga, K. (2007) All APOBEC3 family proteins differentially inhibit LINE-1 retrotransposition. *Nucleic Acids Res.*, **35**, 2955–2964.
25. Arjan-Odedra, S., Swanson, C.M., Sherer, N.M., Wolinsky, S.M. and Malim, M.H. (2012) Endogenous MOV10 inhibits the retrotransposition of endogenous retroelements but not the replication of exogenous retroviruses. *Retrovirology*, **9**, 53.
26. Goodier, J.L., Cheung, L.E. and Kazazian, H.H. Jr. (2012) MOV10 RNA helicase is a potent inhibitor of retrotransposition in cells. *PLoS Genet.*, **8**, e1002941.
27. Li, X., Zhang, J., Jia, R., Cheng, V., Xu, X., Qiao, W., Guo, F., Liang, C. and Cen, S. (2013) The MOV10 helicase inhibits LINE-1 mobility. *J. Biol. Chem.*, **288**, 21146–21160.
28. Sheehy, A.M., Gaddis, N.C., Choi, J.D. and Malim, M.H. (2002) Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral vif protein. *Nature*, **418**, 646–650.
29. Mangeat, B., Turelli, P., Caron, G., Friedli, M., Perrin, L. and Trono, D. (2003) Broad antiretroviral defence by human APOBEC3G through lethal editing of nascent reverse transcripts. *Nature*, **424**, 99–103.
30. Zhang, H., Yang, B., Pomerantz, R.J., Zhang, C., Arunachalam, S.C. and Gao, L. (2003) The cytidine deaminase CEM15 induces hypermutation in newly synthesized HIV-1 DNA. *Nature*, **424**, 94–98.
31. Turelli, P., Mangeat, B., Jost, S., Vianin, S. and Trono, D. (2004) Inhibition of hepatitis B virus replication by APOBEC3G. *Science*, **303**, 1829.
32. Esnault, C., Heidmann, O., Delebecque, F., Dewannieux, M., Ribet, D., Hance, A.J., Heidmann, T. and Schwartz, O. (2005) APOBEC3G cytidine deaminase inhibits retrotransposition of endogenous retrovirus. *Nature*, **433**, 430–433.
33. Hulme, A.E., Bogerd, H.P., Cullen, B.R. and Moran, J.V. (2007) Selective inhibition of Alu retrotransposition by APOBEC3G. *Gene*, **390**, 199–205.
34. Bulliard, Y., Turelli, P., Röhrig, U.F., Zoete, V., Mangeat, B., Michielin, O. and Trono, D. (2009) Functional analysis and structural modeling of human APOBEC3G reveal the role of evolutionarily conserved elements in the inhibition of human immunodeficiency virus type 1 infection and *Alu* transposition. *J. Virol.*, **83**, 12611–12621.
35. Kazazian, H.H. Jr, Wong, C., Youssoufian, H., Scott, A.F., Phillips, D.G. and Antonarakis, S.E. (1988) Haemophilia A resulting from *de novo* insertion of L1 sequence represents a novel mechanism for mutation in man. *Nature*, **332**, 164–166.
36. Miki, Y., Nishisho, I., Horii, A., Miyoshi, Y., Utsunomiya, J., Kinzler, K.W., Vogelstein, B. and Nakamura, Y. (1992) Disruption of the *APC* gene by a retrotransposal insertion of L1 sequence in a colon cancer. *Cancer Res.*, **52**, 643–645.
37. Gilbert, N., Lutz-Prigge, S. and Moran, J. (2002) Genomic deletion created upon LINE-1 retrotransposition. *Cell*, **110**, 315–325.
38. Symer, D.E., Connelly, C., Szak, S.T., Caputo, E.M., Cost, G.J., Parmigiani, G. and Boeke, J.D. (2002) Human L1 retrotransposition is associated with genetic instability in vivo. *Cell*, **110**, 327–338.
39. Gasior, S.L., Wakeman, T.P., Xu, B. and Deininger, P.L. (2006) The human LINE-1 retrotransposon creates DNA double-strand breaks. *J. Mol. Biol.*, **357**, 1383–1393.
40. Hancks, D.C. and Kazazian, H.H. Jr. (2016) Role for retrotransposon insertions in human disease. *Mob. DNA*, **7**, 9.
41. Jeang, K.T., Xiao, H. and Rich, E.A. (1999) Multifaceted activities of the HIV-1 transactivator of transcription, Tat. *J. Biol. Chem.*, **274**, 28837–28840.
42. Berkhout, B., Silverman, R.H. and Jeang, K.T. (1989) Tat *trans*-activates the human immunodeficiency virus through a nascent RNA target. *Cell*, **59**, 273–282.
43. Hope, T.J. and Pomerantz, R.J. (1995) The human immunodeficiency virus type 1 Rev protein: a pivotal protein in the viral life cycle. *Curr. Top. Microbiol. Immunol.*, **193**, 91–105.
44. Pollard, V.W. and Malim, M.H. (1998) The HIV-1 Rev protein. *Annu. Rev. Microbiol.*, **52**, 491–532.
45. Cullen, B.R. (2003) Nuclear mRNA export: insights from virology. *Trends Biochem. Sci.*, **28**, 419–424.
46. Yedavalli, V.S., Neuveut, C., Chi, Y.H., Kleiman, L. and Jeang, K.T. (2004) Requirement of DDX3 DEAD box RNA helicase for HIV-1 Rev-RRE export function. *Cell*, **119**, 381–392.
47. Fang, J., Kubota, S., Yang, B., Zhou, N., Zhang, H., Godbout, R. and Pomerantz, R.J. (2004) A DEAD box protein facilitates HIV-1 replication as a cellular co-factor of Rev. *Virology*, **330**, 471–480.
48. Naji, S., Ambrus, G., Cimermančić, P., Reyes, J.R., Johnson, J.R., Fibrandt, R., Huber, M.D., Vesely, P., Krogan, N.J., Yates, J.R. 3rd *et al.* (2012) Host cell interactome of HIV-1 Rev includes RNA helicases involved in multiple facets of virus production. *Mol. Cell. Proteomics*, **11**:M111.
49. Yasuda-Inoue, M., Kuroki, M. and Ariumi, Y. (2013) Distinct DDX DEAD-box RNA helicases cooperate to modulate the HIV-1 Rev function. *Biochem. Biophys. Res. Commun.*, **434**, 803–808.
50. Ariumi, Y. (2014) Multiple functions of DDX3 RNA helicase in gene regulation, tumorigenesis, and viral infection. *Frontiers Genet.*, **5**, 423.
51. Cohen, E.A., Dehni, G., Sodroski, J.G. and Haseltine, W.A. (1990) Human immunodeficiency virus vpr product is a virion-associated regulatory protein. *J. Virol.*, **64**, 3097–3099.
52. Kogan, M. and Rappaport, J. (2011) HIV-1 accessory protein Vpr: relevance in the pathogenesis of HIV and potential for therapeutic intervention. *Retrovirology*, **8**, 25.
53. Goh, W.C., Rogel, M.E., Kinsey, C.M., Michael, S.F., Fultz, P.N., Nowak, M.A., Hahn, B.H. and Emerman, M. (1998) HIV-1 Vpr increases viral expression by manipulation of the cell cycle: a mechanism for selection of Vpr in vivo. *Nat. Med.*, **4**, 65–71.
54. Ostertag, E.M., Prak, E.T., DeBerardinis, R.J., Moran, J.V. and Kazazian, H.H. Jr. (2000) Determination of L1 retrotransposition kinetics in cultured cells. *Nucleic Acids Res.*, **28**, 1418–1423.
55. Pertel, T., Hausmann, S., Morger, D., Züger, S., Guerra, J., Lascano, J., Reinhard, C., Santoni, F.A., Uchil, P.D., Chatel, L. *et al.* (2011) TRIM5 is an innate immune sensor for the retrovirus capsid lattice. *Nature*, **472**, 316–365.
56. Chen, J., Jackson, P.K., Kirschner, M.W. and Dutta, A. (1995) Separate domains of p21 involved in the inhibition of Cdk kinase and PCNA. *Nature*, **374**, 386–388.
57. Brummelkamp, T.R., Bernard, R. and Agami, R. (2002) A system for stable expression of short interfering RNAs in mammalian cells. *Science*, **296**, 550–553.
58. Bridge, A.J., Pebernard, S., Ducaux, A., Nicoulaz, A.L. and Iggo, R. (2003) Induction of an interferon response by RNAi vectors in mammalian cells. *Nat. Genet.*, **34**, 263–264.
59. Naldini, L., Blömer, U., Gally, P., Ory, D., Mulligan, R., Gage, F.H., Verma, I.M. and Trono, D. (1996) In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science*, **272**, 263–267.
60. Zufferey, R., Nagy, D., Mandel, R.J., Naldini, L. and Trono, D. (1997) Multiply attenuated lentiviral vector achieves efficient gene delivery in vivo. *Nat. Biotechnol.*, **15**, 871–875.
61. Adachi, A., Gendelman, H.E., Koenig, S., Folks, T., Willey, R., Rabson, A. and Martin, M.A. (1996) Production of acquired immunodeficiency syndrome-associated retrovirus in human and non human cells transfected with an infectious molecular clones. *J. Virol.*, **59**, 284–291.
62. Koyanagi, Y., Miles, S., Mitsuyasu, R.T., Merrill, J.E., Vinters, H.V. and Chen, I.S. (1987) Dual infection of the central nervous system by AIDS viruses with distinct cellular tropisms. *Science*, **236**, 819–822.
63. Schubert, U., Bour, S., Willey, R.L. and Strelbel, K. (1999) Regulation of virus release by the macrophage-tropic human immunodeficiency virus type 1 AD8 isolate is redundant and can be controlled by either Vpu or Env. *J. Virol.*, **73**, 887–896.
64. Ariumi, Y. and Trono, D. (2006) Ataxia-telangiectasia-mutated (ATM) protein can enhance human immunodeficiency virus type 1 replication by stimulating Rev function. *J. Virol.*, **80**, 2445–2452.
65. Monot, C., Kuciak, M., Viollet, S., Mir, A.A., Gabus, C., Darlix, J.L. and Cristofari, G. (2013) The specificity and flexibility of L1 reverse

- transcription priming at imperfect T-tracts. *PLoS Genet.*, **9**, e1003499.
66. Krishan, A. (1975) Rapid flow cytofluorometric analysis of mammalian cell cycle by propidium iodide staining. *J. Cell Biol.*, **66**, 188–193.
 67. Viollet, S., Doucet, A.J. and Cristofari, G. (2016) Biochemical approaches to study LINE-1 reverse transcriptase activity in vitro. *Methods Mol. Biol.*, **1400**, 357–376.
 68. Sambrook, J. and Russell, D.W. (2006) Calcium phosphate-mediated transfection of eukaryotic cells with plasmid DNAs. *CSH Protoc.*, **2006**, pdb.prot3871.
 69. Kulpa, D.A. and Moran, J.V. (2006) Cis-preferential LINE-1 reverse transcriptase activity in ribonucleoprotein particles. *Nat. Struct. Mol. Biol.*, **13**, 655–660.
 70. Freeman, J.D., Goodchild, N.L. and Manger, D.L. (1994) A modified indicator gene for selection of retrotransposition events in mammalian cells. *BioTechniques*, **46**, 48–49.
 71. Moran, J.V., Holmes, S.E., Naas, T.P., DeBerardinis, R.J. and Kazazian, H.H. Jr. (1996) High frequency retrotransposition in cultured mammalian cells. *Cell*, **87**, 917–927.
 72. Xie, Y., Rosser, J.M., Thompson, T.L., Boeke, J.D. and An, W. (2011) Characterization of L1 retrotransposition with high-throughput dual-luciferase assays. *Nucleic Acids Res.*, **39**, e16.
 73. Xiong, Y., Hannon, G.J., Zhang, H., Casso, D., Kobayashi, R. and Beach, D. (1993) p21 is a universal inhibitor of cyclin kinases. *Nature*, **366**, 701–704.
 74. Gu, Y., Turck, C.W. and Morgan, D.O. (1993) Inhibition of CDK2 activity in vivo by an associated 20K regulatory subunit. *Nature*, **366**, 707–710.
 75. Harper, J.W., Adami, G.R., Wei, N., Keyomarsi, K. and Elledge, S.J. (1993) The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell*, **75**, 805–816.
 76. Polyak, K., Lee, M.H., Erdjument-Bromage, H., Koff, A., Roberts, J.M., Tempst, P. and Massagué, J. (1994) Cloning of p27Kip1, a cyclin-dependent kinase inhibitor and a potential mediator of extracellular antimitogenic signals. *Cell*, **78**, 59–66.
 77. Toyoshima, H. and Hunter, T. (1994) p27, a novel inhibitor of G1 cyclin-Cdk protein kinase activity, is related to p21. *Cell*, **78**, 67–74.
 78. Wang, X., Han, Y., Dang, Y., Fu, W., Zhou, T., Ptak, R.G. and Zheng, Y.H. (2010) Moloney leukemia virus 10 (MOV10) protein inhibits retrovirus replication. *J. Biol. Chem.*, **285**, 14346–14355.
 79. Burdick, R., Smith, J.L., Chaiphan, C., Friew, Y., Chen, J., Venkatachari, N.J., Delvikis-Frankenberry, K.A., Hu, W.S. and Pathak, V.K. (2010) P body-associated protein Mov10 inhibits HIV-1 replication at multiple stages. *J. Virol.*, **84**, 10241–10253.
 80. Abudu, A., Wang, X., Dang, Y., Zhou, T., Xiang, S.H. and Zheng, Y.H. (2012) Identification of molecular determinants from Moloney leukemia virus 10 homolog (MOV10) protein for virion packaging and anti-HIV-1 activity. *J. Biol. Chem.*, **287**, 1220–1228.
 81. Zheng, Y.H., Irwin, D., Kuroso, T., Tokunaga, K., Sata, T. and Peterlin, B.M. (2004) Human APOBEC3F is another host factor that blocks human immunodeficiency virus type 1 replication. *J. Virol.*, **78**, 6073–6076.
 82. Wiegand, H.L., Doehle, B.P., Bogerd, H.P. and Cullen, B.R. (2004) A second human antiretroviral factor, APOBEC3F, is suppressed by the HIV-1 and HIV-2 Vif proteins. *EMBO J.*, **23**, 2451–2458.
 83. Ariumi, Y., Kaida, A., Hatanaka, M. and Shimotohno, K. (2001) Functional cross-talk of HIV-1 Tat with p53 through its C-terminal domain. *Biochem. Biophys. Res. Commun.*, **287**, 556–561.
 84. Ariumi, Y., Serhan, F., Turelli, P., Telenti, A. and Trono, D. (2006) The integrase interactor 1 (INI1) proteins facilitate Tat-mediated human immunodeficiency virus type 1 transcription. *Retrovirology*, **3**, 47.
 85. Yasuda-Inoue, M., Kuroki, M. and Ariumi, Y. (2013) DDX3 RNA helicase is required for HIV-1 Tat function. *Biochem. Biophys. Res. Commun.*, **441**, 607–611.
 86. Chowdhury, I.H., Wang, X.F., Landau, N.R., Robb, M.L., Polonis, V.R., Birk, D.L. and Kim, J.H. (2003) HIV-1 Vpr activates cell cycle inhibitor p21/Waf1/Cip1: a potential mechanism of G2/M cell cycle arrest. *Virology*, **305**, 371–377.
 87. Amini, S., Saunders, M., Kelley, K., Khalili, K. and Sawaya, B.E. (2004) Interplay between HIV-1 Vpr and Sp1 modulates p21(WAF1) gene expression in human astrocytes. *J. Biol. Chem.*, **279**, 46046–46056.
 88. Moran, J.V., DeBerardinis, R.J. and Kazazian, H.H. Jr. (1999) Exon shuffling by L1 retrotransposition. *Science*, **283**, 1530–1534.
 89. Doucet, A.J., Wilusz, J.E., Miyoshi, T., Liu, Y. and Moran, J.V. (2015) A 3' POLY(A) tract is required for LINE-1 retrotransposition. *Mol. Cell*, **60**, 728–741.
 90. Goodier, J.L., Zhnag, L., Vetter, M.R. and Kazazian, H.H. Jr. (2007) LINE-1 ORF1 protein localizes in stress granules with other RNA-binding proteins, including components of RNA interference RNA-induced silencing complex. *Mol. Cell Biol.*, **27**, 6469–6483.
 91. Taylor, M.S., Lacava, J., Mita, P., Molloy, K.R., Huang, C.R., Li, D., Adney, E.M., Jiang, H., Burns, K.H., Chait, B.T. et al. (2013) Affinity proteomics reveals human host factors implicated in discrete stages of LINE-1 retrotransposition. *Cell*, **155**, 1034–1048.
 92. Alisch, R.S., Garcia-Perez, J.L., Muotri, A.R., Gage, F.H. and Moran, J.V. (2006) Unconventional translation of mammalian LINE-1 retrotransposons. *Genes Dev.*, **20**, 210–224.
 93. Goodier, J.L., Cheung, L.E. and Kazazian, H.H. Jr. (2013) Mapping the LINE1 ORF1 protein interactome reveals associated inhibitors of human retrotransposition. *Nucleic Acids Res.*, **41**, 7401–7419.
 94. Shi, X., Seluanov, A. and Gorbunova, V. (2007) Cell divisions are required for L1 retrotransposition. *Mol. Cell Biol.*, **27**, 1264–1270.
 95. Xie, Y., Mates, L., Ivics, Z., Izsvák, Z., Martin, S.L. and An, W. (2013) Cell division promotes efficient retrotransposition in a stable L1 reporter cell line. *Mob. DNA*, **4**, 10.
 96. Kubo, S., Seleme, M.C., Soifer, H.S., Perez, J.L., Moran, J.V., Kazazian, H.H. Jr. and Kasahara, N. (2006) L1 retrotransposition in nondividing and primary human somatic cells. *Proc. Natl. Acad. Sci. U.S.A.*, **103**, 8036–8041.
 97. Macia, A., Widmann, T.J., Heras, S.R., Ayllon, V., Sanchez, L., Benkaddour-Boumzaouad, M., Muñoz-Lopez, M., Rubio, A., Amador-Cubero, S., Bianco-Jimenez, E. et al. (2017) Engineered LINE-1 retrotransposition in nondividing human neurons. *Genome Res.*, **27**, 335–348.
 98. Mita, P., Wudzinska, A., Sun, X., Andrade, J., Nayak, S., Kahler, D.J., Badri, S., LaCava, J., Ueberheide, B., Yun, C.Y. et al. (2018) LINE-1 protein localization and functional dynamics during the cell cycle. *eLife*, **7**, e30058.
 99. Coufal, N.G., Garcia-Perez, J.L., Peng, G.E., Marchetto, M.C., Muotri, A.R., Mu, Y., Carson, C.T., Macia, A., Moran, J.V. and Gage, F.H. (2011) Ataxia telangiectasia mutated (ATM) modulates long interspersed element-1 (L1) retrotransposition in human neural stem cells. *Proc. Natl. Acad. Sci. U.S.A.*, **108**, 20382–20387.
 100. Wallace, N.A., Gasior, S.L., Faber, Z.J., Howie, H.L., Deininger, P.L. and Galloway, D.A. (2013) HPV 5 and 8 E6 expression reduces ATM protein levels and attenuates LINE-1 retrotransposition. *Virology*, **443**, 69–79.
 101. Harris, C.R., Dewan, A., Zupnick, A., Normart, R., Gabriel, A., Prives, C., Levine, A.J. and Hoh, J. (2009) p53 responsive elements in human retrotransposons. *Oncogene*, **28**, 3857–3865.
 102. Zhang, J., Scadden, D.T. and Crumpacker, C.S. (2007) Primitive hematopoietic cells resist HIV-1 infection via p21^{Waf1/Cip1/Sdi1}. *J. Clin. Invest.*, **117**, 473–481.
 103. Bergamaschi, A., David, A., Le Rouzic, E., Nisole, S., Barré-Sinoussi, F. and Pancino, G. (2009) The CDK inhibitor p21Cip1/WAF1 is induced by FcγR activation and restricts the replication of human immunodeficiency virus type 1 and related primate lentiviruses in human macrophages. *J. Virol.*, **83**, 12253–12265.
 104. Allouch, A., David, A., Amie, S.M., Lahouassa, H., Chartier, L., Margottin-Gouget, F., Barré-Sinoussi, F., Kim, B., Sáez-Cirión, A. and Pancino, G. (2013) p21-mediated RNR2 repression restricts HIV-1 replication in macrophages by inhibiting dNTP biosynthesis. *Proc. Natl. Acad. Sci. U.S.A.*, **110**, E3997–E4006.
 105. Pauls, E., Ruiz, A., Riveira-Muñoz, E., Permanyer, M., Badia, R., Clotet, B., Keppler, O.T., Ballana, E. and Este, J.A. (2014) p21 regulates the HIV-1 restriction factor SAMHD1. *Proc. Natl. Acad. Sci. U.S.A.*, **111**, E1322–E1324.
 106. Hrecka, K., Hao, C., Gierszewska, M., Swanson, S.K., Kesik-Brodacka, M., Strivastava, S., Florens, L., Washburn, A.P. and Skowronski, J. (2011) Vpx relieves inhibition of HIV-1 infection of macrophages mediated by the SAMHD1 protein. *Nature*, **474**, 658–661.
 107. Laguet, N., Sobhian, B., Casartelli, N., Ringeard, M., Chable-Bessia, C., Ségéral, E., Yatim, A., Emiliani, S., Schwartz, O.

- and Benkirane, M. (2011) SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. *Nature*, **474**, 654–657.
108. Goldstone, D.C., Ennis-Adeniran, V., Hedden, J.J., Groom, H.C., Rice, G.I., Christodoulou, E., Walker, P.A., Kelly, G., Haire, L.F., Yap, M.W. *et al.* (2011) HIV-1 restriction factor SAMHD1 is a deoxynucleoside triphosphate triphosphohydrolase. *Nature*, **480**, 379–382.
109. Baldauf, H.M., Pan, X., Erikson, E., Schmidt, S., Daddacha, W., Burggraf, M., Schenkova, K., Ambiel, I., Wabnitz, S., Gramberg, T. *et al.* (2012) SAMHD1 restricts HIV-1 infection in resting CD4(+) T cells. *Nat. Med.*, **18**, 1682–1687.
110. Powell, R.D., Holland, P.J., Hollis, T. and Perrino, F.W. (2011) Aicardi-Goutieres syndrome gene and HIV-1 restriction factor SAMHD1 is a dGTP-regulated deoxynucleotide triphosphohydrolase. *J. Biol. Chem.*, **286**, 43596–43600.
111. Lahouassa, H., Daddacha, W., Hofmann, H., Ayinde, D., Logue, E.C., Dragin, L., Bioch, N., Maudet, C., Bertrand, M., Granberg, T. *et al.* (2012) SAMHD1 restricts the replication of human immunodeficiency virus type 1 by depleting the intracellular pool of deoxynucleoside triphosphate. *Nat. Immunol.*, **13**, 223–228.
112. Zhao, K., Du, J., Han, X., Goodier, J.L., Zhou, X., Wei, W., Evans, S.L., Li, L., Zhang, W., Cheung, L.E. *et al.* (2013) Modulation of LINE-1 and Alu/SVA retrotransposition by Aicardi-Goutieres syndrome-related SAMHD1. *Cell Rep.*, **4**, 1108–1115.
113. Jones, R.B., Song, H., Xu, Y., Garrison, K.E., Buzdin, A.A., Anwar, N., Hunter, D.V., Mujib, S., Mihajlovic, V., Martin, E. *et al.* (2013) LINE-1 retrotransposable element DNA accumulates in HIV-1-infected cells. *J. Virol.*, **87**, 13307–13320.
114. Iijima, K., Okudaira, N., Tamura, M., Doi, A., Saito, Y., Shimura, M., Goto, M., Matsunaga, A., Kawamura, Y.I., Otsubo, T. *et al.* (2013) Viral protein R of human immunodeficiency virus type-1 induces retrotransposition of long interspersed element-1. *Retrovirology*, **10**, 83.
115. Ariumi, Y., Kaida, A., Lin, J.Y., Hirota, M., Masui, O., Yamaoka, S., Taya, Y. and Shimotohno, K. (2000) HTLV-1 Tax oncoprotein represses the p53-mediated trans-activation function through coactivator CBP sequestration. *Oncogene*, **19**, 1491–1499.
116. Ueda, Y., Hijikata, M., Takagi, S., Chiba, T. and Shimotohno, K. (1999) New p73 variants with altered C-terminal structures have varied transcriptional activities. *Oncogene*, **18**, 4993–4998.
117. Kaida, A., Ariumi, Y., Ueda, U., Lin, J.Y., Hijikata, M., Ikawa, S. and Shimotohno, K. (2000) Functional impairment of p73 and p51, the p53-related proteins, by the human T-cell leukemia virus type 1 Tax oncoprotein. *Oncogene*, **19**, 827–830.
118. Kawata, S., Ariumi, Y. and Shimotohno, K. (2003) p21 (Waf1/Cip1/Sdi1) prevents apoptosis as well as stimulates growth in cells transformed or immortalized by human T-cell leukemia virus type 1-encoded tax. *J. Virol.*, **77**, 7291–7299.
119. Matsuoka, S., Huang, M. and Elledge, S.J. (1998) Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science*, **282**, 1893–1897.
120. Matsuoka, S., Rotman, G., Ogawa, A., Shiloh, Y., Tamai, K. and Elledge, S.J. (2000) Ataxia-telangiectasia mutated phosphorylates Chk2 in vivo and in vitro. *Proc. Natl. Acad. Sci. U.S.A.*, **97**, 10389–10394.
121. Ariumi, Y., Kuroki, M., Dansako, H., Abe, K., Ikeda, M., Wakita, T. and Kato, N. (2008) The DNA damage sensors ataxia-telangiectasia mutated kinase and checkpoint kinase 2 are required for hepatitis C virus RNA replication. *J. Virol.*, **82**, 9639–9646.
122. Canman, C.E., Lim, D.S., Cimprich, K.A., Taya, Y., Tamai, K., Sakaguchi, K., Apella, E., Kastan, M.B. and Siliciano, J.D. (1998) Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. *Science*, **281**, 1677–1679.
123. Brés, V., Kiernan, R.E., Linares, L.K., Chable-Bessia, C., Plechakova, O., Treand, C., Emiliani, S., Peloponese, J.M., Jeang, K.T., Coux, O. *et al.* (2003) A non-proteolytic role for ubiquitin in Tat-mediated transactivation of the HIV-1 promoter. *Nat. Cell Biol.*, **5**, 754–761.