Enantioselective Access to S-Chiral 1,2-Benzothiazines by Cp^xRh(III)-Catalyzed C–H Functionalization of Sulfoximines

Yang Sun and Nicolai Cramer*

Laboratory of Asymmetric Catalysis and Synthesis
École Polytechnique Fédérale de Lausanne (EPFL)

SB ISIC LCSA BCH 4305
CH-1015 Lausanne, Switzerland

E-Mail: nicolai.cramer@epfl.ch

Supplementary Information

Table of Content:

General Methods	page 02
Experimental Section	page 03
X-ray Crystallographic Data	page 46
Spectra	page 58

General Methods:

All other reagents were used as obtained from the suppliers. Flash Column Chromatography was performed with Fluka silica gel 60 (0.040-0.063 μm grade). Analytical thin-layer chromatography was performed with commercial glass plates coated with 0.25 mm silica gel (E. Merck, Kieselgel 60 F254). Compounds were visualized by UV-light at 254 nm and by dipping the plates in an ethanolic vanillin/sulfuric acid solution or an aqueous potassium permanganate solution followed by heating. Proton nuclear magnetic resonance (^1H NMR) data were acquired on a Bruker AV400 (400 MHz) and Bruker AV600 (600 MHz). Chemical shifts are reported in delta (δ) units, in parts per million (ppm) relative to residual chloroform (s, 7.26 ppm), residual methylene chloride (t, 5.32 ppm), or residual dimethyl sulfoxide (quint, 2.50 ppm). Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; sept, septet; m, multiplet. Coupling constants (J) are given in Hz. Proton decoupled Carbon-13 nuclear magnetic resonance (^{13}C NMR) data were acquired at 101 MHz on a Bruker AV400 and at 151 MHz on a Bruker AV600. Chemical shifts are reported in ppm relative to CDCl_3 (77.16 ppm), CD_2Cl_2 (53.84 ppm), or $\text{DMSO}-d_6$ (39.52 ppm). Proton decoupled Phosphorus-31 nuclear magnetic resonance (^{31}P NMR) data were acquired at 162 MHz on a Bruker AV400 spectrometer. Proton decoupled Fluorine-19 nuclear magnetic resonance (^{19}F NMR) data were acquired at 376 MHz on a Bruker AV400 spectrometer. Infrared (IR) data were recorded on an Alpha-P Bruker FT-IR Spectrometer. Absorbance frequencies are reported in reciprocal centimeters (cm^{-1}). HRMS measurements were performed by an Agilent LC-MS TOF. High resolution mass are given in m/z . Optical rotations were measured on a Polartronic M polarimeter using a 0.5 cm cell with a Na 589 nm filter. X-ray analysis was performed by Dr. R. Scopelliti and Dr. F. FADAEI at the EPF Lausanne.

Experimental Section

General procedure for preparation of rhodium catalysts

Rhodium catalysts **Rh1-Rh6** were prepared following the reported procedure by Cramer *et. al.*^[1] Rh(III) dimer complexes were prepared from their corresponding Rh(I) diethylene complexes. The Rh(I) ethylene complex was dissolved in DCM, then treated with HCl (10.0 equiv. 2.0 M in diethyl ether). The resulting dark red solution was stirred for 1 more minute, then the volatiles were removed by evaporation under reduced pressure to afford the Rh(III) dimer complex without further purification.

Complex Rh5:

obtained as dark red solid in 97% yield. $^1\text{H NMR}$ (600 MHz, Methylene Chloride- d_2) δ = 8.06 (d, J = 8.2 Hz, 1H), 7.81 (d, J = 8.1 Hz, 1H), 7.50 (t, J = 7.4 Hz, 1H), 7.39 (t, J = 7.2 Hz, 1H), 7.34 (s, 1H), 7.25 (s, 1H), 7.16 – 7.02 (m, 3H), 6.93 (d, J = 8.4 Hz, 1H), 5.28 (s, 1H), 5.21 (s, 1H), 3.95 (s, 3H), 3.81 (d, J = 15.8 Hz, 1H), 3.75 (s, 3H), 3.69 (d, J = 14.3 Hz, 1H), 3.04 (d, J = 14.3 Hz, 1H), 2.50 (sept, J = 6.9 Hz, 1H), 1.95 (d, J = 15.8 Hz, 1H), 1.31 (d, J = 6.8 Hz, 3H), 1.19 (d, J = 6.9 Hz, 3H) ppm; $^{13}\text{C NMR}$ (151 MHz, CD_2Cl_2) δ = 157.13, 155.83, 137.75, 137.27, 134.50, 134.24, 127.59, 127.24, 127.17, 127.15, 127.13, 127.08, 126.76, 126.67, 126.56, 124.72, 124.38, 123.83, 106.36, 106.20, 103.79 (d, $J_{\text{C-Rh}}$ = 6.0 Hz), 103.66 (d, $J_{\text{C-Rh}}$ = 8.0 Hz), 90.86 (d, $J_{\text{C-Rh}}$ = 8.2 Hz), 84.95 (d, $J_{\text{C-Rh}}$ = 6.7 Hz), 81.98 (d, $J_{\text{C-Rh}}$ = 3.4 Hz), 56.13, 55.98, 26.06, 25.35, 24.24, 22.51, 20.73 ppm; **IR (ATR):** $\tilde{\nu}$ = 3650, 3483, 3059, 3002, 2960, 2932, 2828, 1618, 1595, 1573, 1502, 1481, 1451, 1422, 1410, 1388, 1363, 1328, 1296, 1262, 1236, 1225, 1196, 1165, 1149, 1113, 1078, 1019, 944, 930, 904, 878, 863, 831, 787, 770, 745, 700 cm^{-1} ; **HRMS (ESI)** calculated for $[\text{C}_{32}\text{H}_{28}\text{O}_2\text{ClRh}]^+$: 583.0906, found: 583.0900; $[\alpha]_{\text{D}}^{20}$ = +10.2 (1.0 in CH_2Cl_2).

Complex Rh6:

obtained as dark red solid in 91% yield. $^1\text{H NMR}$ (400 MHz, Methylene Chloride- d_2) δ = 8.07 (d, J = 8.2 Hz, 1H), 7.80 (d, J = 8.2 Hz, 1H), 7.50 (t, J = 7.7 Hz, 1H), 7.39 (ddd, J = 8.1, 6.5, 1.4 Hz, 1H), 7.32 (s, 1H), 7.25 (s, 1H), 7.15 – 7.00 (m, 3H), 6.93 (d, J = 8.4 Hz, 1H), 5.37 (s, 1H), 5.19 (s, 1H), 3.95 (s, 3H), 3.78 (d, J = 15.8 Hz, 1H), 3.72 (d, J = 14.2 Hz, 1H), 3.64 (s, 3H), 3.11 (d, J = 14.2 Hz, 1H), 1.91 (d, J = 15.8 Hz, 1H), 1.31 (s, 9H) ppm; $^{13}\text{C NMR}$ (101 MHz, CD_2Cl_2) δ = 157.24, 155.86, 137.64, 137.21, 134.52, 134.24, 127.62, 127.19, 127.16, 127.11, 127.06, 126.75, 126.72, 126.64, 124.56, 124.34, 123.79, 106.41, 106.24, 102.13, 84.95, 81.25, 56.11, 55.92, 31.32, 30.06, 25.48, 24.38 ppm; **IR (ATR):** $\tilde{\nu}$ = 3059, 2958, 2925, 2853, 1722, 1619, 1596, 1573, 1502, 1485, 1454, 1423, 1411, 1389, 1370, 1328, 1295, 1262, 1229, 1197, 1165, 1149, 1114, 1077, 1020, 944, 917, 904, 878, 862, 830, 787, 770, 745 cm^{-1} ; **HRMS (ESI)** calculated for $[\text{C}_{33}\text{H}_{31}\text{O}_2\text{ClRh}]^+$: 597.1062, found: 597.1058; $[\alpha]_D^{20}$ = +30.6 (1.0 in CH_2Cl_2).

General procedure for preparation of chiral carboxylic acids

Chiral carboxylic acids **A1-A5** were prepared following the reported procedure by Hashimoto *et al.*^[2] All spectra were in agreement with the reported data for **A1-2**^[3a], **A3**^[3b], **A4-5**^[3c].

General procedure for preparation of sulfoximines substrates

Diaryl sulfoximines were prepared following the reported procedure.^[4] In a flame-dried round bottom flask, to a solution of thionyl chloride (5.0 mmol, 0.37 mL) in THF (5.0 mL), aryl bromide solution (2.0 equiv., fresh prepared from the corresponding aryl bromide) was added at 0 °C. The reaction mixture was allowed to warm up to room temperature. After the reaction was complete (monitored by TLC), sat. brine was added to quench the reaction. The aqueous phase was washed with AcOEt (20 mL X 3), and the combined organic phase was dried over MgSO_4 . The solvent was removed by evaporation under reduced pressure. The residue was purified by column

chromatography (pentane/AcOEt). Then the pure sulfide was mixed with PhI(OAc)₂ (2.5 equiv.) and ammonium carbamate (2.0 equiv.). The mixture was dissolved in MeOH and stirred at room temperature for overnight. After the sulfide was fully consumed (monitored by TLC), the solvent was removed by evaporation under reduced pressure. The residue was redissolved in DCM, and washed with water. The organic layer was dried over MgSO₄, and solvent was removed. The residue was purified by column chromatography (pentane/AcOEt).

Iminodiphenyl- λ^6 -sulfanone (1a)^[4]

obtained as a white solid in 86% yield. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.01 – 7.93 (m, 4H), 7.61 – 7.49 (m, 6H), 4.97 (s, 1H) ppm; ¹³C NMR (101 MHz, DMSO) δ = 143.96, 132.43, 129.23, 127.61 ppm; IR (ATR): $\tilde{\nu}$ = 3260, 3062, 1580, 1474, 1444, 1222, 1158, 1126, 1093, 1065, 1023, 998, 957, 852, 764, 752, 719 cm⁻¹; flash column chromatography with pentane:AcOEt = 2:1 as eluent; HRMS (ESI) calculated for [C₁₂H₁₂NOS]⁺: 218.0634, found: 218.0635; m.p.:102.5-104.4 °C.

Iminobis(4-methoxyphenyl)- λ^6 -sulfanone (1b)^[9]

obtained as a white solid in 85% yield. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 7.92 – 7.79 (m, 4H), 7.11 – 6.99 (m, 4H), 4.63 (s, 1H), 3.78 (s, 6H) ppm; ¹³C NMR (101 MHz, DMSO) δ = 162.05, 136.21, 129.49, 114.29, 55.63 ppm; IR (ATR): $\tilde{\nu}$ = 3324, 3095, 3068, 2998, 2976, 2947, 2919, 2900, 2838, 1591, 1575, 1493, 1460, 1438, 1411, 1309, 1249, 1214, 1175, 1127, 1111, 1094, 1070, 1018, 975, 844, 830, 817, 797, 716 cm⁻¹; flash column chromatography with pentane:AcOEt = 1:1 as eluent; HRMS (ESI) calculated for [C₁₄H₁₆NO₃S]⁺: 278.0845, found: 278.0848; m.p.:137.5-138.2 °C.

Bis(4-bromophenyl)(imino)- λ^6 -sulfanone (1c)

obtained as a white solid in 78% yield. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ = 7.91 – 7.83 (m, 4H), 7.79 – 7.71 (m, 4H), 5.25 (s, 1H) ppm; $^{13}\text{C NMR}$ (101 MHz, DMSO) δ = 142.83, 132.39, 129.74, 126.77 ppm; **IR (ATR):** $\tilde{\nu}$ = 3294, 3081, 3012, 1567, 1465, 1385, 1227, 1123, 1088, 1061, 1006, 940, 817, 738, 721, 703 cm^{-1} ; flash column chromatography with pentane:AcOEt = 2:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{12}\text{H}_{10}^{79/81}\text{Br}_2\text{NOS}]^+$: 375.8824, found: 375.8826; m.p.: 137.9–139.0 °C.

Bis(4-fluorophenyl)(imino)- λ^6 -sulfanone (1d)

obtained as a colorless oil in 65% yield. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ = 8.06 – 7.98 (m, 4H), 7.43 – 7.34 (m, 4H), 5.13 (s, 1H) ppm; $^{13}\text{C NMR}$ (101 MHz, $\text{DMSO}-d_6$) δ = 164.27 (d, $J_{\text{C-F}}$ = 251.5 Hz), 140.14 (d, $J_{\text{C-F}}$ = 2.8 Hz), 130.66 (d, $J_{\text{C-F}}$ = 9.7 Hz), 116.35 (d, $J_{\text{C-F}}$ = 22.7 Hz) ppm; $^{19}\text{F NMR}$ (376 MHz, DMSO) δ = -107.17 ppm; **IR (ATR):** $\tilde{\nu}$ = 3263, 3101, 3066, 1585, 1485, 1403, 1221, 1152, 1127, 1091, 1012, 973, 831, 813, 706 cm^{-1} ; flash column chromatography with pentane:AcOEt = 4:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{12}\text{H}_{10}\text{F}_2\text{NOS}]^+$: 254.0446, found: 254.0451.

Iminobis(4-(trifluoromethyl)phenyl)- λ^6 -sulfanone (1e)^[9]

obtained as a light yellow solid in 65% yield. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ = 8.20 (d, J = 8.1 Hz, 4H), 7.95 (d, J = 8.3 Hz, 4H), 5.59 (s, 1H) ppm; $^{13}\text{C NMR}$ (101 MHz, $\text{DMSO}-d_6$) δ = 146.99, 132.64 (q, $J_{\text{C-F}}$ = 32.3 Hz), 128.92, 126.60 (q, $J_{\text{C-F}}$ = 3.7 Hz), 123.43 (q, $J_{\text{C-F}}$ = 272.8 Hz) ppm; $^{19}\text{F NMR}$ (376 MHz, DMSO) δ = -61.79 ppm; **IR (ATR):** $\tilde{\nu}$ = 3341, 3315, 3286, 3210, 3098, 3052, 1605, 1403, 1316, 1239, 1166, 1125, 1102, 1057, 1015, 980, 842, 784, 731, 710 cm^{-1} ; flash column chromatography with pentane:AcOEt = 4:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{14}\text{H}_{10}\text{F}_6\text{NOS}]^+$: 354.0382, found: 354.0382; m.p.: 101.6–103.2 °C.

Bis(4-chlorophenyl)(imino)- λ^6 -sulfanone (1f)^[9]

obtained as an off white solid in 51% yield. **¹H NMR** (400 MHz, DMSO-*d*₆) δ = 8.02 – 7.87 (m, 4H), 7.68 – 7.56 (m, 4H), 5.25 (s, 1H) ppm; **¹³C NMR** (101 MHz, DMSO) δ = 142.43, 137.70, 129.62, 129.40 ppm; **IR (ATR):** $\tilde{\nu}$ = 3335, 3306, 3254, 3082, 1916, 1572, 1472, 1391, 1237, 1168, 1132, 1082, 1008, 939, 820, 752, 739, 706 cm⁻¹; flash column chromatography with pentane:AcOEt = 2:1 as eluent; **HRMS (ESI)** calculated for [C₁₂H₁₀Cl₂NOS]⁺: 285.9855, found: 285.9856; m.p.:108.7-110.1 °C.

Bis(3-chlorophenyl)(imino)-λ⁶-sulfanone (1g)

obtained as an off white solid in 66% yield. **¹H NMR** (400 MHz, DMSO-*d*₆) δ = 8.02 (t, *J* = 1.9 Hz, 2H), 7.94 (dt, *J* = 7.9, 1.4 Hz, 2H), 7.70 (ddd, *J* = 8.0, 2.2, 1.0 Hz, 2H), 7.60 (t, *J* = 7.9 Hz, 2H), 5.42 (s, 1H) ppm; **¹³C NMR** (101 MHz, DMSO) δ = 145.27, 133.88, 132.77, 131.41, 127.33, 126.63 ppm; **IR (ATR):** $\tilde{\nu}$ = 3091, 1577, 1452, 1412, 1235, 1137, 1118, 1070, 1012, 988, 970, 884, 783 cm⁻¹; flash column chromatography with pentane:AcOEt = 2:1 as eluent; **HRMS (ESI)** calculated for [C₁₂H₁₀Cl₂NOS]⁺: 285.9855, found: 285.9855; m.p.:81.7-83.8 °C.

Iminodi-*m*-tolyl-λ⁶-sulfanone (1h)

obtained as an off white solid in 88% yield. **¹H NMR** (400 MHz, DMSO-*d*₆) δ = 7.83 – 7.68 (m, 4H), 7.49 – 7.36 (m, 4H), 4.83 (s, 1H), 2.35 (s, 6H) ppm; **¹³C NMR** (101 MHz, DMSO) δ = 143.95, 138.85, 132.96, 129.03, 127.71, 124.82, 20.83 ppm; **IR (ATR):** $\tilde{\nu}$ = 3270, 3081, 3050, 3018, 2918, 2857, 1598, 1474, 1453, 1419, 1226, 1170, 1116, 1091, 1060, 1004, 973, 932, 914, 894, 864, 848, 795, 786 cm⁻¹; flash column chromatography with pentane:AcOEt = 3:1 as eluent; **HRMS (ESI)** calculated for [C₁₄H₁₆NOS]⁺: 246.0947, found: 246.0948; m.p.:92.7-93.9 °C.

Iminodi-*o*-tolyl-λ⁶-sulfanone (1i)

obtained as a white solid in 38% yield. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ = 8.18 (dd, J = 7.9, 1.6 Hz, 2H), 7.50 (dtd, J = 23.4, 7.5, 1.5 Hz, 4H), 7.34 – 7.23 (m, 2H), 5.09 (s, 1H), 2.16 (s, 6H) ppm; $^{13}\text{C NMR}$ (101 MHz, DMSO) δ = 140.77, 136.57, 132.65, 132.54, 128.98, 126.04, 19.53 ppm; **IR (ATR):** $\tilde{\nu}$ = 3311, 3282, 3255, 3057, 3016, 2964, 2927, 1591, 1568, 1453, 1377, 1271, 1215, 1188, 1162, 1137, 1105, 1061, 1033, 951, 805, 764, 756, 704 cm^{-1} ; flash column chromatography with pentane:AcOEt = 3:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{14}\text{H}_{16}\text{NOS}]^+$: 246.0947, found: 246.0946; m.p.:136.5-137.6 °C.

Imino(phenyl)(*p*-tolyl)- λ^6 -sulfanone (4)

obtained as a white solid in 78% yield. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ = 8.01 – 7.91 (m, 2H), 7.85 (d, J = 8.2 Hz, 2H), 7.59 – 7.48 (m, 3H), 7.33 (d, J = 8.1 Hz, 2H), 4.88 (s, 1H), 2.31 (s, 3H) ppm; $^{13}\text{C NMR}$ (101 MHz, DMSO) δ = 144.33, 142.75, 141.02, 132.22, 129.66, 129.14, 127.67, 127.43, 20.85

ppm; **IR (ATR):** $\tilde{\nu}$ = 3263, 3062, 2972, 2917, 1593, 1581, 1488, 1474, 1447, 1393, 1226. 1167, 1112, 1092, 1062, 1015, 1001, 978, 840, 811, 754, 710 cm^{-1} ; flash column chromatography with pentane:AcOEt = 3:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{13}\text{H}_{14}\text{NOS}]^+$: 232.0791, found: 232.0790; m.p.:103.5-105.0 °C.

General procedure for preparation of diazo compounds

Diazo compounds **2a-2c**^[5], **2f-2l**^[5], **2d**^[6], **2g**^[6], **2e**^[7], **2m-2n**^[8], **2o**^[9], **2p**^[10] were prepared following the procedure in the given reference. All spectra were in agreement with the reported data.

Ethyl 2-diazo-3-oxobutanoate (2a)

$^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ = 4.30 (q, J = 7.1 Hz, 2H), 2.47 (s, 3H), 1.33 (t, J = 7.1 Hz, 3H) ppm.

Benzyl 2-diazo-3-oxobutanoate (2b)

^1H NMR (400 MHz, Chloroform-*d*) δ = 7.45 – 7.32 (m, 5H), 5.27 (s, 2H), 2.49 (s, 3H) ppm.

Dimethyl (1-diazo-2-oxopropyl)phosphonate (2c)

^1H NMR (400 MHz, Chloroform-*d*) δ = 3.84 (dd, J = 11.9, 0.9 Hz, 6H), 2.26 (s, 3H) ppm.

1-Diazo-1-tosylpropan-2-one (2d)

^1H NMR (400 MHz, Chloroform-*d*) δ = 7.82 (d, J = 8.4 Hz, 2H), 7.35 (d, J = 8.3 Hz, 2H), 2.43 (s, 3H), 2.24 (s, 3H) ppm.

Ethyl (*E*)-2-diazo-3-oxo-5-phenylpent-4-enoate (2e)

^1H NMR (400 MHz, Chloroform-*d*) δ = 7.88 (d, J = 15.8 Hz, 1H), 7.77 (d, J = 15.8 Hz, 1H), 7.66 – 7.60 (m, 2H), 7.39 (ddt, J = 5.5, 4.1, 2.2 Hz, 3H), 4.35 (q, J = 7.1 Hz, 2H), 1.37 (t, J = 7.1 Hz, 3H) ppm.

Ethyl 2-diazo-3-oxopropanoate (2f)

^1H NMR (400 MHz, Chloroform-*d*) δ = 9.70 (s, 1H), 4.36 (q, J = 7.1 Hz, 2H), 1.35 (t, J = 7.1 Hz, 3H) ppm.

4-Diazoheptane-3,5-dione (2g)

¹H NMR (400 MHz, Chloroform-*d*) δ = 2.75 (q, J = 7.3 Hz, 2H), 1.15 (t, J = 7.3 Hz, 3H) ppm.

Ethyl 2-diazo-3-oxo-3-phenylpropanoate (2h)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.67 – 7.59 (m, 2H), 7.57 – 7.48 (m, 1H), 7.42 (t, J = 7.6 Hz, 2H), 4.25 (q, J = 7.1 Hz, 2H), 1.25 (t, J = 7.1 Hz, 3H) ppm.

Ethyl 2-diazo-3-(4-methoxyphenyl)-3-oxopropanoate (2i)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.77 – 7.61 (m, 2H), 6.99 – 6.87 (m, 2H), 4.26 (q, J = 7.1 Hz, 2H), 3.86 (s, 3H), 1.28 (t, J = 7.1 Hz, 3H) ppm.

Ethyl 2-diazo-3-(4-fluorophenyl)-3-oxopropanoate (2j)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.75 – 7.62 (m, 2H), 7.18 – 7.02 (m, 2H), 4.26 (q, J = 7.1 Hz, 2H), 1.27 (t, J = 7.1 Hz, 3H) ppm.

Ethyl 2-diazo-3-(furan-2-yl)-3-oxopropanoate (2k)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.64 – 7.59 (m, 1H), 7.51 (dd, J = 3.6, 0.7 Hz, 1H), 6.56 (dd, J = 3.6, 1.7 Hz, 1H), 4.34 (q, J = 7.1 Hz, 2H), 1.35 (t, J = 7.1 Hz, 3H) ppm.

Ethyl 3-cyclopropyl-2-diazo-3-oxopropanoate (2l)

¹H NMR (400 MHz, Chloroform-*d*) δ = 4.32 (q, J = 7.1 Hz, 2H), 3.10 (tt, J = 7.9, 4.6 Hz, 1H), 1.34 (t, J = 7.1 Hz, 3H), 1.26 – 1.14 (m, 2H), 1.04 – 0.94 (m, 2H) ppm.

1-Diazo-1-phenylpropan-2-one (2m)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.50 (d, J = 7.3 Hz, 2H), 7.45 – 7.38 (m, 2H), 7.29 – 7.22 (m, 1H), 2.37 (s, 3H) ppm.

2-Diazo-1-(furan-2-yl)-2-phenylethan-1-one (2n)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.64 – 7.55 (m, 2H), 7.54 – 7.50 (m, 1H), 7.44 (t, J = 7.8 Hz, 2H), 7.33 – 7.26 (m, 1H), 7.14 (d, J = 3.6 Hz, 1H), 6.54 (dd, J = 3.5, 1.7 Hz, 1H) ppm.

3-Diazo-4-phenylbutan-2-one (2o)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.39 – 7.29 (m, 2H), 7.29 – 7.25 (m, 1H), 7.25 – 7.18 (m, 2H), 3.65 (s, 2H), 2.27 (s, 3H) ppm.

2-Diazo-1-phenylethan-1-one (2p)

¹H NMR (400 MHz, Chloroform-*d*) δ = 7.83 – 7.71 (m, 2H), 7.60 – 7.51 (m, 1H), 7.45 (dd, J = 8.3, 6.9 Hz, 2H), 5.90 (s, 1H) ppm.

General procedure for Rh(III)-catalyzed asymmetric synthesis of annulated sulfoximines **3**

Without protection from oxygen or moisture, the mixture of **Rh5** (3.1 mg, 2.5 μ mol), chiral carboxylic acid **A4** (9.3 mg, 30.0 μ mol), NaSbF₆ (12.9 mg, 50.0 μ mol), diaryl sulfoximine **1** (1 equiv., 0.10 mmol) and diazo compound **2** (1 equiv., 0.10 mmol) in *t*BuOH (500 μ L) were stirred at 35 °C. After the sulfoximine were fully consumed (monitored by TLC), the solvent was evaporated under reduced pressure. The reaction mixture was purified by column chromatography (pentane/EtOAc) on silica gel to afford the product **3**.

Ethyl (*R*)-3-methyl-1-phenyl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3aa**)**

obtained as a colorless oil in 92% yield (30.0 mg) after 30 h reaction. ¹H NMR (400 MHz, Chloroform-*d*) δ = 7.97 – 7.88 (m, 2H), 7.76 – 7.69 (m, 1H), 7.70 – 7.61 (m, 1H), 7.63 – 7.54 (m, 2H), 7.51 (ddd, *J* = 8.4, 6.8, 1.6 Hz, 1H), 7.31 – 7.18 (m, 2H), 4.42 (q, *J* = 7.2 Hz, 2H), 2.47 (s, 3H) 1.42 (t, *J* = 7.2 Hz, 3H) ppm; ¹³C NMR (101 MHz, CDCl₃) δ = 169.03, 151.87, 139.58, 134.15, 133.89, 132.62, 129.32, 129.26, 126.13, 125.21, 124.41, 118.03, 105.65, 61.09, 25.14, 14.45 ppm; IR (ATR): $\tilde{\nu}$ = 3063, 2980, 2928, 2905, 1707, 1596, 1567, 1522, 1472, 1446, 1425, 1369, 1348, 1310, 1281, 1261, 1220, 1202, 1168, 1137, 1102, 1076, 1058, 1023, 998, 957, 876, 824, 775, 754, 742 cm⁻¹; flash column chromatography with pentane:AcOEt = 5:1 as eluent; HRMS (ESI) calculated for [C₁₈H₁₈NO₃S]⁺: 328.1002, found: 328.1008; [α]_D²⁰ = +27.3 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t*_r (major) = 12.6 min, *t*_r (minor) = 13.6 min), 96 : 4 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	12.476	BV	0.3187	6551.39111	300.08505	49.5947
2	13.570	VB	0.3474	6658.47607	280.07114	50.4053

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	12.635	VV	0.3118	5466.53320	255.29948	95.6228
2	13.753	VV	0.2476	250.23431	12.18417	4.3772

Ethyl (R)-6-methoxy-1-(4-methoxyphenyl)-3-methyl-1 λ^4 -benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3ba)

obtained as a colorless oil in 95% yield (39.0 mg) after 48 h reaction. ^1H NMR (400 MHz, Chloroform- d) δ = 7.85 – 7.74 (m, 2H), 7.21 (dd, J = 5.7, 3.2 Hz, 2H), 7.01 (d, J = 9.0 Hz, 2H), 6.79 (dd, J = 8.9, 2.5 Hz, 1H), 4.41 (q, J = 7.1 Hz, 2H), 3.88 (s, 3H), 3.84 (s, 3H), 2.45 (s, 3H), 1.42 (t, J = 7.1 Hz, 3H) ppm; ^{13}C NMR (101 MHz, CDCl_3) δ = 169.26, 163.88, 162.52, 153.26, 136.52, 131.48, 131.19, 127.19, 115.39, 114.45, 111.46, 105.72, 104.86, 60.91, 55.91, 55.57, 25.51, 14.47 ppm; IR (ATR): $\tilde{\nu}$ = 2975, 2937, 2841, 1701, 1592, 1568, 1495, 1463, 1444, 1399, 1370, 1354, 1315,

1260, 1214, 1191, 1172, 1149, 1136, 1098, 1081, 1064, 1024, 990, 890, 866, 833, 802, 757, 726, 701 cm^{-1} ; flash column chromatography with pentane:AcOEt = 3:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{20}\text{H}_{22}\text{NO}_5\text{S}]^+$: 388.1213, found: 388.1213; $[\alpha]_{\text{D}}^{20} = +26.7$ ($c = 1.0$, CHCl_3); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 20% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (major) = 12.5 min, t_{r} (minor) = 16.5 min), 93 : 7 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	12.558	BV	0.3543	8055.15479	337.65378	49.7177
2	16.481	VV	0.4638	8146.62207	254.24498	50.2823

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	12.511	BV	0.3732	1.74465e4	694.19708	92.8786
2	16.487	VV	0.3829	1337.69885	43.48203	7.1214

Ethyl (R)-6-bromo-1-(4-bromophenyl)-3-methyl-1 λ^4 -benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3ca)

obtained as a colorless oil in 91% yield (44.0 mg) after 72 h reaction. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 7.96 (d, *J* = 1.8 Hz, 1H), 7.73 (s, 4H), 7.35 (dd, *J* = 8.5, 1.8 Hz, 1H), 7.12 (d, *J* = 8.5 Hz, 1H), 4.42 (q, *J* = 7.1 Hz, 2H), 2.46 (s, 3H), 1.42 (t, *J* = 7.2 Hz, 3H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 168.29, 153.78, 138.44, 135.79, 132.73, 130.61, 129.85, 129.48, 128.17, 127.27, 126.63, 116.13, 104.93, 61.32, 25.42, 14.41 ppm; **IR (ATR):** $\tilde{\nu}$ = 3086, 2979, 2933, 2903, 1705, 1572, 1551, 1509, 1459, 1385, 1339, 1306, 1278, 1258, 1224, 1189, 1145, 1101, 1070, 1049, 1008, 970, 879, 822, 752, 722 cm⁻¹; flash column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for [C₁₈H₁₆^{79/81}Br₂NO₃S]⁺: 485.9192, found: 485.9191; **[α]_D²⁰** = -29.3 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t*_r (minor) = 7.6 min, *t*_r (major) = 8.9 min), 96 : 4 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	7.594	VV	0.1711	1437.59045	126.02926	50.0340
2	8.959	BV	0.2038	1435.63892	107.50364	49.9660

Ethyl (R)-6-fluoro-1-(4-fluorophenyl)-3-methyl-1 λ^4 -benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3da)

 obtained as a colorless oil in 94% yield (34.0 mg) after 72 h reaction. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 7.99 – 7.86 (m, 2H), 7.50 (dd, *J* = 11.6, 2.4 Hz, 1H), 7.34 – 7.21 (m, 3H), 6.97 (ddd, *J* = 8.9, 7.6, 2.5 Hz, 1H), 4.42 (q, *J* = 7.1 Hz, 2H), 2.47 (s, 3H), 1.42 (t, *J* = 7.1 Hz, 3H) ppm; **¹³C NMR** (101 MHz, Chloroform-*d*) δ = 168.46, 166.11 (d, *J*_{C-F} = 258.0 Hz), 164.91 (d, *J*_{C-F} = 253.1 Hz), 153.99, 137.18 (d, *J*_{C-F} = 10.9 Hz), 135.62 (d, *J*_{C-F} = 3.0 Hz), 132.00 (d, *J*_{C-F} = 9.9 Hz), 128.20 (d, *J*_{C-F} = 10.3 Hz), 116.75 (d, *J*_{C-F} = 22.8 Hz), 114.98 (d, *J*_{C-F} = 24.7 Hz), 114.24 (d, *J*_{C-F} = 2.1 Hz), 110.34 (d, *J*_{C-F} = 24.9 Hz), 105.30 (d, *J*_{C-F} = 2.9 Hz), 61.26, 25.48, 14.42 ppm; **¹⁹F NMR** (376 MHz, Chloroform-*d*) δ = -108.28, -108.72 ppm; **IR (ATR)**: $\tilde{\nu}$ = 3102, 3070, 2980, 2928, 2852, 1705, 1609, 1588, 1568, 1491, 1467, 1406, 1370, 1346, 1316, 1276, 1220, 1171, 1128, 1102, 1077, 1055, 1012, 900, 871, 838, 817, 760, 725 cm⁻¹; flash column chromatography with pentane:AcOEt = 7:1 as eluent; **HRMS (ESI)** calculated for [C₁₈H₁₆F₂NO₃S]⁺: 364.0813, found: 364.0819; **[α]_D²⁰** = +7.8 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t*_r (major) = 7.5 min, *t*_r (minor) = 10.0 min), 96 : 4 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	7.611	VV	0.1729	3993.89941	356.01556	50.1062
2	10.156	BV	0.2391	3976.97119	259.12573	49.8938

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	7.527	BV	0.1697	3746.09717	336.97174	96.1850
2	10.015	VV	0.1964	148.58083	9.40001	3.8150

Ethyl (R) - 3 -methyl-6-(trifluoromethyl)-1-(4-(trifluoromethyl)phenyl)-1 λ^4 - benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3ea)

obtained as a light yellow solid in 86% yield (40.0 mg) after 72 h reaction. ¹H NMR (400 MHz, Chloroform-*d*) δ = 8.14 (s, 1H), 8.05 (d, *J* = 8.2 Hz, 2H), 7.88 (d, *J* = 8.4 Hz, 2H), 7.46 (dd, *J* = 8.5, 1.7 Hz, 1H), 7.38 (d, *J* = 8.4 Hz, 1H), 4.46 (qd, *J* = 7.2, 1.5 Hz, 2H), 2.52 (s, 3H), 1.44 (t, *J* = 7.2 Hz, 3H) ppm; ¹³C NMR (101 MHz, Chloroform-*d*) δ = 168.01, 154.38, 142.50, 136.06 (q, *J*_{C-F} = 33.4 Hz), 134.78, 134.60 (q, *J*_{C-F} = 32.8 Hz), 129.95, 126.63 (q, *J*_{C-F} = 3.7 Hz), 126.15, 123.28 (q, *J*_{C-F} = 273.4 Hz), 123.10 (q, *J*_{C-F} = 273.3 Hz), 122.54 (q, *J*_{C-F} = 3.2 Hz), 122.43 (q, *J*_{C-F} = 4.2 Hz), 119.17,

105.94, 61.48, 25.43, 14.35 ppm; ^{19}F NMR (376 MHz, Chloroform-*d*) δ = -63.24, -63.55 ppm; IR (ATR): $\tilde{\nu}$ = 3099, 2984, 2934, 1711, 1608, 1571, 1523, 1469, 1404, 1371, 1319, 1281, 1230, 1175, 1132, 1105, 1091, 1075, 1062, 1015, 979, 889, 836, 768, 722, 706 cm^{-1} ; flash column chromatography with pentane:AcOEt = 12:1 as eluent; HRMS (ESI) calculated for $[\text{C}_{20}\text{H}_{16}\text{F}_6\text{NO}_3\text{S}]^+$: 464.0750, found: 464.0749; $[\alpha]_{\text{D}}^{20}$ = +6.7 (c = 1.0, CHCl_3); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 2% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (minor) = 12.1 min, t_{r} (major) = 17.1 min), 96 : 4 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	11.728	VV	0.2976	2240.37305	107.30575	50.1711
2	16.363	VV	0.3618	2225.09033	80.18044	49.8289

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	12.182	VV	0.2691	182.58217	8.11876	3.8984
2	17.129	VV	0.4338	4500.91992	153.61111	96.1016

Ethyl (R)-6-chloro-1-(4-chlorophenyl)-3-methyl-1 λ^4 -benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3fa)

obtained as a colorless oil in 91% yield (36.0 mg) after 72 h reaction. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 7.86 – 7.77 (m, 3H), 7.60 – 7.52 (m, 2H), 7.21 (m, 2H), 4.43 (q, *J* = 7.1 Hz, 2H), 2.47 (s, 3H), 1.43 (t, *J* = 7.1 Hz, 3H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 168.35, 153.80, 141.20, 139.55, 137.94, 135.76, 130.57, 129.73, 126.73, 124.21, 115.82, 105.08, 61.33, 25.42, 14.42 ppm; **IR (ATR):** $\tilde{\nu}$ = 3091, 2982, 2935, 2902, 1707, 1580, 1556, 1511, 1474, 1460, 1389, 1341, 1307, 1275, 1225, 1189, 1144, 1087, 1075, 1049, 1011, 978, 880, 826, 761, 740 cm⁻¹; flash column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for [C₁₈H₁₆³⁵Cl₂NO₃S]⁺: 396.0222, found: 396.0225; **[α]_D²⁰** = +3.8 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t_r* (major) = 12.3 min, *t_r* (minor) = 16.7 min), 94 : 6 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	11.329	VV	0.3266	2.14096e4	996.79681	50.4438
2	16.001	VV	0.4132	2.10329e4	715.01465	49.5562

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	12.275	VV	0.2578	4953.67432	295.17117	93.6286
2	16.672	VB	0.2848	337.09760	15.19682	6.3714

Ethyl (R)-7-chloro-1-(3-chlorophenyl)-3-methyl-1 λ ⁴-benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3ga)

obtained as a colorless oil in 61% yield (24.0 mg) after 72 h reaction. ¹H NMR (400 MHz, Chloroform-*d*) δ = 7.86 (t, *J* = 2.0 Hz, 1H), 7.81 (ddd, *J* = 7.8, 1.9, 1.1 Hz, 1H), 7.73 (d, *J* = 9.0 Hz, 1H), 7.66 (ddd, *J* = 8.0, 2.1, 1.1 Hz, 1H), 7.56 (t, *J* = 8.0 Hz, 1H), 7.48 (dd, *J* = 9.0, 2.2 Hz, 1H), 7.23 (d, *J* = 2.2 Hz, 1H), 4.42 (q, *J* = 7.0 Hz, 2H), 2.47 (s, 3H), 1.42 (t, *J* = 7.1 Hz, 3H) ppm; ¹³C NMR (101 MHz, CDCl₃)

δ = 168.45, 152.35, 140.86, 135.92, 134.46, 133.43, 132.75, 131.59, 130.65, 129.35, 127.44, 126.53, 124.28, 118.39, 105.62, 61.32, 25.18, 14.42 ppm; IR (ATR): $\tilde{\nu}$ = 3069, 2980, 2929, 2857, 1800, 1709, 1575, 1471, 1409, 1370, 1347, 1301, 1258, 1225, 1200, 1145, 1122, 1087, 1057, 1019, 996, 877, 860, 840, 787, 727 cm⁻¹; flash column chromatography with pentane:AcOEt = 7:1 as eluent; HRMS (ESI) calculated for [C₁₈H₁₆³⁵Cl₂NO₃S]⁺: 396.0222, found: 396.0218; [α]_D²⁰ = -26.0 (c = 1.0, CHCl₃); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_r (major) = 6.3 min, t_r (minor) = 8.5 min), 95 : 5 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.309	BV	0.1430	2910.67847	311.52979	50.0347
2	8.412	VB	0.1950	2906.63818	230.74068	49.9653

Ethyl (*R*)-3,7-dimethyl-1-(*m*-tolyl)-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3ha)

obtained as a colorless oil in 90% yield (32.0 mg) after 48 h reaction. $^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ = 7.74 (dd, J = 6.7, 2.1 Hz, 1H), 7.68 (s, 1H), 7.63 (d, J = 8.5 Hz, 1H), 7.51 – 7.43 (m, 2H), 7.33 (dd, J = 8.6, 1.9 Hz, 1H), 7.05 (d, J = 1.8 Hz, 1H), 4.41 (q, J = 7.1 Hz, 2H), 2.45 (s, 3H), 2.44 (s, 3H), 2.28 (s, 3H), 1.41 (t, J = 7.1 Hz, 3H) ppm; $^{13}\text{C NMR}$ (101 MHz, CDCl_3) δ = 169.19, 150.91, 139.67, 139.56, 136.31, 134.67, 134.17, 131.86, 129.65, 129.04, 126.50, 124.47, 124.41, 118.04, 105.35, 61.01, 25.06, 21.49, 21.14, 14.46 ppm; **IR (ATR):** $\tilde{\nu}$ = 2978, 2922, 2871, 1704, 1595, 1561, 1488, 1445, 1368, 1346, 1307, 1265, 1214, 1194, 1169, 1148, 1094, 1077, 1057, 997, 957, 880, 853, 825, 786, 765, 731, 714 cm^{-1} ; flash column chromatography with pentane:AcOEt = 6:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{20}\text{H}_{22}\text{NO}_3\text{S}]^+$: 356.1315, found: 356.1317; $[\alpha]_{\text{D}}^{20}$ = -9.3 (c = 1.0, CHCl_3); HPLC separation (Chiralpak ID, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (major) = 15.0 min, t_{r} (minor) = 16.0 min), 93 : 7 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.196	VB	0.3049	1275.84753	55.77475	50.4966
2	16.219	BB	0.3422	1250.75146	51.70996	49.5034

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.002	VV	0.3480	4272.87012	187.38246	92.9028
2	16.010	VB	0.3081	326.41971	13.59388	7.0972

Ethyl (*R*)-3,8-dimethyl-1-(*o*-tolyl)-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3ia)

obtained as an off white solid in 96% yield (34.0 mg) after 48 h reaction. ¹H NMR

(400 MHz, Chloroform-*d*) δ = 8.32 (dd, *J* = 7.7, 1.7 Hz, 1H), 7.57 – 7.38 (m, 4H), 7.23 (dd, *J* = 7.5, 1.6 Hz, 1H), 7.04 (d, *J* = 7.0 Hz, 1H), 4.40 (q, *J* = 7.1 Hz, 2H), 2.35 (s, 3H), 2.05 (s, 3H), 1.71 (s, 3H), 1.40 (t, *J* = 7.1 Hz, 3H) ppm; ¹³C NMR (101 MHz, CDCl₃) δ = 169.59, 149.76, 139.64, 135.93, 135.55, 133.98, 133.16, 132.83, 129.26, 128.85, 126.54, 122.69, 115.71, 105.63, 61.11, 24.47, 20.51, 19.36, 14.42 ppm; IR (ATR):

$\tilde{\nu}$ = 3061, 2979, 2929, 2872, 1708, 1577, 1533, 1468, 1404, 1365, 1348, 1303, 1274, 1246, 1213, 1197, 1138, 1092, 1026, 887, 865, 844, 806, 791, 759, 732, 709 cm⁻¹; flash column chromatography with pentane:AcOEt = 6:1 as eluent; HRMS (ESI) calculated for [C₂₀H₂₂NO₃S]⁺: 356.1315, found: 356.1319; [α]_D²⁰ = -141.8 (c = 1.0, CHCl₃); HPLC separation (Chiralpak ID, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t*_r (major) = 14.6 min, *t*_r (minor) = 16.5 min), 79 : 21 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.494	VV	0.3240	3699.95215	163.51950	50.1452
2	16.204	BB	0.3898	3678.53174	147.05231	49.8548

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.581	BV	0.3368	5879.50342	259.08182	79.1075
2	16.490	BV	0.3583	1552.79126	59.92065	20.8925

Benzyl (*R*)-3-methyl-1-phenyl-1λ⁴-benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3ab)

obtained as an off white solid in 90% yield (35.0 mg) after 48 h reaction. For ***big scale*** (1.0 mmol), 83% yield (323.0 mg) were obtained. ¹H NMR (400 MHz, Chloroform-*d*) δ = 7.96 – 7.87 (m, 2H), 7.72 – 7.63 (m, 2H), 7.61 – 7.54 (m, 2H), 7.51 – 7.45 (m, 3H), 7.44 – 7.33 (m, 3H), 7.29 – 7.18 (m, 2H), 5.40 (s, 2H), 2.44 (s, 3H) ppm; ¹³C NMR (101 MHz, CDCl₃) δ = 168.82, 152.42, 139.52, 135.87, 134.15,

133.91, 132.63, 129.30, 129.26, 128.78, 128.73, 128.51, 126.15, 125.20, 124.46, 118.01, 105.24, 67.03, 25.28 ppm; IR (ATR): $\tilde{\nu}$ = 3063, 3032, 2953, 2928, 1708, 1596, 1565, 1520, 1497, 1471, 1447, 1425, 1376, 1348, 1311, 1280, 1261, 1218, 1199, 1134, 1101, 1074, 1056, 998, 906, 868,

790, 775, 752, 741 cm^{-1} ; flash column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{23}\text{H}_{20}\text{NO}_3\text{S}]^+$: 390.1158, found: 390.1156; $[\alpha]_{\text{D}}^{20} = +11.3$ ($c = 1.0$, CHCl_3); HPLC separation (Chiralpak IF, 4.6 x 250 mm; 20% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (major) = 15.2 min, t_{r} (minor) = 16.6 min), 95 : 5 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.107	BB	0.2910	690.40375	30.39860	49.5127
2	16.445	BB	0.3364	703.99255	29.92832	50.4873

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.214	BB	0.3588	2854.09570	119.42443	94.9613
2	16.645	BV	0.2841	151.44049	6.35263	5.0387

Big scale HPLC spectra

Dimethyl (R)-(3-methyl-1-oxido-1-phenyl-1 λ^4 -benzo[e][1,2]thiazin-4-yl)phosphonate (3ac)

obtained as a colorless oil in 63% yield (23.0 mg) after 48 h reaction. ^1H NMR (400 MHz, Chloroform-*d*) δ = 8.38 – 8.28 (m, 1H), 7.97 – 7.86 (m, 2H), 7.72 – 7.66 (m, 1H), 7.63 – 7.58 (m, 2H), 7.54 (ddd, J = 8.7, 6.1, 2.6 Hz, 1H), 7.30 – 7.21 (m, 2H), 3.79 (d, J = 11.4 Hz, 3H), 3.75 (d, J = 11.4 Hz, 3H), 2.75 (d, J = 1.9 Hz, 3H) ppm;

^{13}C NMR (101 MHz, Chloroform-*d*) δ = 162.85 (d, $J_{\text{C-P}}$ = 19.4 Hz), 138.32, 136.08 (d, $J_{\text{C-P}}$ = 14.6 Hz), 134.16, 132.41, 129.37, 129.26, 126.52 (d, $J_{\text{C-P}}$ = 2.8 Hz), 126.13, 124.56, 118.65 (d, $J_{\text{C-P}}$ = 11.3 Hz), 95.32 (d, $J_{\text{C-P}}$ = 200.8 Hz), 52.21 (d, $J_{\text{C-P}}$ = 5.4 Hz), 27.39 (d, $J_{\text{C-P}}$ = 1.5 Hz) ppm; ^{31}P NMR (162 MHz, CDCl_3) δ = 24.65 ppm; IR (ATR): $\tilde{\nu}$ = 2981, 2952, 2913, 2847, 1594, 1555, 1495, 1464, 1448, 1416, 1328, 1302, 1276, 1228, 1181, 1107, 1044, 1021, 930, 856, 821, 777, 759, 747 cm^{-1} ; flash column chromatography with pentane:AcOEt = 1:2 as eluent; HRMS (ESI) calculated for $[\text{C}_{17}\text{H}_{19}\text{NO}_4\text{PS}]^+$: 364.0767, found: 364.0772; $[\alpha]_{\text{D}}^{20}$ = -25.3 (c = 1.0, CHCl_3); HPLC separation (Chiralpak IG, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (minor) = 18.5 min, t_{r} (major) = 24.0 min), 77 : 23 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	17.838	MM	0.5457	4023.81689	122.90402	49.8172
2	22.962	MM	0.7278	4053.35425	92.82514	50.1828

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	18.538	MM	0.6073	775.91791	21.29501	22.9506
2	24.010	MM	0.7377	2604.89551	58.85146	77.0494

(R)-3-Methyl-1-phenyl-4-tosylbenzo[e][1,2]thiazine 1-oxide (3ad)

obtained as a colorless oil in 93% yield (38.0 mg) after 16 h reaction at 60 °C. ¹H

NMR (400 MHz, Chloroform-*d*) δ = 8.42 (d, *J* = 8.7 Hz, 1H), 7.97 – 7.87 (m, 2H), 7.85 – 7.77 (m, 2H), 7.77 – 7.69 (m, 1H), 7.63 (dd, *J* = 8.6, 7.0 Hz, 2H), 7.49 (dq, *J* = 8.6, 4.3 Hz, 1H), 7.35 – 7.23 (m, 4H), 2.84 (s, 3H), 2.40 (s, 3H) ppm; ¹³C **NMR**

(101 MHz, CDCl₃) δ = 160.19, 143.36, 141.36, 137.85, 137.82, 134.48, 133.00, 132.70, 129.80, 129.53, 129.01, 126.59, 126.08, 125.15, 124.77, 119.20, 111.00, 27.48, 21.61 ppm; **IR (ATR):** $\tilde{\nu}$ = 3063, 3027, 2973, 2923, 1594, 1555, 1490, 1463, 1447, 1415, 1374, 1327, 1294, 1228, 1183, 1153, 1137, 1104, 1085, 1054, 1022, 998, 910, 855, 811, 758, 744, 726 cm⁻¹; flash column

chromatography with pentane:AcOEt = 4:1 as eluent; **HRMS (ESI)** calculated for $[C_{22}H_{20}NO_3S_2]^+$: 410.0879, found: 410.0886; $[\alpha]_D^{20} = -76.5$ ($c = 1.0$, $CHCl_3$); HPLC separation (Chiralpak IF, 4.6 x 250 mm; 40% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_r (major) = 17.7 min, t_r (minor) = 23.2 min), 92 : 8 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	17.774	BB	0.5466	8447.04785	232.09398	50.2366
2	23.089	BB	0.8540	8367.49316	146.58586	49.7634

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	17.657	BB	0.5272	863.63867	24.62192	92.0953
2	23.210	MM	0.8835	74.12771	1.39843	7.9047

Ethyl (*R,E*)-1-phenyl-3-styryl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3ae)

obtained as a yellow oil in 82% yield (34.0 mg) after 48 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 8.01 – 7.94 (m, 2H), 7.73 – 7.63 (m, 3H), 7.62 – 7.56 (m, 2H), 7.54 – 7.45 (m, 3H), 7.38 – 7.29 (m, 4H), 7.29 – 7.21 (m, 2H), 4.50 (qd, J = 7.2, 2.1 Hz, 2H), 1.46 (t, J = 7.1 Hz, 3H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 168.81, 146.66, 139.84, 136.83, 136.33, 133.96, 133.85, 132.54, 129.50, 129.28, 128.78, 128.62, 127.60, 126.72, 125.34, 125.12, 124.79, 120.49, 107.71, 61.50, 14.61 ppm; **IR (ATR):** $\tilde{\nu}$ = 3060, 3025, 2979, 2932, 2902, 2871, 1703, 1623, 1592, 1575, 1559, 1505, 1467, 1446, 1427, 1368, 1351, 1312, 1283, 1258, 1227, 1194, 1137, 1104, 1079, 1024, 998, 964, 908, 865, 848, 824, 776, 753, 728 cm⁻¹; flash column chromatography with pentane:AcOEt = 7:1 as eluent; **HRMS (ESI)** calculated for [C₂₅H₂₂NO₃S]⁺: 416.1315, found: 416.1317; **[α]_D²⁰** = +190.5 (c = 1.0, CHCl₃); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_r (major) = 10.2 min, t_r (minor) = 21.1 min), 95 : 5 e.r.

Ethyl (*R*)-1-phenyl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3af)

obtained as a colorless oil in 70% yield (22.0 mg) after 36 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. ¹H NMR (400 MHz, Chloroform-*d*) δ = 8.86 (d, *J* = 8.2 Hz, 1H), 8.51 (s, 1H), 7.90 – 7.83 (m, 2H), 7.72 – 7.55 (m, 4H), 7.37 (dd, *J* = 8.2, 1.6 Hz, 1H), 7.31 (ddd, *J* = 8.1, 7.0, 1.2 Hz, 1H), 4.36 (q, *J* = 6.9 Hz, 2H), 1.40 (t, *J* = 7.1 Hz, 3H) ppm; ¹³C NMR (101 MHz, CDCl₃)

δ = 166.66, 150.13, 139.86, 134.26, 134.15, 133.27, 129.38, 128.76, 126.85, 125.65, 125.62, 118.90, 102.73, 60.34, 14.59 ppm; IR (ATR): $\tilde{\nu}$ = 3064, 2979, 2932, 2903, 2870, 1691, 1595, 1565, 1520, 1471, 1446, 1429, 1390, 1365, 1278, 1249, 1212, 1176, 1157, 1131, 1095, 1072, 1042, 998, 940, 861, 823, 779, 740 cm⁻¹; flash column chromatography with pentane:AcOEt = 10:1 as eluent; HRMS (ESI) calculated for [C₁₇H₁₆NO₃S]⁺: 314.0845, found: 314.0848; [α]_D²⁰ = -172.8 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t*_r (major) = 16.0 min, *t*_r (minor) = 19.5 min), 95.5 : 4.5 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.928	MM	0.3776	534.07257	23.57295	50.1186
2	19.381	MM	0.4116	531.54407	21.52294	49.8814

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.991	BB	0.3361	1091.39185	48.20981	95.5215
2	19.502	MM	0.3771	51.16965	2.26152	4.4785

(R)-1-(3-Ethyl-1-oxido-1-phenyl-1 λ^4 -benzo[e][1,2]thiazin-4-yl)propan-1-one (3ag)

obtained as a colorless oil in 67% yield (22.0 mg) after 48 h reaction at 60 °C. ^1H

NMR (400 MHz, Chloroform-*d*) δ = 8.00 – 7.86 (m, 2H), 7.70 – 7.63 (m, 1H), 7.63 – 7.55 (m, 2H), 7.47 (ddd, J = 8.4, 6.9, 1.6 Hz, 1H), 7.31 – 7.25 (m, 1H), 7.25 – 7.15 (m, 2H), 2.82 (qd, J = 7.3, 2.5 Hz, 2H), 2.60 – 2.42 (m, 2H), 1.30 (t, J = 7.4 Hz, 3H),

1.24 (t, J = 7.3 Hz, 3H) ppm; ^{13}C **NMR** (101 MHz, CDCl_3) δ = 208.91, 151.30, 139.54, 133.84, 133.46, 132.47, 129.53, 129.24, 126.27, 125.41, 123.55, 118.55, 113.47, 39.34, 30.25, 13.45, 8.67 ppm;

IR (ATR): $\tilde{\nu}$ = 3063, 2972, 2934, 2874, 1685, 1594, 1564, 1524, 1470, 1447, 1430, 1408, 1373, 1352, 1308, 1234, 1212, 1177, 1100, 1067, 1038, 1018, 997, 944, 895, 872, 761, 746 cm^{-1} ; flash

column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for $[C_{19}H_{20}NO_2S]^+$: 326.1209, found: 326.1211; $[\alpha]_D^{20} = -36.7$ ($c = 1.0$, $CHCl_3$); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_r (major) = 9.8 min, t_r (minor) = 18.2 min), 80 : 20 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	9.862	BB	0.2342	5729.76025	375.04480	50.0267
2	18.373	BB	0.4628	5723.64355	191.46288	49.9733

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	9.799	BB	0.2371	6515.91113	424.46030	79.9459
2	18.242	BB	0.4628	1634.49048	55.30733	20.0541

Ethyl (*R*)-1,3-diphenyl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3ah)

obtained as a yellow oil in 80% yield (31.0 mg) after 72 h reaction. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 8.05 (d, J = 8.5 Hz, 1H), 8.02 – 7.94 (m, 2H), 7.65 (dd, J = 6.8, 2.8 Hz, 3H), 7.59 (tq, J = 6.4, 2.1, 1.4 Hz, 3H), 7.44 – 7.28 (m, 5H), 4.00 (ddq, J = 38.6, 10.7, 7.1 Hz, 2H), 0.85 (t, J = 7.1 Hz, 3H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 169.30, 152.73, 140.83, 139.26, 133.98, 133.89, 132.84, 129.61, 129.26, 128.96, 128.59, 128.18, 126.74, 125.21, 124.58, 118.42, 105.98, 61.12, 13.55 ppm; **IR (ATR):** $\tilde{\nu}$ = 3061, 2981, 2935, 2901, 1700, 1639, 1594, 1581, 1561, 1517, 1470, 1446, 1427, 1368, 1347, 1310, 1271, 1234, 1213, 1171, 1128, 1096, 1073, 1038, 999, 934, 913, 825, 762, 744, 727, 701 cm⁻¹; flash column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for [C₂₃H₂₀NO₃S]⁺: 390.1158, found: 390.1156; **[α]_D²⁰** = +137.2 (c = 1.0, CHCl₃); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 20% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_r (major) = 10.0 min, t_r (minor) = 10.8 min), 95: 5 e.r.

Ethyl (*R*)-3-(4-methoxyphenyl)-1-phenyl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3ai)

obtained as a yellow oil in 86% yield (36.0 mg) after 72 h reaction. ^1H NMR (400 MHz, Chloroform-*d*) δ = 8.07 – 7.98 (m, 1H), 7.98 (dd, J = 7.1, 1.5 Hz, 2H), 7.71 – 7.52 (m, 6H), 7.35 – 7.24 (m, 2H), 6.96 – 6.87 (m, 2H), 4.06 (ddq, J = 38.6, 10.7, 7.1 Hz, 2H), 3.83 (s, 3H), 0.95 (t, J = 7.1 Hz, 3H) ppm; ^{13}C NMR (101 MHz, CDCl_3) δ = 169.61, 160.46, 152.17, 139.27, 134.02, 133.93, 133.30, 132.77, 130.18, 129.62, 129.23, 126.46, 125.14, 124.42, 118.31, 113.58, 105.44, 61.13, 55.53, 13.80 ppm; IR (ATR): $\tilde{\nu}$ = 3066, 2980, 2938, 2901, 2837, 1698, 1635, 1606, 1579, 1560, 1497, 1467, 1446, 1429, 1416, 1368, 1346, 1295, 1250, 1211, 1175, 1126, 1095, 1071, 1034, 998, 922, 835, 778, 752, 730 cm^{-1} ; flash column chromatography with pentane:AcOEt = 5:1 as eluent; HRMS (ESI) calculated for $[\text{C}_{24}\text{H}_{22}\text{NO}_4\text{S}]^+$: 420.1264, found: 420.1258; $[\alpha]_{\text{D}}^{20}$ = +123.2 (c = 1.0, CHCl_3); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (major) = 10.5 min, t_{r} (minor) = 12.1 min), 91: 9 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	10.503	BB	0.2563	732.51428	43.97911	50.0186
2	12.107	BB	0.3007	731.96881	38.34032	49.9814

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	10.505	BV	0.2427	204.82986	12.02873	8.6993
2	12.112	BV	0.2923	2149.73193	113.82376	91.3007

Ethyl (*R*)-3-(4-fluorophenyl)-1-phenyl-1 λ^4 -benzo[e][1,2]thiazine-4-carboxylate 1-oxide (**3aj**)

obtained as a yellow oil in 88% yield (36.0 mg) after 72 h reaction. The

diazo compound was added slowly for 24 hours with a syringe pump. ^1H

NMR (400 MHz, Chloroform-*d*) δ = 8.04 (d, J = 8.5 Hz, 1H), 8.01 – 7.94 (m, 2H), 7.71 – 7.54 (m, 6H), 7.36 – 7.29 (m, 2H), 7.15 – 7.04 (m, 2H), 4.04

(ddq, J = 38.4, 10.8, 7.1 Hz, 2H), 0.94 (t, J = 7.2 Hz, 3H) ppm; ^{13}C **NMR** (101

MHz, Chloroform-*d*) δ = 169.18, 163.33 (d, $J_{\text{C-F}}$ = 248.6 Hz), 151.34, 139.05, 136.89 (d, $J_{\text{C-F}}$ = 3.3 Hz), 134.08, 133.75, 132.90, 130.58 (d, $J_{\text{C-F}}$ = 8.4 Hz), 129.60, 129.32, 126.87, 125.18, 124.57,

118.50, 115.13 (d, $J_{\text{C-F}}$ = 21.6 Hz), 106.08, 61.20, 13.70 ppm; ^{19}F **NMR** (376 MHz, CDCl_3) δ = -112.51

ppm; **IR (ATR):** $\tilde{\nu}$ = 3066, 2980, 2938, 2900, 1699, 1595, 1561, 1497, 1468, 1447, 1429, 1406, 1368, 1345, 1312, 1269, 1213, 1157, 1127, 1096, 1071, 1030, 998, 946, 921, 879, 840, 804, 778, 751, 727 cm^{-1} ; flash column chromatography with pentane:AcOEt = 7:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{23}\text{H}_{19}\text{FNO}_3\text{S}]^+$: 408.1064, found: 408.1065; $[\alpha]_{\text{D}}^{20}$ = +86.7 (c = 1.0, CHCl_3); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (major) = 15.8 min, t_{r} (minor) = 18.3 min), 93: 7 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.823	BB	0.3345	1345.61243	61.69125	50.2824
2	18.255	BB	0.3533	1330.49756	53.65547	49.7176

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.830	BV	0.3008	424.81265	19.45478	7.3615
2	18.272	BV	0.3859	5345.96143	213.63925	92.6385

Ethyl (*R*)-3-(furan-2-yl)-1-phenyl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (3ak)

obtained as a yellow oil in 79% yield (30.0 mg) after 72 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 8.04 – 7.96 (m, 2H), 7.77 (d, *J* = 8.4 Hz, 1H), 7.71 – 7.65 (m, 1H), 7.62 – 7.52 (m, 3H), 7.49 (dd, *J* = 1.8, 0.9 Hz, 1H), 7.33 – 7.27 (m, 2H), 6.97 (dd, *J* = 3.4, 0.9 Hz, 1H), 6.49 (dd, *J* = 3.4, 1.8 Hz, 1H), 4.39 (q, *J* = 7.2 Hz, 2H), 1.29 (t, *J* = 7.1 Hz, 3H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 169.07, 152.71, 143.90, 139.23, 138.51, 133.98, 133.49, 132.76, 129.76, 129.28, 126.84, 125.19, 124.37, 119.52, 112.86, 112.01, 104.67, 61.56, 14.36 ppm; **IR (ATR)**: $\tilde{\nu}$ = 3147, 3063, 2982, 2936, 2901, 2173, 2006, 1975, 1902, 1715, 1583, 1557, 1517, 1471, 1447, 1426, 1392, 1367, 1348, 1313, 1274, 1243, 1221, 1175, 1163, 1134, 1096, 1073, 1034, 1000, 940, 885, 867, 825, 753, 727, 703 cm⁻¹; flash column chromatography with pentane:AcOEt = 6:1 as eluent; **HRMS (ESI)** calculated for [C₂₁H₁₈NO₄S]⁺: 380.0951, found: 380.0952; **[α]_D²⁰** = +200.3 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IF, 4.6 x 250 mm; 30% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t_r* (major) = 14.3 min, *t_r* (minor) = 15.3 min), 95: 5 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.279	BV	0.3079	1111.86707	51.50311	4.9236
2	15.306	VV	0.4004	2.14707e4	801.89526	95.0764

Ethyl (*R*)-3-cyclopropyl-1-phenyl-1 λ^4 -benzo[e][1,2]thiazine-4-carboxylate 1-oxide (3al)

obtained as a yellow oil in 74% yield (26.0 mg) after 72 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 7.90 – 7.83 (m, 2H), 7.68 – 7.53 (m, 4H), 7.47 (ddd, *J* = 8.5, 7.0, 1.5 Hz, 1H), 7.28 – 7.23 (m, 1H), 7.17 (ddd, *J* = 8.1, 6.9, 1.1 Hz, 1H), 4.44 (q, *J* = 7.1 Hz, 2H), 2.29 (tt, *J* = 8.0, 4.7 Hz, 1H), 1.42 (t, *J* = 7.1 Hz, 3H), 1.26 (tdd, *J* = 6.4, 5.2, 3.0 Hz, 1H), 1.11 (dddd, *J* = 8.9, 5.9, 4.7, 3.2 Hz, 1H), 0.94 – 0.77 (m, 2H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 169.44, 155.80, 139.95, 134.16, 133.68, 132.45, 129.18, 129.16, 125.63, 125.13, 124.03, 118.34, 105.37, 61.13, 15.96, 14.47, 9.73, 7.43 ppm; **IR (ATR):** $\tilde{\nu}$ = 3089, 3064, 2979, 2928, 2901, 1703, 1596, 1562, 1516, 1470, 1446, 1433, 1369, 1353, 1308, 1275, 1262, 1228, 1208, 1187, 1138, 1110, 1071, 1040, 998, 954, 929, 866, 844, 824, 773, 754, 702 cm⁻¹; flash column chromatography with pentane:AcOEt = 11:1 as eluent; **HRMS (ESI)** calculated for [C₂₀H₂₀NO₃S]⁺: 354.1158, found: 354.1160; **[α]_D²⁰** = +132.2 (*c* = 1.0, CHCl₃); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t_r* (major) = 10.7 min, *t_r* (minor) = 12.5 min), 96: 4 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	11.208	BV	0.2356	3038.68018	197.42853	50.0746
2	13.038	BB	0.2739	3029.62793	173.31265	49.9254

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	10.655	BB	0.2789	1.11691e4	589.85651	95.9324
2	12.451	BB	0.2878	473.57742	23.43478	4.0676

(*R*)-3-Methyl-1,4-diphenylbenzo[*e*][1,2]thiazine 1-oxide (3am)

obtained as a yellow oil in 75% yield (25.0 mg) after 24 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. ¹H NMR (400 MHz, Chloroform-*d*) δ = 8.07 – 8.01 (m, 2H), 7.70 – 7.64 (m, 1H), 7.61 (dd, *J* = 8.3, 6.5 Hz, 2H), 7.47 (q, *J* = 5.0, 2.7 Hz, 2H), 7.42 – 7.36 (m, 1H), 7.35 – 7.27 (m, 4H), 7.16 (ddd, *J* = 8.2, 6.9, 1.2 Hz, 1H), 6.94 (d, *J* = 8.3 Hz, 1H), 2.12 (s, 3H) ppm;

¹³C NMR (101 MHz, CDCl₃) δ = 145.25, 140.18, 138.68, 137.01, 133.55, 131.86, 129.62, 129.13, 127.18, 125.24, 125.19, 124.95, 118.74, 111.99, 24.25 ppm; IR (ATR): $\tilde{\nu}$ = 3057, 3024, 2917, 1597, 1570, 1529, 1492, 1465, 1445, 1426, 1374, 1348, 1310, 1271, 1252, 1216, 1174, 1125, 1100, 1082,

1069, 1038, 1008, 996, 928, 912, 858, 765, 745, 702 cm^{-1} ; flash column chromatography with pentane:AcOEt = 10:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{21}\text{H}_{18}\text{NOS}]^+$: 332.1104, found: 332.1110; $[\alpha]_{\text{D}}^{20} = +102.4$ ($c = 1.0$, CHCl_3); HPLC separation (Chiralpak IA, 4.6 x 250 mm; 10% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (major) = 8.8 min, t_{r} (minor) = 10.2 min), 95: 5 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	8.751	BB	0.1961	1096.54614	86.39272	50.0579
2	10.168	BB	0.2118	1094.00879	77.92001	49.9421

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	8.785	VB	0.1942	2677.17847	210.76527	94.7242
2	10.222	VB	0.2144	149.11017	10.71417	5.2758

(R)-3-(Furan-2-yl)-1,4-diphenylbenzo[e][1,2]thiazine 1-oxide (3an)

obtained as a yellow oil in 73% yield (28.0 mg) after 24 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 8.17 – 8.07 (m, 2H), 7.72 – 7.64 (m, 1H), 7.64 – 7.57 (m, 2H), 7.55 – 7.43 (m, 3H), 7.39 – 7.28 (m, 5H), 7.21 (ddd, J = 8.1, 6.9, 1.2 Hz, 1H), 7.03 (dd, J = 8.4, 1.2 Hz, 1H), 6.18 (dd, J = 3.5, 1.8 Hz, 1H), 5.58 (d, J = 3.5 Hz, 1H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 152.21, 142.73, 139.84, 138.17, 136.98, 135.60, 133.68, 131.89, 131.81, 130.02, 129.16, 127.77, 126.17, 125.99, 124.94, 120.18, 112.83, 111.80, 111.37 ppm; **IR (ATR):** $\tilde{\nu}$ = 3141, 3116, 3058, 3023, 2925, 2854, 1766, 1594, 1577, 1562, 1521, 1460, 1446, 1427, 1391, 1347, 1311, 1261, 1237, 1204, 1160, 1135, 1097, 1070, 1025, 1015, 989, 915, 902, 885, 814, 768, 736, 702 cm⁻¹; flash column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for [C₂₄H₁₈NO₂S]⁺: 384.1053, found: 384.1055; **[α]_D²⁰** = -26.8 (c = 1.0, CHCl₃); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 20% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_r (major) = 17.6 min, t_r (minor) = 20.3 min), 94: 6 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	17.693	VV	0.4089	1969.03845	69.00771	50.2486
2	20.350	VV	0.4534	1949.55469	60.27168	49.7514

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	17.615	BB	0.4230	3435.99634	121.84868	94.1179
2	20.332	BB	0.3611	214.74092	7.09589	5.8821

(R)-4-Benzyl-3-methyl-1-phenylbenzo[e][1,2]thiazine 1-oxide (3ao)

obtained as a yellow oil in 64% yield (22.0 mg) after 24 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 8.07 – 7.95 (m, 2H), 7.69 – 7.61 (m, 1H), 7.64 – 7.55 (m, 2H), 7.42 – 7.33 (m, 2H), 7.33 – 7.25 (m, 3H), 7.25 – 7.11 (m, 4H), 4.11 (d, *J* = 2.5 Hz, 2H), 2.39 (s, 3H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 146.44, 140.43, 140.20, 136.63, 133.48, 132.16, 129.60, 129.10, 128.72, 127.92, 126.16, 125.36, 125.06, 123.52, 119.99, 105.39, 33.77, 23.62 ppm; **IR (ATR):** $\tilde{\nu}$ = 3063, 2981, 2934, 2901, 1698, 1640, 1595, 1561, 1496, 1467, 1447, 1429, 1406, 1368, 1344, 1311, 1266, 1210, 1171, 1156, 1124, 1094, 1071, 1028, 998, 945, 921, 879, 839, 803, 777, 730, 700 cm⁻¹; flash column chromatography with pentane:AcOEt = 8:1 as eluent; **HRMS (ESI)** calculated for [C₂₂H₂₀NOS]⁺: 346.1260, found: 346.1261; **[α]_D²⁰** = -3.4 (c = 1.0, CHCl₃); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 20% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; *t_r* (major) = 14.4 min, *t_r* (minor) = 19.5 min), 92: 8 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.518	VB	0.3212	1033.23706	48.37625	50.2171
2	19.643	BB	0.3727	1024.30249	35.70605	49.7829

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.408	BV	0.3243	9197.65527	442.89362	91.6174
2	19.464	BV	0.3564	841.54932	29.90088	8.3826

(R)-1,3-Diphenylbenzo[e][1,2]thiazine 1-oxide (3ap)

obtained as a yellow oil in 69% yield (22.0 mg) after 24 h reaction. The diazo compound was added slowly for 24 hours with a syringe pump. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 8.01 (ddt, *J* = 7.0, 4.1, 1.7 Hz, 4H), 7.69 – 7.60 (m, 1H), 7.62 – 7.53 (m, 2H), 7.53 – 7.37 (m, 4H), 7.40 – 7.31 (m, 2H), 7.27 – 7.19 (m, 1H), 6.82 (s, 1H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 147.43, 140.84, 139.01, 136.74,

133.46, 132.22, 129.46, 129.13, 128.93, 128.49, 127.05, 126.80, 126.39, 125.11, 119.83, 98.31 ppm; **IR (ATR):** $\tilde{\nu}$ = 3411, 3063, 2919, 2335, 2196, 2153, 2056, 2025, 1997, 1964, 1903, 1680, 1586, 1576, 1532, 1490, 1473, 1447, 1430, 1362, 1285, 1219, 1181, 1160, 1133, 1109, 1068, 1022,

998, 962, 946, 910, 859, 847, 808, 767, 741, 725 cm^{-1} ; flash column chromatography with pentane:AcOEt = 5:1 as eluent; **HRMS (ESI)** calculated for $[\text{C}_{20}\text{H}_{16}\text{NOS}]^+$: 318.0947, found: 318.0947; $[\alpha]_{\text{D}}^{20} = -11.3$ ($c = 1.0$, CHCl_3); HPLC separation (Chiralpak IC, 4.6 x 250 mm; 20% *i*-PrOH / hexane, 1.0 mL/min, 254 nm; t_{r} (minor) = 13.2 min, t_{r} (major) = 20.9 min), 94: 6 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	13.332	VB	0.2949	2213.28052	117.98558	49.9674
2	21.552	VV	0.4188	2216.17065	72.87543	50.0326

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	13.217	BV	0.2806	602.56769	33.06842	5.9599
2	20.898	BB	0.4649	9507.73340	316.08322	94.0401

Ethyl (*R*)-3-methyl-1-(*p*-tolyl)-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (5) &

Ethyl (*R*)-3,6-dimethyl-1-phenyl-1 λ^4 -benzo[*e*][1,2]thiazine-4-carboxylate 1-oxide (6)

obtained as a colorless oil in 97% yield (33.0 mg) after 48 h reaction. **¹H NMR** (400 MHz, Chloroform-*d*) δ = 7.91 – 7.84 (m, 1H), 7.82 – 7.75 (m, 1H), 7.71 (dd, J = 8.2, 1.0 Hz, 0.5H), 7.66 – 7.60 (m, 0.5H), 7.59 – 7.53 (m, 1H), 7.52 – 7.46 (m, 1H), 7.37 (d, J = 8.1 Hz, 1H), 7.29 – 7.24 (m, 0.5H), 7.24 – 7.15 (m, 1H), 7.04 (dd, J = 8.2, 1.6 Hz, 1H), 4.42 (q, J = 7.2 Hz, 1H), 4.41 (q, J = 7.1 Hz, 1H), 2.46 (s, 1.5H), 2.45 (s, 1.5H), 2.44 (s, 1.5H), 2.38 (s, 1.5H), 1.42 (t, J = 7.2 Hz, 1.5H), 1.41 (t, J = 7.1 Hz, 1.5H) ppm; **¹³C NMR** (101 MHz, CDCl₃) δ = 169.17, 169.07, 151.89, 151.63, 145.08, 143.42, 140.08, 136.35, 134.31, 133.98, 133.70, 132.44, 129.89, 129.36, 129.17, 129.08, 127.54, 126.00, 125.25, 125.10, 124.31, 124.02, 118.32, 115.51, 105.53, 105.32, 61.02, 25.13, 25.08, 22.22, 21.72, 14.42 ppm; **IR (ATR)**: $\tilde{\nu}$ 3064, 2978, 2925, 1706, 1595, 1567, 1523, 1471, 1447, 1397, 1369, 1348, 1312, 1282, 1262, 1220, 1170, 1149, 1137, 1101, 1076, 1057, 1018, 956, 875, 813, 773, 753, 737, 724 cm⁻¹; flash column chromatography with pentane:AcOEt = 5:1 as eluent; **HRMS (ESI)** calculated for [C₁₉H₂₀NO₃S]⁺: 342.1158, found: 342.1147; HPLC separation (Chiralpak IA, 4.6 x 250 mm x 4; 20% *i*-PrOH / hexane, 0.8 mL/min, 254 nm;) **6**: t_r (major) = 95.3 min, t_r (minor) = 103.2 min, 90: 10 e.r.; **5**: t_r (major) = 98.1 min, t_r (minor) = 116.4 min, 93.5: 6.5 e.r.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	95.258	MF	1.1388	9437.89551	138.12851	44.1978
2	98.078	FM	1.2030	1.01685e4	140.87538	47.6190
3	103.233	BB	0.8838	1038.69312	13.96487	4.8642
4	116.368	BB	1.0080	708.74640	8.40304	3.3191

X-ray Crystallographic Data of (R)-3ab (CCDC 1872022):

Table 1. Crystal data and structure refinement for sya03-269.

Identification code	sya03-269	
Empirical formula	C ₂₃ H ₁₉ NO ₃ S	
Formula weight	389.45	
Temperature	140.01(10) K	
Wavelength	1.54184 Å	
Crystal system	Orthorhombic	
Space group	P2 ₁ 2 ₁ 2 ₁	
Unit cell dimensions	a = 8.63582(16) Å	α = 90°.
	b = 10.3887(3) Å	β = 90°.
	c = 20.7055(5) Å	γ = 90°.
Volume	1857.60(8) Å ³	
Z	4	
Density (calculated)	1.393 Mg/m ³	
Absorption coefficient	1.752 mm ⁻¹	
F(000)	816	
Crystal size	0.475 x 0.129 x 0.094 mm ³	
Θ range for data collection	4.271 to 73.536°.	
Index ranges	-7 ≤ h ≤ 10, -12 ≤ k ≤ 12, -25 ≤ l ≤ 24	
Reflections collected	6469	
Independent reflections	3625 [<i>R</i> _{int} = 0.0286]	
Completeness to θ = 67.684°	99.7 %	

Absorption correction	Analytical
Max. and min. transmission	0.892 and 0.566
Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	3625 / 0 / 254
Goodness-of-fit on F^2	1.029
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0417$, $wR_2 = 0.1047$
R indices (all data)	$R_1 = 0.0462$, $wR_2 = 0.1082$
Absolute structure parameter	0.007(19)
Largest diff. peak and hole	0.424 and -0.315 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for sya03-269. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
S(1)	6723(1)	5129(1)	3364(1)	28(1)
O(1)	7762(3)	4944(3)	2824(1)	36(1)
O(2)	1469(3)	6802(3)	2484(1)	36(1)
O(3)	1489(3)	7609(3)	3489(1)	34(1)
N(1)	6252(3)	6509(3)	3543(2)	33(1)
C(1)	4834(4)	7032(3)	3373(2)	28(1)
C(2)	3589(4)	6305(4)	3181(2)	29(1)
C(3)	3655(3)	4912(4)	3153(1)	28(1)
C(4)	2328(4)	4136(4)	3039(2)	33(1)
C(5)	2407(5)	2818(4)	3043(2)	34(1)
C(6)	3805(4)	2178(4)	3148(2)	35(1)
C(7)	5132(4)	2893(4)	3257(2)	32(1)
C(8)	5042(4)	4230(4)	3263(2)	29(1)
C(9)	4833(4)	8462(4)	3436(2)	34(1)
C(10)	7653(4)	4517(4)	4059(2)	31(1)
C(11)	8914(4)	3708(4)	3974(2)	37(1)
C(12)	9649(5)	3240(5)	4522(2)	47(1)
C(13)	9120(5)	3553(5)	5129(2)	48(1)
C(14)	7849(5)	4359(5)	5202(2)	45(1)
C(15)	7103(4)	4843(4)	4665(2)	37(1)
C(16)	2092(4)	6925(4)	3000(2)	29(1)
C(17)	30(4)	8257(4)	3348(2)	36(1)
C(18)	-683(4)	8697(4)	3968(2)	34(1)
C(19)	-1273(4)	9932(4)	4019(2)	41(1)
C(20)	-2071(5)	10299(5)	4576(2)	47(1)
C(21)	-2263(5)	9440(5)	5077(2)	45(1)
C(22)	-1669(5)	8209(4)	5028(2)	42(1)
C(23)	-874(4)	7845(4)	4477(2)	36(1)

Table 3. Bond lengths [Å] and angles [°] for sya03-269.

S(1)-O(1)	1.446(2)
S(1)-N(1)	1.536(3)
S(1)-C(8)	1.739(3)
S(1)-C(10)	1.766(4)
O(2)-C(16)	1.203(4)
O(3)-C(16)	1.343(4)
O(3)-C(17)	1.458(4)
N(1)-C(1)	1.386(4)
C(1)-C(2)	1.372(5)
C(1)-C(9)	1.491(5)
C(2)-C(3)	1.450(5)
C(2)-C(16)	1.492(5)
C(3)-C(8)	1.411(5)
C(3)-C(4)	1.421(5)
C(4)-C(5)	1.370(6)
C(4)-H(4)	0.9500
C(5)-C(6)	1.395(6)
C(5)-H(5)	0.9500
C(6)-C(7)	1.385(5)
C(6)-H(6)	0.9500
C(7)-C(8)	1.391(5)
C(7)-H(7)	0.9500
C(9)-H(9A)	0.9800
C(9)-H(9B)	0.9800
C(9)-H(9C)	0.9800
C(10)-C(15)	1.384(5)
C(10)-C(11)	1.387(5)
C(11)-C(12)	1.387(6)
C(11)-H(11)	0.9500
C(12)-C(13)	1.377(7)
C(12)-H(12)	0.9500
C(13)-C(14)	1.388(7)
C(13)-H(13)	0.9500
C(14)-C(15)	1.379(6)

C(14)-H(14)	0.9500
C(15)-H(15)	0.9500
C(17)-C(18)	1.496(5)
C(17)-H(17A)	0.9900
C(17)-H(17B)	0.9900
C(18)-C(19)	1.385(6)
C(18)-C(23)	1.386(6)
C(19)-C(20)	1.396(6)
C(19)-H(19)	0.9500
C(20)-C(21)	1.378(7)
C(20)-H(20)	0.9500
C(21)-C(22)	1.381(7)
C(21)-H(21)	0.9500
C(22)-C(23)	1.384(6)
C(22)-H(22)	0.9500
C(23)-H(23)	0.9500

O(1)-S(1)-N(1)	118.34(18)
O(1)-S(1)-C(8)	110.68(16)
N(1)-S(1)-C(8)	108.04(16)
O(1)-S(1)-C(10)	107.47(16)
N(1)-S(1)-C(10)	105.06(18)
C(8)-S(1)-C(10)	106.53(17)
C(16)-O(3)-C(17)	115.3(3)
C(1)-N(1)-S(1)	122.6(3)
C(2)-C(1)-N(1)	123.4(3)
C(2)-C(1)-C(9)	125.0(3)
N(1)-C(1)-C(9)	111.7(3)
C(1)-C(2)-C(3)	122.1(3)
C(1)-C(2)-C(16)	120.9(3)
C(3)-C(2)-C(16)	117.0(3)
C(8)-C(3)-C(4)	115.3(3)
C(8)-C(3)-C(2)	121.9(3)
C(4)-C(3)-C(2)	122.8(3)
C(5)-C(4)-C(3)	121.7(4)
C(5)-C(4)-H(4)	119.2

C(3)-C(4)-H(4)	119.2
C(4)-C(5)-C(6)	121.4(4)
C(4)-C(5)-H(5)	119.3
C(6)-C(5)-H(5)	119.3
C(7)-C(6)-C(5)	119.1(3)
C(7)-C(6)-H(6)	120.5
C(5)-C(6)-H(6)	120.5
C(6)-C(7)-C(8)	119.4(3)
C(6)-C(7)-H(7)	120.3
C(8)-C(7)-H(7)	120.3
C(7)-C(8)-C(3)	123.2(3)
C(7)-C(8)-S(1)	119.4(3)
C(3)-C(8)-S(1)	117.3(3)
C(1)-C(9)-H(9A)	109.5
C(1)-C(9)-H(9B)	109.5
H(9A)-C(9)-H(9B)	109.5
C(1)-C(9)-H(9C)	109.5
H(9A)-C(9)-H(9C)	109.5
H(9B)-C(9)-H(9C)	109.5
C(15)-C(10)-C(11)	122.2(3)
C(15)-C(10)-S(1)	119.7(3)
C(11)-C(10)-S(1)	118.2(3)
C(10)-C(11)-C(12)	117.9(4)
C(10)-C(11)-H(11)	121.0
C(12)-C(11)-H(11)	121.0
C(13)-C(12)-C(11)	120.7(4)
C(13)-C(12)-H(12)	119.6
C(11)-C(12)-H(12)	119.6
C(12)-C(13)-C(14)	120.3(4)
C(12)-C(13)-H(13)	119.9
C(14)-C(13)-H(13)	119.9
C(15)-C(14)-C(13)	120.1(4)
C(15)-C(14)-H(14)	119.9
C(13)-C(14)-H(14)	119.9
C(14)-C(15)-C(10)	118.7(4)
C(14)-C(15)-H(15)	120.6

C(10)-C(15)-H(15)	120.6
O(2)-C(16)-O(3)	123.5(3)
O(2)-C(16)-C(2)	124.4(3)
O(3)-C(16)-C(2)	112.0(3)
O(3)-C(17)-C(18)	108.9(3)
O(3)-C(17)-H(17A)	109.9
C(18)-C(17)-H(17A)	109.9
O(3)-C(17)-H(17B)	109.9
C(18)-C(17)-H(17B)	109.9
H(17A)-C(17)-H(17B)	108.3
C(19)-C(18)-C(23)	119.3(4)
C(19)-C(18)-C(17)	120.1(4)
C(23)-C(18)-C(17)	120.4(4)
C(18)-C(19)-C(20)	119.9(4)
C(18)-C(19)-H(19)	120.1
C(20)-C(19)-H(19)	120.1
C(21)-C(20)-C(19)	120.2(4)
C(21)-C(20)-H(20)	119.9
C(19)-C(20)-H(20)	119.9
C(20)-C(21)-C(22)	120.0(4)
C(20)-C(21)-H(21)	120.0
C(22)-C(21)-H(21)	120.0
C(21)-C(22)-C(23)	119.8(4)
C(21)-C(22)-H(22)	120.1
C(23)-C(22)-H(22)	120.1
C(22)-C(23)-C(18)	120.8(4)
C(22)-C(23)-H(23)	119.6
C(18)-C(23)-H(23)	119.6

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for sya03-269. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
S(1)	19(1)	32(1)	33(1)	1(1)	-1(1)	0(1)
O(1)	25(1)	46(2)	38(1)	4(1)	3(1)	2(1)
O(2)	28(1)	45(2)	34(1)	-4(1)	-5(1)	4(1)
O(3)	23(1)	44(1)	35(1)	-3(1)	-1(1)	7(1)
N(1)	22(1)	34(2)	43(2)	-2(1)	-4(1)	-6(1)
C(1)	24(2)	31(2)	29(2)	0(1)	1(1)	2(1)
C(2)	19(2)	38(2)	28(2)	-1(1)	-1(1)	2(1)
C(3)	19(1)	36(2)	28(1)	-1(1)	-1(1)	-1(1)
C(4)	24(2)	41(2)	33(2)	-2(1)	-2(1)	-2(2)
C(5)	31(2)	38(2)	33(2)	-2(1)	-3(2)	-8(2)
C(6)	39(2)	28(2)	36(2)	-1(1)	-3(1)	-1(2)
C(7)	29(2)	35(2)	31(2)	-1(1)	-1(1)	1(2)
C(8)	26(2)	32(2)	29(2)	-2(1)	0(1)	0(1)
C(9)	26(2)	33(2)	41(2)	1(2)	-2(1)	-1(1)
C(10)	20(1)	36(2)	39(2)	3(1)	-6(1)	-4(1)
C(11)	26(2)	43(2)	44(2)	1(2)	-1(2)	1(2)
C(12)	29(2)	53(3)	59(3)	11(2)	-9(2)	6(2)
C(13)	34(2)	61(3)	49(2)	15(2)	-14(2)	-8(2)
C(14)	35(2)	62(3)	38(2)	4(2)	-3(2)	-9(2)
C(15)	25(2)	45(2)	41(2)	0(2)	0(1)	-3(2)
C(16)	24(2)	30(2)	32(2)	0(1)	1(1)	0(1)
C(17)	23(2)	46(2)	38(2)	0(2)	-2(1)	11(2)
C(18)	19(2)	44(2)	37(2)	-4(2)	-2(1)	0(2)
C(19)	29(2)	41(2)	53(2)	3(2)	3(2)	2(2)
C(20)	34(2)	43(2)	63(3)	-9(2)	7(2)	5(2)
C(21)	32(2)	61(3)	42(2)	-10(2)	6(2)	-2(2)
C(22)	33(2)	53(2)	39(2)	1(2)	0(2)	-4(2)
C(23)	28(2)	40(2)	41(2)	-2(2)	-3(1)	2(2)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for sya03-269.

	x	y	z	U(eq)
H(4)	1361	4540	2957	39
H(5)	1492	2331	2974	41
H(6)	3845	1264	3145	41
H(7)	6096	2475	3327	38
H(9A)	5018	8699	3887	50
H(9B)	3827	8801	3297	50
H(9C)	5653	8825	3164	50
H(11)	9264	3481	3555	45
H(12)	10528	2696	4478	56
H(13)	9626	3217	5500	58
H(14)	7492	4576	5622	54
H(15)	6229	5391	4711	44
H(17A)	-681	7658	3123	43
H(17B)	218	9005	3062	43
H(19)	-1135	10529	3677	49
H(20)	-2485	11144	4610	56
H(21)	-2803	9694	5455	54
H(22)	-1805	7614	5371	50
H(23)	-454	7002	4447	44

Table 6. Torsion angles [°] for sya03-269.

O(1)-S(1)-N(1)-C(1)	101.5(3)
C(8)-S(1)-N(1)-C(1)	-25.2(3)
C(10)-S(1)-N(1)-C(1)	-138.6(3)
S(1)-N(1)-C(1)-C(2)	17.0(5)
S(1)-N(1)-C(1)-C(9)	-163.9(3)
N(1)-C(1)-C(2)-C(3)	1.2(5)
C(9)-C(1)-C(2)-C(3)	-177.7(3)
N(1)-C(1)-C(2)-C(16)	-179.6(3)
C(9)-C(1)-C(2)-C(16)	1.4(5)
C(1)-C(2)-C(3)-C(8)	-6.1(5)
C(16)-C(2)-C(3)-C(8)	174.8(3)
C(1)-C(2)-C(3)-C(4)	171.4(3)
C(16)-C(2)-C(3)-C(4)	-7.7(5)
C(8)-C(3)-C(4)-C(5)	0.7(5)
C(2)-C(3)-C(4)-C(5)	-177.0(3)
C(3)-C(4)-C(5)-C(6)	-1.3(6)
C(4)-C(5)-C(6)-C(7)	0.8(5)
C(5)-C(6)-C(7)-C(8)	0.3(5)
C(6)-C(7)-C(8)-C(3)	-1.0(5)
C(6)-C(7)-C(8)-S(1)	-177.2(3)
C(4)-C(3)-C(8)-C(7)	0.4(5)
C(2)-C(3)-C(8)-C(7)	178.1(3)
C(4)-C(3)-C(8)-S(1)	176.8(2)
C(2)-C(3)-C(8)-S(1)	-5.5(4)
O(1)-S(1)-C(8)-C(7)	64.8(3)
N(1)-S(1)-C(8)-C(7)	-164.2(3)
C(10)-S(1)-C(8)-C(7)	-51.7(3)
O(1)-S(1)-C(8)-C(3)	-111.7(3)
N(1)-S(1)-C(8)-C(3)	19.3(3)
C(10)-S(1)-C(8)-C(3)	131.8(3)
O(1)-S(1)-C(10)-C(15)	164.2(3)
N(1)-S(1)-C(10)-C(15)	37.4(3)
C(8)-S(1)-C(10)-C(15)	-77.1(3)
O(1)-S(1)-C(10)-C(11)	-16.3(4)

N(1)-S(1)-C(10)-C(11)	-143.2(3)
C(8)-S(1)-C(10)-C(11)	102.3(3)
C(15)-C(10)-C(11)-C(12)	-1.2(6)
S(1)-C(10)-C(11)-C(12)	179.4(3)
C(10)-C(11)-C(12)-C(13)	1.2(7)
C(11)-C(12)-C(13)-C(14)	-0.8(7)
C(12)-C(13)-C(14)-C(15)	0.4(7)
C(13)-C(14)-C(15)-C(10)	-0.4(6)
C(11)-C(10)-C(15)-C(14)	0.8(6)
S(1)-C(10)-C(15)-C(14)	-179.8(3)
C(17)-O(3)-C(16)-O(2)	-2.5(5)
C(17)-O(3)-C(16)-C(2)	179.4(3)
C(1)-C(2)-C(16)-O(2)	122.0(4)
C(3)-C(2)-C(16)-O(2)	-58.8(5)
C(1)-C(2)-C(16)-O(3)	-59.9(4)
C(3)-C(2)-C(16)-O(3)	119.3(3)
C(16)-O(3)-C(17)-C(18)	166.3(3)
O(3)-C(17)-C(18)-C(19)	133.9(3)
O(3)-C(17)-C(18)-C(23)	-51.4(5)
C(23)-C(18)-C(19)-C(20)	-1.1(6)
C(17)-C(18)-C(19)-C(20)	173.8(4)
C(18)-C(19)-C(20)-C(21)	0.6(6)
C(19)-C(20)-C(21)-C(22)	-0.3(6)
C(20)-C(21)-C(22)-C(23)	0.4(6)
C(21)-C(22)-C(23)-C(18)	-0.9(6)
C(19)-C(18)-C(23)-C(22)	1.2(6)
C(17)-C(18)-C(23)-C(22)	-173.6(4)

Symmetry transformations used to generate equivalent atoms:

References

- [1] a) B. Ye, N. Cramer, *J. Am. Chem. Soc.* **2013**, *135*, 636; b) G. Smits, B. Audic, M. D. Wodrich, C. Corminboeuf, N. Cramer, *Chem. Sci.* **2017**, *8*, 7174; c) Y. Sun, N. Cramer, *Chem. Sci.* **2018**, *9*, 2981.
- [2] H. Tsutsui, T. Abe, S. Nakamura, M. Anada, S. Hashimoto, *Chem. Pharm. Bull.* **2005**, *53*, 1366.
- [3] a) K. Al-Farhan, M. Ghazzali, H. M. A. Al-Hazimi, A. El-Faham, J. Reedijk, *J. Mol. Struct.* **2011**, *994*, 269; b) Z. Bourhani, A. V. Malkov, *Chem. Commun.* **2005**, 4592; c) S. Kitagaki, M. Anada, O. Kataoka, K. Matsuno, C. Umeda, N. Watanabe, S.-i. Hashimoto, *J. Am. Chem. Soc.* **1999**, *121*, 1417.
- [4] A. Tota, M. Zenzola, S. J. Chawner, S. S. John-Campbell, C. Carlucci, G. Romanazzi, L. Degennaro, J. A. Bull, R. Luisi, *Chem. Commun.* **2017**, *53*, 348.
- [5] H. Lv, J. Shi, B. Wu, Y. Guo, J. Huang, W. Yi, *Org. Biomol. Chem.* **2017**, *15*, 8054.
- [6] X. Chen, Y. Xie, X. Xiao, G. Li, Y. Deng, H. Jiang, W. Zeng, *Chem. Commun.* **2015**, *51*, 15328.
- [7] Y. Jiang, V. Z. Y. Khong, E. Lourdasamy, C. M. Park, *Chem. Commun.* **2012**, *48*, 3133.
- [8] Y.-T. Tsoi, Z. Zhou, A. S. C. Chan, W.-Y. Yu, *Org. Lett.* **2010**, *12*, 4506–4509.
- [9] Q. Liu, M. Li, R. Xiong, F. Mo, *Org. Lett.* **2017**, *19*, 6756.
- [10] S. Yoshida, H. Nakajima, K. Uchida, T. Yano, M. Kondo, T. Matsushita, T. Hosoya, *Chem. Lett.* **2017**, *46*, 77.

¹⁹F NMR

e)

19F NMR

31P NMR