GeoMed 2017, Porto

Healthy Urban Planning:

Body Mass Index association with exposure to night-time noise in the canton of Geneva







Andrea Salmi¹, Jérôme Chenal¹, Idris Guessous^{2,4}, Stéphane Joost^{1,2,3}

1 Urban and regional planning community (CEAT), School of Architecture, Civil and Environmental Engineering (ENAC), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland 2 Unit of Population Epidemiology, Division of Primary Care Medicine, Department of Community Medicine, Primary Care and Emergency Medicine, Geneva University Hospitals, Geneva, Switzerland 3 Laboratory of Geographic Information Systems (LASIG), School of Architecture, Civil and Environmental Engineering (ENAC), Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

4 Department of Ambulatory Care and Community Medicine, University of Lausanne

Health and Land-Use Planning project (HeLP):

The HeLP project purpose is to produce an improved and adapted tool for urban planning to make Switzerland healthier; promoting healthy housing, active travel, healthy environment and vibrant neighbourhoods₁. The study of the spatial dependence of health determinants, e.g. obesity, and the study of the association with patterns of urbanization, allow us to validate the approach and identify the priorities.

DATASET

Bus Santé study₂: geo-referenced participants,N > 15000 in the canton of Geneva, on cardiovascular diseases, ongoing from 1995


SonBase: night-noise geo-data throughout Switzerland, computed by road, rail and airport traffic, resolution 10 m² 5

OBESITY & NOISE IN SWITZERLAND



- 10% obese population: percentage doubled from 1992 to 2012₃
- In Europe, in 2011, 20% report disturbed sleep4
- 13% of the population is exposed to noise levels over 50 dB(A) during the night, and 34% to 45 dB(A) (WHO recommendation)₅
- CHF 1174 million: Health cost and loss of rent in 2005 given by the noise₅

DATA TREATMENT



Noise sampling from SonBase around each participant

> Variables: Mean, Median, Max,

RAW BMI

• 1 High-High (HH) 2 Low-Low (LL)

3 Low-High (LH) 4 High-Low (HL)

ADJUSTED BMI By: age, gender, median revenue, education, smoking habits

N=15544

Spatial lag: 800m

p<0.05


Permutations: 999

Adjusted by: median revenue

CLUSTERING Of BMI

Univariate Local Moran statistic with GeoDa ©: creation of LISA clusters

(Local Indicators of Spatial Association)₆



GENERAL RESULTS

- Heterogeneous distribution: 64% of participants belongs to clusters
- •10.4% are obese and 43.2% overweight
- •28% are characterised by more then 50 dB(A) and 51% more then 45 dB(A)
- The Low-Low cluster is located in the town center despite different adjustments and is characterised by high road traffic
- The High-High clusters are located in the poorer districts and between train station and airport

Red clusters show locations where a high BMI is significantly associated with a high average noise level (HH), and blue clusters show locations where a low BMI is significantly associated with a low average noise level (LL). The pink and purple classes show high-low and low-high relationships respectively. White dots show sampling areas where BMI are randomly distributed and are not significant in term of spatial dependence.

STATISTICAL **ANALYSIS**

ANOVA (analysis of variance) and multiple T-test of noise variables between clusters p<0.05

Clusters are significantly different. Low-Low cluster are exposed to higher noise level compared to the High-High (e.g. 50.7 vs 45.6 dB(A) of mean noise level with adjusted BMI by median revenue). The resulting association is inverse.

Observations:

- Geneva is listed as one of the city with the best quality of life in the world (8th₇): no health impact of noise pollution due to the presence of other positive health determinants located in the town center
- Annoyance from noise may be caused by high variability when mean is under the thresholds

REFERENCES

- 1. Healthy Urban Development Unit (HUDU), 2015. Healthy Urban Planning Checklist. National Health Service (NHS).
- 2. Guessous, I., et al., (2014). A comparison of the spatial dependence of body mass index among adults and children in a Swiss general population. Nutritional Diabetes, 4, e111.
- 3. Office Fédéral de la Statistique (OFS). (2014). Enquête suisse la sante 2012 : Surpoids et Obésité. 4. EEA 2007: Presenting Noise Mapping Information to the Public. A Position Paper from the European Environment Agency Working Group on the Assessment of Exposure to Noise (WG-AEN).
- 5. Federal Office for the Environment (FOEN), 2014. Noise pollution in Switzerland. SonBase. 6. Anselin, L., (1995). Local indicators of spatial association-Lisa. Geographic Analysis 27, 93-115.
- 7. Mercer, (2017). Quality of Living Rankings.
- 8. Sim, C.S et al,(2015). The Effects of Different Noise Types on Heart Rate Variability in Men. Yonsei Med journal, 56 (1), 235-243.