

Exploitation of Unconventional Electrophiles in Enantioselective Pd(0)-Catalyzed C–H Functionalizations

Daria Grosheva[§] and Nicolai Cramer*

[§]SCS-Metrohm Award for best oral presentation in Organic Chemistry

Abstract: Asymmetric metal-catalyzed C–H functionalization is a powerful strategy for the rapid generation of complex target molecules from simple precursors. While aryl (pseudo)halides have been used as electrophilic substrates in enantioselective Pd(0)-catalyzed syntheses of *N*-heterocycles, alternative partners for C–H functionalizations remain underexplored. Herein we describe the first application of ketene aminal phosphates as competent electrophiles in a Pd(0)-catalyzed desymmetrization. This transformation allowed the efficient synthesis of chiral isoindolines in a highly enantioselective fashion. Furthermore, intending to introduce sought-after perfluoroalkyl substituents, we explored intramolecular C–H functionalization of imidoyl chlorides resulting in the asymmetric preparation of previously inaccessible 1*H*-isoindoles. Tailored chiral monodentate ligands afforded high levels of enantioinduction in both transformations.

Keywords: Asymmetric catalysis · C–H functionalization · Heterocycles · Ligand design · Palladium




Daria Grosheva was born and raised in Saint Petersburg, Russia. While completing her Specialist of Chemistry degree at St Petersburg University she conducted a summer internship at the University of Texas at San Antonio with Prof. Oleg Larionov and a research stay at Ecole Polytechnique, Paris, with Dr. Yvan Six, supported by a Scholarship of the President of Russian Federation. Daria commenced her doctoral

studies at the group of Prof. Cramer in late 2014, working on the development of asymmetric Pd-catalyzed C–H functionalizations. In mid-2019 she will join the Hyster group at Princeton University as an SNF Postdoctoral Fellow.

Introduction


Over the past decade, metal-catalyzed C–H functionalization has become a key research area in synthetic organic chemistry.^[1] In particular, the stereocontrolled functionalization of C–H bonds has emerged as a powerful tool for the rapid construction of elaborate structures from simple and readily available starting materials.^[2] For example, the enantioselective synthesis of *N*-heterocycles by Pd(0)-catalyzed direct arylation has been extensively explored, typically employing aryl (pseudo)halides as electrophilic substrates (Scheme 1).^[3] Related asymmetric C–H alkenylations,^[4] alkylations^[5] and imidoylations^[6] remain far less developed, resulting in the confined scope of the obtainable products. We aimed to explore unconventional electrophilic partners in order to access a broader diversity of elaborate *N*-heterocyclic scaffolds.

C–H Functionalization of Aryl (Pseudo)halides



* many precedents
* limited scope of available products

C–H Functionalization of Complex *N*-Containing Electrophiles



* underexplored
* more elaborate products are accessible

Scheme 1. Aryl (pseudo)halides vs complex *N*-containing electrophiles for asymmetric Pd(0)-catalyzed C–H functionalizations.

Results and Discussion

Synthesis of Chiral Electron-Rich Monodentate Phosphine Ligands^[7]


Efforts towards successful development of enantioselective metal-catalyzed transformations, including Pd(0)-catalyzed C–H functionalizations, go hand-in-hand with the design of suitable chiral ligands. Mechanistically, while the oxidative addition of electrophiles to Pd(0) complexes at ambient temperature is accelerated by electron-rich phosphines, the concerted metalation-deprotonation (CMD) step requires a monoligated Pd(0) species (Scheme 2). Overall, a successful catalytic asymmetric process is, therefore, favored by a ligand combining both properties; however, only a limited range of efficient chiral electron-rich monodentate phosphines is currently available.

Available chiral electron-rich monodentate phosphines include the SagePhos ligand family, previously developed in the Cramer group; it combines a 2',6'-dialkoxybiaryl core with the *C*₂-symmetric electron-rich phospholane moiety (Scheme 3).^[3d] In

*Correspondence: Prof. Dr. N. Cramer

E-mail: nicolai.cramer@epfl.ch

Ecole Polytechnique Fédérale de Lausanne, EPFL SB ISIC LCSA, BCH 4305, CH-1015 Lausanne



Scheme 2. Generic catalytic cycle for Pd(0)-catalyzed C–H functionalization.


order to have an extra handle for further reactivity and selectivity tuning, we sought to introduce an additional modification position in the SagePhos scaffold; namely, the powerful atrop-chiral biaryl unit of the MOP ligand.^[8] The synthesis of methoxy-substituted ligand **L1** has been reported;^[9] however, the previous approach is low yielding and rather inconvenient, and while it was suitable for the preparation of a single member of the ligand family, it did not fit our purpose of generating a small library of BINOL-derived phosphines. To this end, our strategy was based on the Pd(0)-catalyzed coupling of BINOL bis-triflate (**R-1** or **S-1**) with secondary phosphine oxide **2**, which resulted in the selective formation of products **3** or **4**. Saponification of the triflyl moiety revealed the hydroxyl group, which was alkylated or silylated with a substituent of variable steric demand. Finally, reduction of phosphine oxide moiety with Cl_3SiH gave access to bulky monodentate ligands **L1–L6**, as well as their diastereoisomeric

counterparts **L7** and **L8**. The overall route proved to be scalable, reproducible and divergent with respect to the introduction of the substituent R on the lower aryl ring of the ligand scaffold.


Asymmetric C–H Functionalization of Ketene Aminal Phosphates^[7]

Regarding the reaction development, we first envisioned a methodology based on the Pd(0)-catalyzed desymmetrization of ketene aminal electrophiles delivering fused *N*-heterocycles (Scheme 4). We chose ketene aminal phosphates as substrates as they can be readily accessed by deprotonation of the corresponding imides and trapping with a phosphoryl chloride.^[10] Moreover, in stark contrast to ketene aminal triflates, the corresponding phosphates are stable compounds that could be isolated and characterized.^[10,11] We were also interested whether the built-in phosphate moiety could serve as a competent base in the concerted metalation-deprotonation step and what influence it would have on the enantioselectivity of the process. Finally, the successful alkenylation followed by functional group manipulations would afford enantioselective access *via* a new disconnection to the valuable indolizidine scaffold that is present in many natural products.


Phenyl-substituted ketene aminal phosphate **5a** was chosen as the prototype substrate and the system including Pd(dba)₂, 2,2-diphenylpropionic acid and Cs_2CO_3 was identified as the optimal set of reaction conditions after the preliminary optimization studies. With newly developed phospholanes **L1–L6** a general trend was observed: introducing a bulkier substituent R on the ligand backbone resulted in higher enantioselectivities, accompanied by a loss in reactivity; and TBDPS-substituted ligand **L6** was the most selective (Table 1, entries 1–6). When axially diastereoisomeric phosphines **L7** and **L8** were employed, the enantioselectivities slightly dropped, suggesting that the axial chirality of the ligand scaffold has a minor influence on the selectivity, and the dominant portion of enantioinduction primarily arises from the dimethyl phospholane unit (entries 7 and 8). This hypothesis was further

Scheme 3. Design and synthesis of chiral electron-rich monodentate ligands **L1–L8**.

Scheme 4. Enantioselective C–H functionalization of ketene aminal phosphates.



Selected examples of indolizidine alkaloids





proved when the reaction with axially chiral Cy-MOP resulted in a virtually racemic formation of the desired product (entry 9), in striking contrast to the highly enantioselective transformation enabled by SagePhos-type ligand **L10** (entry 10). Notably, more electron-poor TADDOL-derived phosphoramidites **L11** and **L12** that have provided for a number of asymmetric Pd(0)-catalyzed C–H functionalizations^[12] required higher reaction temperature and displayed only modest enantiocontrol (entries 11 and 12). The selectivity of the transformation could be improved by using benzoic or pivalic acids (entries 13 and 14). Carrying out a reaction under carboxylate-free conditions (carboxylate-free Pd source and no external carboxylic acid) we tested the hypothesis on the phosphate anion acting as a base in the CMD step. Conversion and enantioselectivity were inferior compared to the original conditions, suggesting the diphenyl phosphate was a poor promoter of the concerted metalation-deprotonation process (entry 15). Replacing Pd(dba)₂ with Pd(OAc)₂ resulted in a faster reaction, albeit with lower enantioselectivity: the presence of acetate most likely influenced the selectivity of the enantiodetermining CMD step (entry 16). Employing commercially available Pd(OPiv)₂ circumvented this issue and allowed for an efficient and highly enantioselective transformation at ambient temperature with the catalyst loading reduced to 5 mol% of palladium precursor (entry 17). The nature of the employed phosphate ester was crucial for the reactivity: thus, diethyl phosphate **5a'** remained inert, even at an elevated temperature of 80 °C (entry 18). Finally, the transformation could be carried out on a larger scale, delivering 1.05 g of isoindoline **6a** without affecting the enantioselectivity (entry 19).

With the optimal conditions in hand, we evaluated the scope and limitations of the transformation (Scheme 5). Electron-poor and electron-rich substrates could be functionalized delivering corresponding heterocycles **6b** and **6c** in comparable yields and similar enantioselectivities. Furthermore, an aryl chloride moiety in substrate **5d** remained intact during the Pd-catalyzed transformation, allowing for a potential orthogonal late-stage functionalization. The cyclizations of *meta*-substituted substrates **5e** and **5f** proceeded with a remarkable level of regioselectivity, and the more accessible *ortho*-C–H groups were functionalized exclusively. The substrate featuring electron-rich thiophene was well tolerated, and the corresponding heterocycle **6g** was furnished in excellent enantioselectivity. *ortho*-Substituted substrates (e.g. **5h**) could also be functionalized, however, in this case, higher reaction temperature was required, resulting in reduced enantioselectivity of the obtained isoindoline. Spirocyclic ketene aminal phosphate **5i** was efficiently functionalized at ambient temperature, but with somewhat reduced selectivity.


The alkenyl moiety in isoindoline **6a** could be selectively hydrogenated from the more accessible site, delivering a single diastereoisomer of lactam **7a** (Scheme 6). Subsequent nickel-

Table 1. Optimization of the enantioselective isoindoline synthesis.




Entry	[Pd]	L*	Acid	Yield of 6a ^a	er
1	Pd(dba) ₂	L1	Ph ₂ MeCCO ₂ H	78%	78.5:21.5
2	Pd(dba) ₂	L2	Ph ₂ MeCCO ₂ H	65%	87:13
3	Pd(dba) ₂	L3	Ph ₂ MeCCO ₂ H	48%	87:13
4	Pd(dba) ₂	L4	Ph ₂ MeCCO ₂ H	32%	90:10
5	Pd(dba) ₂	L5	Ph ₂ MeCCO ₂ H	16%	77.5:22.5
6	Pd(dba) ₂	L6	Ph ₂ MeCCO ₂ H	29%	90.5:9.5
7	Pd(dba) ₂	L7	Ph ₂ MeCCO ₂ H	43%	70:30
8	Pd(dba) ₂	L8	Ph ₂ MeCCO ₂ H	38%	86.5:13.5
9	Pd(dba) ₂	L9	Ph ₂ MeCCO ₂ H	19%	51:49
10	Pd(dba) ₂	L10	Ph ₂ MeCCO ₂ H	82%	89:11
11 ^b	Pd(dba) ₂	L11	Ph ₂ MeCCO ₂ H	12%	70.5:29.5
12 ^b	Pd(dba) ₂	L12	Ph ₂ MeCCO ₂ H	60%	79.5:20.5
13	Pd(dba) ₂	L6	PhCO ₂ H	37%	93:7
14	Pd(dba) ₂	L6	PivOH	36%	93.5:6.5
15	Pd(dba) ₂	L6	–	21%	73:27
16	Pd(OAc) ₂	L6	PivOH	82%	75.5:24.5
17 ^c	Pd(OPiv) ₂	L6	PivOH	93%	94.5:5.5
18 ^{c,d}	Pd(OPiv) ₂	L6	PivOH	8%	not determined
19 ^e	Pd(OPiv) ₂	L6	PivOH	80%	95.5:4.5

^aDetermined by ¹H NMR with an internal standard. ^bAt 80 °C. ^cWith 5 mol% Pd(OPiv)₂, 10 mol% **L6** at 0.5 M and 23 °C. ^d**5a'** instead of **5a** at 80 °C. ^e5 mmol **5a** [2.56 g], 5 mol% Pd(OPiv)₂, 10 mol% **L6**, 7.5 mmol Cs₂CO₃, 0.5 M in PhMe, at 23 °C for 48 h.



Scheme 5. Selected scope entries of the asymmetric ketene aminal phosphate cyclization. ^[a]With 10 mol% Pd(OPiv)₂, 20 mol% **L6**. ^[b]At 60 °C.



Scheme 6. Two-step preparation of indolizidine **8a**.

catalyzed amide reduction with phenylsilane^[13] afforded tertiary amine **8a** featuring the sought-after indolizidine scaffold.

Enantioselective Synthesis of Perfluoroalkyl-Substituted 1*H*-Isoindoles^[14]




Having developed an enantioselective approach toward isoindolines, we aspired to employ Pd(0)-catalyzed C–H functionalization to access other types of *N*-heterocycles, namely, 1*H*-isoindoles. The isoindole framework is found in natural products, pharmaceuticals and materials (Scheme 7).^[15] However, synthetic approaches to 1*H*-isoindoles remain largely limited to conventional methods.^[16] As active molecules often contain stereogenic centers, catalytic asymmetric strategy to access 1*H*-isoindoles would be highly desirable. We proposed an approach based on the intramolecular enantioselective C–H functionalization of perfluoroalkyl-substituted imidoyl chlorides, electrophilic partners, which were introduced by Uneyama,^[17] and subsequently probed in an asymmetric functionalization by our group.^[6b] If successful, our strategy would allow for an introduction of the quaternary stereogenic center, furnishing an attractive three-dimensional framework with functional exit vectors. Employing perfluoroalkylated imidoyl chlorides as substrates would result in the introduction of a perfluoroalkyl substituent in the product. Given the interest in fluorine-containing compounds in active-ingredient discovery,^[18] the construction of such patterns would be an additional advantage of the approach.

Furthermore, while several methods enabling synthesis of perfluoroalkylated isoindolines have been reported,^[19] the corresponding 1*H*-isoindoles bearing this appealing substitution pattern remain inaccessible through conventional methods.^[16a]

The initial ligand screening was carried out with isopropyl-substituted imidoyl chloride **9a** (Scheme 8). The standard TADDOL-derived ligands **L11–L13** provided the desired product in excellent yields, albeit, with only modest enantioselectivities. When we switched to the use of bulky diazaphospholane ligands **L14–L16** developed in the Cramer group previously,^[20] the reaction was operating already at 40 °C. Methoxy-substituted diazaphospholane **L16** bearing 1-naphthyl and *tert*-butyl groups on the chiral flanking arms was the ligand of choice and resulted in a highly enantioselective transformation. The catalyst loading could be lowered to 5 mol% of Pd(dba)₂ with the reaction time prolonged to 36 hours.

In regard to the scope of the transformation, a wide variety of substrates were well-tolerated (Scheme 9). Again, electron-rich and electron-poor substrates reacted well, however, the functionalization of *para*-methoxy substituted imidoyl chloride **9c** required longer reaction time, which is consistent with a CMD-type activation mechanism.^[21] Selective functionalization of 2-naphthyl-substituted trifluoroacetimidoyl chloride delivered **10d** featuring the extended aromatic system of 1*H*-benzo[*f*]isoindole. Substrate **9e**, bearing electron-rich thienyl substitu-

Biologically active molecules with an isoindole motif


Scheme 7. Enantioselective C–H functionalization approach toward chiral 1*H*-isoindoles.

Scheme 8. Optimization of the enantioselective C–H functionalization of trifluoroacetimidoyl chloride.


ents, was efficiently functionalized, and the corresponding thienopyrrole **10e** was obtained in excellent enantioselectivity. Moreover, X-ray analysis of **10e** enabled the assignment of the absolute configuration. Imidoyl chlorides homologated with longer perfluoroalkylated chains reacted equally well, and the corresponding thienopyrroles **10f** and **10g** were obtained with

comparable levels of yield and enantioselectivity. Various alkyl groups on the quaternary stereocenter were well-tolerated. For example, cyclopropyl-substituted imidoyl chloride **9h** was cyclized with excellent chemoselectivity, and no product of competing C(sp³)–H functionalization was observed. Functionalized substituents could also be introduced: azetidynyl-substituted

Scheme 9. Selected scope entries of the asymmetric 1*H*-isoindole synthesis. ^[a]Reaction time: 80 h. ^[b]10 mol% Pd(dba)₂, 20 mol% L16.



Scheme 10. Scope of the asymmetric synthesis of 1*H*-isoindole with small substituents R.



isoindole **10i** was obtained in high yield and good enantioselectivity. However, when substrates that bear less bulky substituents, such as **9j**, were tested in the reaction, a significant decrease in enantioinduction was observed.

Identifying the set of reaction conditions suitable for the preparation of 1*H*-isoindoles with small substituent R required considerable additional optimization efforts. Essentially, the combination of TADDOL-phosphoramidite ligand **L11** with the cesium salt of *ortho*-anisic acid enabled highly enantioselective cyclization of the corresponding substrates (Scheme 10). As before, the reaction rate and the enantioselectivity were largely independent of the electronic characteristics of the functionalized arenes. Electron-neutral, electron-rich and electron-poor substrates all reacted well, furnishing the corresponding isoindoles **10j**, **10k** and **10l**, as well as benzothiophene substituted product **10m**. Furthermore, isoindole **10n** featuring a synthetically valuable benzl ether chain was accessed in excellent yield and high enantioselectivity.

Summary

In conclusion, we developed a short and divergent route toward modular and tunable chiral phosphines filling the gap of accessible electron-rich monodentate ligands. These ligands were vital for the successful enantioselective C–H alkenylations at ambient temperature, where the potential of ketene aminal phosphates as electrophilic partners in Pd(0)-catalyzed transformations was showcased. Furthermore, we discovered that employing imidoyl chlorides as substrates in C(sp²)–H functionalization resulted in the formation of highly functionalized perfluoroalkylated 1*H*-isoindoles, which were previously inaccessible. Various substitution patterns were well tolerated, and high levels of stereocontrol were enabled by virtue of using a bulky phosphordiamidite ligand.

Acknowledgements

Daria Grosheva is grateful to Metrohm and the Swiss Chemical Society for being granted with the SCS-Metrohm Best Oral Presentation Award. We thank the Swiss National Science Foundation (no. 155967) for financial support.

Received: January 30, 2019

- [1] a) J. Yamaguchi, A. D. Yamaguchi, K. Itami, *Angew. Chem. Int. Ed.* **2012**, *51*, 8960; b) J. He, M. Wasa, K. S. L. Chan, Q. Shao, J.-Q. Yu, *Chem. Rev.* **2017**, *117*, 8754; c) J. R. Hummel, J. A. Boerth, J. A. Ellman, *Chem. Rev.* **2017**, *117*, 9163; d) Z. Dong, Z. Ren, S. J. Thompson, Y. Xu, G. Dong, **2017**, *117*, 9333; e) K. Murakami, S. Yamada, T. Kaneda, K. Itami, *Chem. Rev.* **2017**, *117*, 9302.
- [2] a) C. G. Newton, S.-G. Wang, C. C. Oliveira, N. Cramer, *Chem. Rev.* **2017**, *117*, 8908; b) T. G. Saint-Denis, R.-Y. Zhu, G. Chen, Q.-F. Wu, J.-Q. Yu, *Science* **2018**, *359*, eaao4798.
- [3] a) S. Anas, A. Cordi, H. B. Kagan, *Chem. Commun.* **2011**, *47*, 11483; b) M. Nakanishi, D. Katayev, C. Besnard, E. P. Kündig, *Angew. Chem. Int. Ed.* **2011**, *50*, 7438; c) D. Katayev, M. Nakanishi, T. Burgi, E. P. Kündig, *Chem. Sci.* **2012**, *3*, 1422; d) T. Saget, S. J. Lemouzy, N. Cramer, *Angew. Chem. Int. Ed.* **2012**, *51*, 2238; e) T. Saget, N. Cramer, *Angew. Chem. Int. Ed.* **2012**, *51*, 12842; f) E. Larionov, M. Nakanishi, D. Katayev, C. Besnard, E. P. Kündig, *Chem. Sci.* **2013**, *4*, 1995; g) T. Saget, N. Cramer, *Angew. Chem. Int. Ed.* **2013**, *52*, 7865; h) D. Katayev, E. Larionov, M. Nakanishi, C. Besnard, E. P. Kündig, *Chem. Eur. J.* **2014**, *20*, 15021; i) Z.-Q. Lin, W.-Z. Wang, S.-B. Yan, W.-L. Duan, *Angew. Chem. Int. Ed.* **2015**, *54*, 6265; j) J. Pedroni, T. Saget, P. A. Donets, N. Cramer, *Chem. Sci.* **2015**, *6*, 5164; k) C. He, M. Hou, Z. Zhu, Z. Gu, *ACS Catal.* **2017**, *7*, 5316; l) L. Yang, R. Melot, M. Neuburger, O. Baudoin, *Chem. Sci.* **2017**, *8*, 1344; m) L. Yang, M. Neuburger, O. Baudoin, *Angew. Chem. Int. Ed.* **2018**, *57*, 1394; n) R. Rocaboy, D. Dailler, O. Baudoin, *Org. Lett.* **2018**, *20*, 772; o) C. G. Newton, E. Braconi, J. Kuziola, M. D. Wodrich, N. Cramer, *Angew. Chem. Int. Ed.* **2018**, *57*, 11040.
- [4] a) M. R. Albicker, N. Cramer, *Angew. Chem. Int. Ed.* **2009**, *48*, 9139; b) P. M. Holstein, D. Dailler, J. Vantourout, J. Shaya, A. Millet, O. Baudoin, *Angew. Chem. Int. Ed.* **2016**, *55*, 2805.
- [5] a) J. Pedroni, M. Boghi, T. Saget, N. Cramer, *Angew. Chem. Int. Ed.* **2014**, *53*, 9064; b) J. Pedroni, N. Cramer, *Angew. Chem. Int. Ed.* **2015**, 11826.
- [6] a) J. Wang, D.-W. Gao, J. Huang, S. Tang, Z. Xiong, H. Hu, S.-L. You, Q. Zhu, *ACS Catal.* **2017**, *7*, 3832; b) J. Pedroni, N. Cramer, *J. Am. Chem. Soc.* **2017**, *139*, 12398; c) S. Luo, Z. Xiong, Y. Lu, Q. Zhu, *Org. Lett.* **2018**, *20*, 1837.
- [7] D. Grosheva, N. Cramer, *ACS Catal.* **2017**, *7*, 7417.
- [8] T. Hayashi, *Acc. Chem. Res.* **2000**, *33*, 354.
- [9] B. Saha, T. V. RajanBabu, *J. Org. Chem.* **2007**, *72*, 2357.
- [10] K. C. Nicolaou, K. Namoto, *Chem. Commun.* **1998**, 1757.
- [11] a) J. W. Coe, *Org. Lett.* **2000**, *2*, 4205; b) F. L. Galbo, E. G. Occhiato, A. Guarna, C. Faggi, *J. Org. Chem.* **2003**, *68*, 6360.
- [12] J. Pedroni, N. Cramer, *Chem. Commun.* **2015**, *51*, 17647.
- [13] B. J. Simmons, M. Hoffmann, J. Hwang, M. K. Jackl, N. K. Garg, *Org. Lett.* **2017**, *19*, 1910.
- [14] D. Grosheva, N. Cramer, *Angew. Chem. Int. Ed.* **2018**, *57*, 13644.
- [15] a) K. Speck, T. Magauer, *Beilstein J. Org. Chem.* **2013**, *9*, 2048; b) B.-M. Swahn, K. Kolmodin, S. Karlström, S. von Berg, P. Söderman, J. Holenz, S. Berg, J. Lindström, M. Sundström, D. Turek, J. Kihlström, C. Slivo, L. Andersson, D. Pyring, D. Rotticci, L. Öhberg, A. Kers, K. Bogar, F. von Kieseritzky, M. Bergh, L.-L. Olsson, J. Janson, S. Eketjäll, B. Georgievskaa, F. Jeppsson, J. Fälting, *J. Med. Chem.* **2012**, *55*, 9346; c) G. E. Morse, T. P. Bender, *ACS Appl. Mater. Interfaces* **2012**, *4*, 5055.
- [16] a) C. G. Savarin, C. Grisé, J. A. Murry, R. A. Reamer, D. L. Hughes, *Org. Lett.* **2007**, *9*, 981; b) B. Hatano, K. Miyoshi, H. Sato, T. Ito, T. Ogata, T. Kijima, *Tetrahedron Lett.* **2010**, *51*, 5399; c) M. Kuroda, K. Kobayashi, *Helv. Chim. Acta* **2015**, *98*, 279.
- [17] a) K. Tamura, H. Mizukami, K. Maeda, H. Watanabe, K. Uneyama, *J. Org. Chem.* **1993**, *58*, 32; b) K. Uneyama, *J. Fluorine Chem.* **1999**, *97*, 11.
- [18] a) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, *37*, 320; b) T. Liang, C. N. Neumann, T. Ritter, *Angew. Chem. Int. Ed.* **2013**, *52*, 8214.
- [19] a) S. Fustero, J. Moscardó, M. Sánchez-Roselló, E. Rodríguez, P. Barrio, *Org. Lett.* **2010**, *12*, 5494; b) S. Fustero, I. Ibáñez, P. Barrio, M. A. Maestro, S. Catalán, *Org. Lett.* **2013**, *15*, 832; c) S. Fustero, L. Herrera, R. Lázaro, E. Rodríguez, M. A. Maestro, N. Mateu, P. Barrio, *Chem. Eur. J.* **2013**, *19*, 11776.
- [20] P. A. Donets, N. Cramer, *J. Am. Chem. Soc.* **2013**, *135*, 11772.
- [21] D. García-Cuadrado, P. de Mendoza, A. A. C. Braga, F. Maseras, A. M. Echavarren, *J. Am. Chem. Soc.* **2007**, *129*, 6880.