Rh^I, Ir^{III} and Co^{III} Complexes with Atropchiral Biaryl Cyclopentadienyl Ligands: Syntheses, Structures and Catalytic Activities

Coralie Duchemin,^a Gints Smits,^b and Nicolai Cramer^{*a}

^a Laboratory of Asymmetric Catalysis and Synthesis, Institute of Chemical Sciences and Engineering, Ecole Polytechnique Fédérale de Lausanne (EPFL), SB-ISIC, BCH4305, 1015 Lausanne, Switzerland

^b Latvian Institute of Organic Synthesis, Aizkraukles 21, Riga LV-1006, Latvia

*e-mail: nicolai.cramer@epfl.ch

Supplementary Information

Table of Content:

<i>General Methods</i>	S2
<i>Ligands synthesis</i>	S4
<i>Complexes synthesis</i>	S22
<i>Benchmark Rh^{III}-catalyzed reactions</i>	S27
<i>Crystallographic data</i>	S34
<i>NMR-Spectra</i>	S35
<i>References</i>	S63

General Methods

Experimental Procedures, Reagents and Glassware

All reactions were carried out under an atmosphere of nitrogen in oven-dried glassware with magnetic stirring, unless otherwise indicated. Toluene, dichloromethane, tetrahydrofuran, acetonitrile and diethyl ether were purified by an Innovative Technology Solvent Delivery System. Chemicals were used as obtained from the suppliers unless otherwise stated.

Chromatography

Flash chromatography was performed with Silicycle silica gel 60 (0.040-0.063 μm grade). Analytical thin-layer chromatography was performed with commercial glass plates coated with 0.25 mm silica gel (E. Merck, Kieselgel 60 F254). Compounds were either visualised under UV-light at 254 nm or by dipping the plates in an aqueous potassium permanganate solution followed by heating.

NMR Spectroscopy

Proton nuclear magnetic resonance (^1H NMR) data were acquired at 400 MHz on a Bruker AVANCEIII-400 spectrometer or at 600 MHz on a Bruker DRX-600. Chemical shifts (δ) are reported in parts per million (ppm) relative to incompletely deuterated CDCl_3 (s, 7.26 ppm), CD_2Cl_2 (t, 5.32 ppm, $J = 1.1\text{Hz}$) or C_6D_6 (s, 7.16 ppm). Proton decoupled Carbon-13 nuclear magnetic resonance (^{13}C $\{^1\text{H}\}$ NMR) data were acquired at 101 MHz on a Bruker AVANCEIII-400 spectrometer or at 151 MHz on a Bruker DRX-600 spectrometer. Chemical shifts are reported in ppm relative to residual CDCl_3 (77.16 ppm), CD_2Cl_2 (53.84) or C_6D_6 (128.06). The assignment of proton and carbon signals was assisted by COSY, HSQC, HMBC and DEPT-135 experiments where necessary. Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; p, pentet; hept, heptet; dd, doublet of doublets; dt, doublet of triplets; ddd, doublet of doublets of doublets; tt, triplet of triplets; tq, triplet of quartets; qt, quartet of triplets; m, multiplet. All NMR data were recorded at 298 K, unless otherwise indicated.

Infrared Spectroscopy

Infrared (IR) data were recorded on an Alpha-P Bruker FT-IR Spectrometer. Absorbance frequencies are reported in reciprocal centimeters (cm^{-1}).

Mass Spectrometry

HRMS measurements were performed on an Agilent LC-MS TOF (Multimode: ESI + APCI) or Waters Xevo G2-S QTOF (APPI or APCI). High resolution mass are given in m/z .

Melting Points

Melting points were measured on a Büchi B-540 and are uncorrected.

Enantiomeric excesses

Enantiomeric excesses were measured on an Agilent or Waters HPLC, or on a Thar SFC Investigator system using chiral stationary phase columns. Optical rotations were measured on a Polartronic M polarimeter using a 0.5 cm cell with a Na 589 nm filter.

X-Ray analyses

X-ray analysis of compounds **Rh1-Rh4**, **Rh6**, **Co1**, **Co2**, **Ir1** and **Ir2** was performed by Dr. R. Scopelliti and Dr. F. Fadaei Tirani at the EPF Lausanne.

Ligands synthesis

Compounds (*R*)-**1**^[1] and (*R*)-**4**^[2] were prepared following the literature procedure.

Synthesis of compound (*S*)-**3a**:

Scheme S1: Synthesis of (S)-3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-dicarboxylic acid

(S)-3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-dicarboxylic acid:^[3] To a solution of freshly distilled over CaH₂ 2,2,6,6-tetramethylpiperidine (6 ml, 35 mmol) in dry THF (20 ml) at 0°C under nitrogen, *n*-BuLi (1.6M in Hexanes, 14 ml, 35 mmol) was slowly added. The reaction was stirred at this temperature for 15min. The reaction was cooled down to -78 °C and freshly distilled over CaH₂ TMSCl (7.5 ml, 58 mmol) was added dropwise and stirred at -78°C for 15min. A solution of [1,1'-binaphthalene]-2,2'-dicarboxylic acid (2 g, 5.8 mmol) in dry THF (9 ml) was added at -78°C and the reaction was allowed to warm up to room temperature overnight. The reaction was quenched with a 1N HCl solution and extracted with Et₂O. The organic fraction was washed with a 1N NaOH solution and the aqueous resulting part was acidified with a 2N HCl solution to pH = 1, then extracted with Et₂O. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum to give 3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-dicarboxylic acid (2.6 g, 5.3 mmol, 91 % yield) as an off-white solid.

¹H NMR (400 MHz, CDCl₃) δ 8.22 (s, 2H), 7.91 (d, *J* = 8.2 Hz, 2H), 7.51 (t, *J* = 7.5 Hz, 2H), 7.31 (t, *J* = 7.8 Hz, 2H), 7.01 (d, *J* = 8.5 Hz, 2H), 0.41 (s, 18H); **¹³C NMR** (101 MHz, CDCl₃) δ 173.9, 137.0, 135.6, 134.4, 133.3, 132.9, 132.4, 128.3, 128.1, 127.7, 126.7, 77.2, 0.1; **IR (ATR):** ν_{max} = 2953, 1686, 1250, 840, 748 cm⁻¹; **HRMS (ESI/QTOF) *m/z*:** [M-H]⁻ Calcd for [C₂₈H₂₉O₄Si₂]⁻ 485.1610; Found 485.1616; **[α]_D²⁰** = -183.3 (*c* = 1.0, CHCl₃); **R_f**: 0.45 (Pentane:EtOAc:AcOH, 7:3:1%); **Mp** = 188 °C.

Scheme S2: Synthesis of (S)-3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-diyl dimethanol

(S)-3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-diyl dimethanol.^[3] A solution of 3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-dicarboxylic acid (1.0 g, 2.1 mmol) in dry THF (10.0 ml) under N₂ was cooled down to 0 °C. BH₃.THF complex (9.3 ml, 9.3 mmol, 1M in THF) was slowly added and then the reaction was heated to 70°C and stirred for 15 h. The mixture was cooled down to 0 °C and carefully quenched with water before being diluted with EtOAc. The organic layer was washed with a 2N HCl solution and the resulting aqueous layer was extracted with EtOAc. The combined organic layers were washed with a saturated solution of NaHCO₃, dried over Na₂SO₄ and concentrated under vacuum. The crude product was purified by column chromatography (SiO₂, Pentane:EtOAc 20:1) affording (3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-diyl dimethanol (0.9 g, 2.0 mmol, 95 % yield) as a white foam.

¹H NMR (400 MHz, CDCl₃) δ 8.22 (s, 2H), 7.92 (d, *J* = 8.1 Hz, 2H), 7.45 (ddd, *J* = 8.1, 6.8, 1.1 Hz, 2H), 7.23 (ddd, *J* = 8.2, 6.8, 1.2 Hz, 2H), 6.94 (d, *J* = 8.4 Hz, 2H), 4.57 (d, *J* = 11.6 Hz, 2H), 4.20 (d, *J* = 11.6 Hz, 2H), 2.79 (s, 2H), 0.49 (s, 18H); **¹³C NMR** (101 MHz, CDCl₃) δ 141.2, 138.8, 136.6, 135.9, 133.6, 132.5, 128.4, 127.2, 126.3, 126.2, 62.5, 1.1; **IR (ATR):** ν_{max} = 3550-3092 (br), 2952, 1249, 1026, 837, 753, 732 cm⁻¹; **HRMS (APCI/QTOF) *m/z*:** [M+Na]⁺ Calcd for [C₂₈H₃₄NaO₂Si₂]⁺ 481.1990; Found 481.1986; **[α]_D²⁰** = -107.5 (*c* = 1.0, CHCl₃); **R_f**: 0.58 (Pentane: EtOAc, 9:1); **Mp** = 130 °C;

Scheme S3: Synthesis of (S)-2a

(S)-(2,2'-bis(bromomethyl)-[1,1'-binaphthalene]-3,3'-diyl)bis(trimethylsilane) (R)-2a:^[3] A solution of PBr₃ (0.16 ml, 1.66 mmol) was added dropwise at 0 °C under N₂ to a solution (3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-diyl)dimethanol (0.76mg, 1.66 mmol) in THF (17 ml). The reaction was warmed up to room temperature and stirred for 3 h. The mixture was cooled down to 0 °C and carefully quenched with a saturated solution of NaHCO₃ before being diluted with EtOAc. The mixture was extracted with EtOAc, dried over Na₂SO₄ and concentrated under vacuum. The crude product was purified by column chromatography (SiO₂, Pentane/Ethyl Acetate, 20:1) to give (2,2'-bis(bromomethyl)-[1,1'-binaphthalene]-3,3'-diyl)bis(trimethylsilane) (R)-2a (0,9 g. 1.54 mmol, 93 % yield) as a white solid.

¹H NMR (400 MHz, CDCl₃) δ 8.24 (s, 2H), 7.91 (d, *J* = 8.1 Hz, 2H), 7.42 (t, *J* = 6.8 Hz, 2H), 7.23 (t, *J* = 8.1 Hz, 2H), 7.01 (d, *J* = 8.4 Hz, 2H), 4.44 (d, *J* = 10.3 Hz, 2H), 4.34 (d, *J* = 10.3 Hz, 2H), 0.55 (s, 18H); ¹³C NMR (101 MHz, CDCl₃) δ 138.4, 137.8, 137.6, 136.3, 133.0, 132.7, 128.1, 127.5, 126.9, 34.8, 1.4; IR (ATR): ν_{max} = 3050, 2953, 1251, 839, 750, 733 cm⁻¹; HRMS (ESI/QTOF) *m/z*: [M-Br]⁺ Calcd for [C₂₈H₃₂BrSi₂]⁺ 503.1220; Found 503.1209; [α]_D²⁰ = -111.8 (*c* = 1.0, CHCl₃); R_f: 0.52 (Pentane: EtOAc, 20:1); M_p = 136 °C.

Scheme S4: Synthesis of (S)-3a

(S)-(4,17-dihydro-1H-cyclopenta[6,7]cycloocta[2,1-a:3,4-a']dinaphthalene-5,16-

diyl)bis(trimethylsilane): To a stirred solution of NaH (23 mg, 0.941 mmol), 15-Crown-5 ether (377 mg, 1.71 mmol) and NaCp (470 μ l, 0.941 mmol, 2M in THF) in dry THF (9 ml) at -78 °C was added a solution of (S)-(2,2'-bis(bromomethyl)-[1,1'-binaphthalene]-3,3'-diyl)bis(trimethylsilane) (0.5 g, 0.86 mmol) in dry THF (9 ml). The mixture was stirred for 2h at -78°C and allowed to warm up to room temperature overnight. The reaction was quenched with a saturated solution of NH₄Cl and diluted with Et₂O. The aqueous layer was extracted with Et₂O, dried over Na₂SO₃ and concentrated under vacuum. The crude product was purified by column chromatography (SiO₂, Pentane/EtOAc, 100:0 to 50:1) to afford a mixture of **8** and **Rac-7** (ratio: 4:1, 0.390 g). This mixture was separate and charged under N₂ into two dried 30 ml dry Micro-wave tube. Both tubes were filled with K₂CO₃ (55 mg, 0.40 mmol) and freshly dry and degassed *o*-dichlorobenzene (20 ml) under N₂. The reaction tubes were heated by microwaves irradiation under stirring (*Biotage Initiator* reactor) at 250°C for 30 min. After being cooled to 25 °C, the reaction mixtures were combined and concentrated under vacuum at 50 °C. The residue was purified by column chromatography (SiO₂, Pentane/EtOAc, 100:0 to 20:1) affording (S)- (4,17-dihydro-1H-cyclopenta[6,7]cycloocta[2,1-a:3,4-a']dinaphthalene-5,16-diyl)bis(trimethylsilane) (0.38 g, 0.78 mmol, 90 % yield) as a slightly yellow foam. Slightly yellow solid, two inseparable double-bond isomers, ratio = 23:77.

¹H NMR (400 MHz, C₆D₆) δ 8.30 (s, 2H), 7.79 (d, *J* = 8.1 Hz, 2H), 7.25 (d, *J* = 8.3 Hz, 2H), 7.20 (t, *J* = 7.5 Hz, 2H), 6.96 (ddd, *J* = 8.2, 6.9, 1.1 Hz, 2H), 6.41 (d, *J* = 5.3 Hz, 0.23H), 6.16 (d, *J* = 5.1 Hz, 0.23H), 6.06 – 5.92 (m, 1.54H), 4.01 (d, *J* = 14.7 Hz, 1.54H), 3.78 (dd, *J* = 21.0, 15.2 Hz, 0.5H), 3.62 (d, *J* = 13.4 Hz, 1.54H), 3.57 – 3.42 (m, 0.5H), 3.11 (d, *J* = 23.6 Hz, 0.22H), 2.75 (s, 0.23H), 2.69 (dp, *J* = 3.3, 1.7 Hz, 2H), 0.42 (s, 16H), 0.35 (s, 2H); **¹³C NMR** (101 MHz, C₆D₆) δ 145.4, 141.2, 139.9, 139.7, 139.4, 138.9, 138.7, 138.0, 137.2, 137.0, 136.9, 136.7, 136.5, 133.5, 133.4, 132.4, 132.0, 129.6, 128.7, 128.6, 127.2, 127.0, 126.9, 125.7, 125.5, 125.5, 47.0, 38.6, 35.6, 34.2, 33.8, 1.4, 1.3, 1.2; **IR (ATR):** ν_{max} = 2923, 1558, 1444, 907, 883, 729 cm⁻¹; **HRMS (ESI/QTOF) m/z:** [M+H]⁺ Calcd for [C₃₃H₃₇Si₂]⁺ 489.2428; Found 489.2423; **[α]_D²⁰** = -378.7 (*c* = 0.3, CHCl₃); **R_f**: 0.36 (Pentane: EtOAc, 50:1); **Mp** = 238 °C;

Synthesis of compound (R)-3b:

(R)-diisopropyl 3,3'-diiodo-[1,1'-binaphthalene]-2,2'-dicarboxylate:

The compound was prepared (83 % yield), following the reported procedure and the analytical data was consistent with the literature.^[2] **¹H NMR** (400 MHz, CDCl₃) δ 8.52 (s, 2H), 7.78 (d, *J* = 8.3 Hz, 2H), 7.51 (ddd, *J* = 8.2, 6.9, 1.2 Hz, 2H), 7.34 (ddd, *J* = 8.2, 6.6, 1.3 Hz, 2H), 7.17 (d, *J* = 8.5 Hz, 2H), 4.76 (hept, *J* = 6.3 Hz, 2H), 0.76 (d, *J* = 6.2 Hz, 6H), 0.69 (d, *J* = 6.2 Hz, 6H); **¹³C NMR** (101 MHz, CDCl₃) δ 166.8, 139.6, 137.2, 134.5, 133.8, 132.1, 128.0, 127.8, 127.6, 126.8, 88.4, 69.2, 21.1, 20.9; **IR (ATR):** $\nu_{\max}$ = 2979, 1723, 1276, 1163, 1139, 1102, 749 cm⁻¹; **HRMS (ESI/QTOF) m/z:** [M+Na]⁺ Calc'd for [C₂₈H₂₄I₂NaO₄]⁺ 700.9656; Found 700.9660; **[α]_D²⁰** = +56.1 (*c* = 0.3, CHCl₃); **Rf:** 0.48 (Pentane: EtOAc, 10:1); **Mp** = 144 °C.

Scheme S5: Synthesis of (R)-(3,3'-diiodo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol

(R)-(3,3'-diiodo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol: To a stirred solution of diisopropyl 3,3'-diiodo-[1,1'-binaphthalene]-2,2'-dicarboxylate (2.25 g, 3.2 mmol) in THF (20 ml) at -78 °C under N₂, DIBALH (20% in Toluene, 22 ml, 26.5 mmol) was added dropwise. The reaction was slowly warmed up to room temperature and stirred overnight. The mixture was cooled down to 0 °C and carefully quenched with water, diluted with a saturated solution of Rochelle salt and stirred overnight. The remaining white suspension was filtered over Celite® and washed with warm THF. The aqueous layer was saturated with NaCl and extracted with CHCl₃ and then Et₂O. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The crude product was purified by trituration into EtOAc/Pentane (1:5) affording (3,3'-diiodo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol (1.75 g, 3.09 mmol, 93 % yield) as a white solid.

¹H NMR (400 MHz, CDCl₃) δ 8.61 (s, 2H), 7.82 (d, *J* = 8.3 Hz, 2H), 7.49 (ddd, *J* = 8.2, 6.8, 1.2 Hz, 2H), 7.27 (ddd, *J* = 8.3, 6.8, 1.3 Hz, 2H), 6.90 (d, *J* = 8.4 Hz, 2H), 4.61 (d, *J* = 12.2 Hz, 2H), 4.16 (d, *J* = 12.2 Hz, 2H); **¹³C NMR** (101 MHz, CDCl₃) δ 140.1, 138.0, 136.4, 134.4, 132.7, 127.7, 127.5, 127.1, 126.9, 98.3, 66.0; **IR (ATR):** $\nu_{\max}$ = 3286, 1027, 967, 890, 753 cm⁻¹; **HRMS (APPI/LTQ-Orbitrap) m/z:** [M-H₂O]⁺ Calc'd for [C₂₂H₁₅I₂O]⁺ 548.9207; Found 548.9208; **[α]_D²⁰** = +55.0 (*c* = 0.3, CHCl₃); **Rf:** 0.32 (Pentane: EtOAc, 4:1); **Mp** = 276 °C (decomposition);

Scheme S6: Synthesis of (R)-2b

(R)-2,2'-bis(bromomethyl)-3,3'-diiodo-1,1'-binaphthalene (R)-2b: To a solution of (3,3'-diiodo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol (1,5 g, 2,7 mmol) in dry THF (26 ml) under N₂ at 0 °C, PBr₃ (250 µl, 2.26 mmol) was added slowly. The reaction was stirred for 2 h at room temperature. The reaction was quenched at 0 °C with a saturated solution of NaHCO₃, diluted with EtOAc and extracted with EtOAc and then with CDCl₃. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The desired 3,3'-dibromo-2,2'-bis(bromomethyl)-1,1'-binaphthalene (0.130 g, 0.217 mmol, 79 % yield) was obtained in good purity as a white powder. Trituration into Pentane afford 2,2'-bis(bromomethyl)-3,3'-diiodo-1,1'-binaphthalene (1,5 g, 2,168 mmol, 82 % yield) as an off-white solid.

¹H NMR (400 MHz, CDCl₃) δ 8.65 (s, 2H), 7.82 (d, *J* = 8.3 Hz, 2H), 7.51 (ddd, *J* = 8.2, 6.8, 1.2 Hz, 2H), 7.29 (ddd, *J* = 8.3, 6.8, 1.3 Hz, 2H), 6.98 (d, *J* = 8.5 Hz, 2H), 4.41 (d, *J* = 10.4 Hz, 2H), 4.32 (d, *J* = 10.4 Hz, 2H); **¹³C NMR** (101 MHz, CDCl₃) δ 141.0, 136.5, 134.7, 134.4, 131.9, 128.0, 127.6, 127.5, 127.0, 97.9, 38.6; **IR (ATR):** ν_{max} = 1551, 1427, 1209, 983, 907, 887, 856, 749, 730, 571, 515 cm⁻¹; **HRMS (APPI/LTQ-Orbitrap) m/z:** Calc'd for [M-Br]⁺ = [C₂₂H₁₄BrI₂]⁺ 610.8363; Found 610.8361; **[α]_D²⁰** = +76.7 (c = 0.3, CHCl₃); **Mp** = 276 °C (decomposition).

Scheme S7: Synthesis of (R)-3b

(R)-5,16-diiodo-4,17-dihydro-2H-cyclopenta[6,7]cycloocta[2,1-a:3,4-a']dinaphthalene: To a stirred solution of NaH (11 mg, 0.477 mmol), 15-Crown-5 ether (191 mg, 0.87 mmol) and NaCp (238 μ l, 0.477 mmol, 2M in THF) in dry THF (4 ml) at -78 °C was added a solution of (R)-2,2'-bis(bromomethyl)-3,3'-diiodo-1,1'-binaphthalene (0.3 g, 0.43 mmol) in dry THF (4 ml). The mixture was stirred for 2h at -78°C and allowed to warm up to 0°C over 3h. The reaction was quenched with a saturated solution of NH₄Cl and diluted with Et₂O. The aqueous layer was extracted with Et₂O, dried over Na₂SO₃ and concentrated under vacuum. The crude product was purified by column chromatography (SiO₂, Pentane/EtOAc, 100:0 to 20:1) to afford a mixture of Cp and Spiro compounds (ratio: 1.5:1, 0.17 g). This mixture was separate and charged under N₂ into two dried 30 ml dry Micro-wave tube. Both tubes were filled with K₂CO₃ (20 mg, 0.15 mmol) and freshly dry and degassed *o*-dichlorobenzene (15 ml) under N₂. The reaction tubes were heated by microwaves irradiation under stirring (*Biotage Initiator* reactor) at 250°C for 45 min. After being cooled to 25 °C, the reaction mixtures were combined and concentrated under vacuum at 50 °C. The residue was purified by column chromatography (SiO₂, Pentane/EtOAc, 100:0 to 20:1) affording (R)-5,16-diiodo-4,17-dihydro-2H-cyclopenta[6,7]cycloocta[2,1-a:3,4-a']dinaphthalene (0.16 g, 0.26 mmol, 60 % yield) as a slightly yellow foam.

Slightly yellow solid, two inseparable double-bond isomers, ratio = 71:29. **¹H NMR** (400 MHz, C₆D₆) δ 8.41 (d, *J* = 5.9 Hz, 2H), 7.38 (dd, *J* = 8.2, 3.6 Hz, 2H), 7.14 (s, 1H), 7.11 – 7.02 (m, 3H), 6.94 – 6.83 (m, 2H), 6.60 (d, *J* = 5.3 Hz, 0.4H), 6.23 (d, *J* = 5.2 Hz, 0.4H), 6.21 – 6.18 (m, 1.2H), 4.04 (d, *J* = 14.5 Hz, 1.2H), 3.85 (dd, *J* = 29.4, 14.8 Hz, 0.8H), 3.37 (t, *J* = 16.8 Hz, 1.6H), 3.14 (d, *J* = 14.7 Hz, 0.8H), 2.74 (d, *J* = 25.2 Hz, 1.6H); **¹³C NMR** (101 MHz, C₆D₆) δ 141.7, 140.2, 140.1, 140.0, 138.9, 138.2, 137.8, 137.6, 137.2, 137.0, 136.8, 136.6, 136.5, 134.2, 133.8, 133.7, 132.0, 131.5, 131.5, 131.1, 130.2, 128.7, 128.6, 127.3, 127.3, 127.1, 126.4, 126.3, 126.3, 99.9, 99.8, 47.1, 40.4, 39.4, 38.8, 38.3; **IR (ATR):** ν_{max} = 3058, 2919, 1556, 976, 883, 748, 499 cm⁻¹; **HRMS (APPI/LTQ-Orbitrap) m/z:** [M]⁺ Calc'd for [C₂₇H₁₈I₂]⁺ 595.9492; Found 595.9513; [α]_D²⁰ = +172.8 (c = 0.52, CHCl₃); **Rf:** 0.2 (Pentane: EtOAc, 50:1); **Mp** = 279 °C (decomposition).

Synthesis of compound (R)-3c:

(R)-diisopropyl 3,3'-dibromo-[1,1'-binaphthalene]-2,2'-dicarboxylate

The compound was prepared (78 % yield) following the reported procedure and the analytical data was consistent with the literature.^[2] **¹H NMR** (400 MHz, CDCl₃) δ 8.24 (s, 2H), 7.82 (d, *J* = 8.2 Hz, 2H), 7.53 (t, *J* = 7.5 Hz, 2H), 7.35 (t, *J* = 8.2 Hz, 2H), 7.19 (d, *J* = 8.5 Hz, 2H), 4.78 (p, *J* = 6.3 Hz, 2H), 0.78 (d, *J* = 6.3 Hz, 6H), 0.67 (d, *J* = 6.3 Hz, 6H); **¹³C NMR** (101 MHz, CDCl₃) δ 165.6, 134.3, 134.3, 134.0, 132.4, 131.6, 128.2, 127.8, 127.4, 127.0, 116.1, 69.2, 21.1, 20.9; **IR (ATR):** $\nu_{\max}$ = 2980, 1724, 1275, 1102, 748 cm⁻¹; **HRMS (APCI/QTOF) m/z:** [M + H]⁺ Calcd for [C₂₈H₂₅Br₂O₄]⁺ 583.0114; Found 583.0114; **[α]_D²⁰** = +87.2 (*c* = 1.0, CHCl₃); **Rf:** 0.54 (Pentane: EtOAc, 9:1); **Mp** = 147 °C.

Scheme S8: Synthesis of (R)-(3,3'-dibromo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol

(R)- (3,3'-dibromo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol:^[4] To a stirred solution of diisopropyl 3,3'-dibromo-[1,1'-binaphthalene]-2,2'-dicarboxylate (120 mg, 0.2 mmol) in DCM (1 ml) at -78°C under N₂, DIBALH (20% in Toluene, 1.5 ml, 1.8 mmol) was added dropwise. The reaction was slowly warmed up to room temperature and stirred overnight. The mixture was cooled down to 0 °C and carefully quenched with saturated solution of NH₄Cl and extracted with Et₂O. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The crude product was purified by trituration into EtOAc/Pentane (1:5) affording (3,3'-dibromo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol (0.090 g, 0.191 mmol, 93 % yield) as a white solid.

¹H NMR (400 MHz, CDCl₃) δ 8.31 (s, 2H), 7.85 (d, *J* = 8.2 Hz, 2H), 7.50 (t, *J* = 8.0 Hz, 2H), 7.27 (t, *J* = 8.3 Hz, 2H), 6.92 (d, *J* = 8.5 Hz, 2H), 4.76 (d, *J* = 12.2 Hz, 2H), 4.07 (d, *J* = 12.2 Hz, 2H), 2.56 (br s, 2H); **¹³C NMR** (101 MHz, CDCl₃) δ 137.1, 136.1, 134.0, 132.7, 132.1, 127.7, 127.4, 127.4, 126.9, 123.2, 62.3; **IR (ATR):** $\nu_{\max}$ = 3274, 2987, 2901, 1385, 1029, 857, 755 cm⁻¹; **HRMS (APPI/LTQ-Orbitrap) m/z:** [M-OH]⁺ Calcd for [C₂₂H₁₅Br₂O]⁺ 452.9484; Found 452.9482; **[α]_D²⁰** = +62.8 (*c* = 0.3, CHCl₃); **Rf:** 0.32 (Pentane: EtOAc, 4:1); **Mp** = 278 °C.

Scheme S9: Synthesis of (R)-2c

(R)-3,3'-dibromo-2,2'-bis(bromomethyl)-1,1'-binaphthalene:^[4] To a solution of (3,3'-dibromo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol (130 mg, 0.275 mmol) in dry DCM (3 ml) under N₂ at 0 °C, PBr₃ (51.9 µl, 0.551 mmol) was added slowly. The reaction was stirred for 1h at room temperature. The reaction was quenched at 0 °C with a saturated solution of NaHCO₃, diluted with EtOAc and extracted with EtOAc. The combined organic layers were dried over Na₂SO₄ and concentrated under vacuum. The desired 3,3'-dibromo-2,2'-bis(bromomethyl)-1,1'-binaphthalene (0.130 g, 0.217 mmol, 79 % yield) was obtained in good purity as a white powder.

¹H NMR (400 MHz, CDCl₃) δ 8.35 (s, 2H), 7.86 (d, *J* = 8.2 Hz, 2H), 7.56 – 7.49 (m, 2H), 7.30 (ddd, *J* = 8.2, 6.9, 1.1 Hz, 2H), 7.00 (d, *J* = 8.5 Hz, 2H), 4.40 (d, *J* = 10.3 Hz, 2H), 4.33 (d, *J* = 10.3 Hz, 2H); **¹³C NMR** (101 MHz, CDCl₃) δ 137.0, 134.3, 133.4, 132.8, 131.4, 128.2, 127.5, 127.4, 127.2, 122.8, 33.1; **IR (ATR):** ν_{max} = 2987, 1554, 1212, 905, 729cm⁻¹; **HRMS (LTQ-Orbitrap) m/z:** [M-Br]⁺ Calcd for [C₂₂H₁₄Br₃]⁺ 514.8640; Found 514.8647; **[α]_D²⁰** = +106.5 (c = 1.0, CHCl₃); **Mp** = 214 °C.

Scheme S10: Synthesis of (R)-3c

(R)-5,16-dibromo-4,17-dihydro-2H-cyclopenta[6,7]cycloocta[2,1-a:3,4-a']dinaphthalene: To a stirred solution of NaH (7.4 mg, 0.31 mmol), 15-Crown-5 ether (125 mg, 1.71 mmol) and NaCp (156 μ l, 0.31 mmol, 2M in THF) in dry THF (3 ml) at -78 °C was added a solution of (R)-3,3'-dibromo-2,2'-bis(bromomethyl)-1,1'-binaphthalene (170 g, 0.28 mmol) in dry THF (3 ml). The mixture was stirred for 2h at -78°C and allowed to warm up to room temperature overnight. The reaction was quenched with a saturated solution of NH₄Cl and diluted with Et₂O. The aqueous layer was extracted with Et₂O, dried over Na₂SO₃ and concentrated under vacuum. The crude product was purified by column chromatography (SiO₂, Pentane/EtOAc, 100:0 to 50:1) to afford a mixture of Cp and spiro compounds (ratio = 1:1.5, 80 mg). This mixture was charged under N₂ into two dried 2-5 ml dry Micro-wave tubes. Both tubes were filled with K₂CO₃ (11 mg, 0.08 mmol) and freshly dry and degassed *o*-dichlorobenzene (4 ml) under N₂. The reaction tubes were heated by microwaves irradiation under stirring (*Biotage Initiator* reactor) at 250°C for 45 min. After being cooled to 25 °C, the reaction mixtures were combined and concentrated under vacuum at 50 °C. The residue was purified by column chromatography (SiO₂, Pentane/EtOAc, 100:0 to 20:1) affording (R)-5,16-dibromo-4,17-dihydro-2H-cyclopenta[6,7]cycloocta[2,1-a:3,4-a']dinaphthalene (74 mg, 0.12 mmol, 43 % yield) as a slightly yellow foam.

Two inseparable double-bond isomers, ratio = 1:1; **¹H NMR** (400 MHz, C₆D₆) δ 8.10 – 8.04 (m, 2H), 7.49 – 7.38 (m, 2H), 7.21 – 7.17 (m, 2H), 7.12 – 7.06 (m, 2H), 6.96 – 6.84 (m, 2H), 6.49 (d, *J* = 5.3 Hz, 0.5H), 6.19 (d, *J* = 5.1 Hz, 0.5H), 6.13 (s, 1H), 4.18 (d, *J* = 14.4 Hz, 1H), 4.01 (d, *J* = 14.8 Hz, 0.5H), 3.92 (d, *J* = 15.0 Hz, 0.5H), 3.30 – 3.13 (m, 1.5H), 2.97 (t, *J* = 11.7 Hz, 1H), 2.76 – 2.67(m, 1.5H); **¹³C NMR** (101 MHz, C₆D₆) δ 141.9, 138.9, 138.0, 137.9, 137.8, 137.5, 137.5, 136.1, 134.9, 134.5, 133.7, 133.3, 133.2, 132.7, 132.6, 132.5, 131.5, 131.1, 131.0, 130.6, 130.2, 127.5, 127.3, 127.1, 126.7, 126.5, 124.2, 123.7, 123.6, 46.8, 39.3, 35.7, 33.8, 33.4; **IR (ATR):** ν_{max} = 3057, 939, 748, 499 cm⁻¹; **HRMS (APPI/LTQ-Orbitrap) m/z:** [M]⁺ Calcd for [C₂₇H₁₈Br₂]⁺ 499.9770; Found 499.9775; **[α]_D²⁰** = + 152.6 (*c* = 1.0, CHCl₃); **R_f**: 0.92 (Pentane: EtOAc, 9:1); **Mp** = 264 °C;

Synthesis of compounds (S)-9 and (R)-9:

Scheme S11: Synthesis of 2-iodo-3,6-dimethoxybenzoic acid

2-iodo-3,6-dimethoxybenzoic acid: *n*-Butyllithium (2.5M in hexanes, 35.1 ml, 88.0 mmol) was added to a solution of freshly distilled 2,2,6,6-tetramethylpiperidine (15.0 ml, 88.0 mmol) in THF (10ml) at 0°C under nitrogen and the reaction was stirred for 30min. The reaction was cooled down to -78 °C and a solution of 2,5-dimethoxybenzoic acid **5** (4.0 g, 22.0 mmol) in dry THF (10ml) was added. The reaction was warm up to 0 °C and stirred for 2.5 h. The reaction was cooled to -78°C and a solution of iodine (22.3 g, 88.0 mmol) in THF (60ml) was added and the reaction was warm up to room temperature over 1 h. The reaction was diluted into HCl 1N and NaHSO₃ (10% solution) and extracted with ethyl acetate. The organic layer was washed with a saturated solution of Na₂CO₃ (3 times) and the aqueous layer was acidified with a 37% aqueous HCl solution until a white precipitate appears. This solid was filtered off, was washed with toluene and dried to afford 2-iodo-3,6-dimethoxybenzoic acid (5.3 g, 14.0 mmol, 78 % yield) as a white solid.

¹H NMR (400 MHz, CD₃OD) δ 7.05 (d, *J* = 9.0 Hz, 1H), 6.95 (d, *J* = 9.0 Hz, 1H), 3.83 (s, 3H), 3.80 (s, 3H); **¹³C NMR** (101 MHz, CD₃OD) δ 171.1, 154.1, 151.6, 134.5, 113.4, 112.6, 85.1, 57.4, 57.1; **IR (ATR):** ν_{max} = 3004, 2937, 2838, 1706, 1571, 1473, 1420, 1259, 1056, 1029, 804, 716 cm⁻¹; **HRMS (ESI/QTOF) *m/z*:** [M-H]⁻ calc'd for [C₉H₈IO₄]⁻ 306.9473; Found 306.9471; **R_f:** 0.32 (Pentane: EtOAc:AcOH, 1:1:1%); **Mp** = 201-202 °C.

Scheme S12: Synthesis of **6**

methyl iodo-3,6-dimethoxybenzoate **6:** To a stirred solution of 2-iodo-3,6-dimethoxybenzoic acid (5.00 g, 16.23 mmol) in DCM (50 mL), oxalyl chloride (1.5 mL, 17.04 mmol) was added followed by three drops of dry DMF. The mixture was stirred for 3 h, after which the solvent and excess oxalyl chloride were removed in vacuo. Dry Methanol (30 mL) was added to the residue, and the resulting solution was stirred for 30 min. The reaction was diluted into DCM (70 mL) and the mixture was washed with water, brine, dried over Na_2SO_4 and then concentrated under vacuum. Purification of the reaction crude by column chromatography (Hexane/EtOAc, 5/1) afforded 2-methyl iodo-3,6-dimethoxybenzoate **6** as a white solid (4.7 g, 14.59 mmol, 90%).

^1H NMR (400 MHz, CDCl_3) δ 6.89 (d, $J = 9.0$ Hz, 1H), 6.81 (d, $J = 9.0$ Hz, 1H), 3.96 (s, 3H), 3.84 (s, 3H), 3.79 (s, 3H); **^{13}C NMR** (101 MHz, CDCl_3) δ 167.9, 152.6, 150.8, 131.9, 112.0, 111.9, 85.6, 57.3, 56.8, 53.1; **IR (ATR):** $\nu_{\text{max}} = 3001, 2950, 2838, 1735, 1570, 1474, 1430, 1281, 1259, 1058, 1030, 803, 725$ cm^{-1} ; **HRMS (APCI/QTOF) m/z :** $[\text{M}+\text{H}]^+$ calc'd for $\text{C}_{10}\text{H}_{12}\text{IO}_4^+$ 322.9775; Found 322.9773; **Rf:** 0.71 (Pentane: EtOAc, 1:1); **Mp** = 115-116 $^\circ\text{C}$.

Scheme S13: Synthesis of **7**

dimethyl 3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-dicarboxylate 7: A sealed tube was charged with methyl 2-iodo-3,6-dimethoxybenzoate **6** (5 g, 15.52 mmol) and triphenylphosphine (4.07 g, 15.52 mmol) and a nitrogen atmosphere was built up. Then activated zinc powder (3.04 g, 46.6 mmol), $\text{NiCl}_2(\text{PPh}_3)_2$ (5.08 g, 7.76 mmol) and Toluene (Volume: 60 ml) were added under nitrogen atmosphere in a glovebox and the tube was sealed. The reaction mixture was heated at 90 °C for 6 h. After cooling down to room temperature, the reaction was quenched with 1M HCl, and then extracted with ethyl acetate (2 x 50 ml). The combined organic extracts were washed by a saturated solution of NaHCO_3 , dried over (MgSO_4), filtered and then evaporated under vacuum. The crude product was purified by column chromatography (Pentane/EtOAc, 1:1 to 0:1) affording dimethyl 3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-dicarboxylate **7** (2.5 g, 6.4 mmol, 82 % yield) as a white foam.

^1H NMR (400 MHz, CDCl_3) δ 6.95 (d, $J = 9.0$ Hz, 2H), 6.88 (d, $J = 9.0$ Hz, 2H), 3.81 (s, 6H), 3.70 (s, 6H), 3.56 (s, 6H); **^{13}C NMR** (101 MHz, CDCl_3) δ 166.9, 151.7, 150.8, 125.5, 124.7, 113.5, 111.8, 57.0, 56.6, 51.9; **IR (ATR):** ν_{max} = 2950, 2837, 1731, 1478, 1431, 1278, 1257, 1062, 1026, 809, 726 cm^{-1} ; **HRMS (ESI/QTOF) m/z :** $[\text{M}+\text{Na}]^+$ calc'd for $[\text{C}_{20}\text{H}_{22}\text{NaO}_8]^+$ 413.1207; Found 413.1202; **Rf:** 0.45 (Pentane: EtOAc, 1:1); **Mp** = 133 °C.

Scheme S14: Synthesis of (3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-diyl)dimethanol

(3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-diyl)dimethanol: To a solution of dimethyl 3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-dicarboxylate **7** (3 g, 7.68 mmol) in dry THF (71.7 ml) under nitrogen at 0°C, LAH (2.4 M in THF, 3.29 ml, 7.89 mmol) was added slowly. The reaction was run at 0°C for 2h. The reaction was carefully quenched with an aqueous Rochelle salt solution and extracted with DCM. The combined organic layers were dried over MgSO₄ and concentrated under vacuum. The crude product was purified by column chromatography (DCM/MeOH 20:1 to 10:1) to afford (3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-diyl)dimethanol (2.26 g, 6.76 mmol, 88 % yield) as a white solid.

¹H NMR (400 MHz, CDCl₃) δ 6.90 (s, 4H), 4.50 (d, *J* = 11.6 Hz, 2H), 4.05 (d, *J* = 11.6 Hz, 2H), 3.88 (s, 6H), 3.65 (s, 6H), 3.15 (s, 2H); **¹³C NMR** (101 MHz, CDCl₃) δ 152.6 (2C), 151.1 (2C), 129.2 (2C), 126.6 (2C), 111.5 (2C), 110.4 (2C), 58.0 (2C), 56.7 (2C), 55.9 (2C); **IR (ATR):** *v*_{max} = 3340, 2940, 2834, 1593, 1477, 1464, 1434, 1253, 1080, 1030, 802, 725 cm⁻¹; **HRMS (ESI)** calc'd for [M+Na]⁺ = [C₁₈H₂₂O₆Na]⁺: 357.1309, found : 357.1321; **Rf**: 0.24 (Pentane: EtOAc, 1:1); **Mp** = 191 °C.

Scheme S15: Synthesis of 8

2,2'-bis(bromomethyl)-3,3',6,6'-tetramethoxy-1,1'-biphenyl **8:** To a solution of (3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-diyl)dimethanol (1.75 g, 5.23 mmol) in dry DCM (52 ml) at 0 °C under nitrogen, phosphorus tribromide (0.494 ml, 5.23 mmol) was added dropwise. The reaction was stirred at 0 °C for 2 h. The reaction was quenched with a saturated solution of NaHCO₃ and extracted with DCM. The combined organic layers were dried over MgSO₄ and concentrated under vacuum. The residue was dissolved in DCM (2 ml) and filtered over a Silica pad and eluted with DCM to afford 2,2'-bis(bromomethyl)-3,3',6,6'-tetramethoxy-1,1'-biphenyl **8** (2.04 g, 4.43 mmol, 85 % yield) as a white solid.

¹H NMR (400 MHz, Benzene-*d*₆) δ 6.53 (d, *J* = 9.0 Hz, 2H), 6.45 (d, *J* = 9.0 Hz, 2H), 4.55 (d, *J* = 9.3 Hz, 2H), 4.48 (d, *J* = 9.4 Hz, 2H), 3.37 (s, 6H), 3.29 (s, 6H); **¹³C NMR** (101 MHz, Benzene-*d*₆) δ 152.6, 151.5, 126.8, 126.3, 111.6, 111.0, 55.5, 55.4, 27.3; **IR (ATR):** *v*_{max} = 2997, 2936, 2833, 1592, 1477, 1435, 1257, 802, 717, 577 cm⁻¹; **HRMS (ESI)** calc'd. for [M-Br]⁺ = [C₁₈H₂₀BrO₄]⁺: 379,0539, found: 379,0535; **Rf**: 0.79 (Pentane: EtOAc, 1:1); **Mp** = 195-196 °C.

Scheme S16: Synthesis of *Rac-9* and *Rac-10*

(*Rac*)-5,8,9,12-tetramethoxy-4,13-dihydro-2H-dibenzo[a,c]cyclopenta[f][8]annulene *Rac-9* and (*Rac*)-1',4',8',11'-tetramethoxy-5',7'-dihydrospiro[cyclopenta[2,4]diene-1,6'-dibenzo[a,c][7]annulene] *Rac-10*: To a cold (-78 °C) suspension of sodium hydride (57.4 mg, 2.390 mmol) and 15-CROWN-5 (957 mg, 4.35 mmol) in THF (20 ml) was added dropwise a solution of sodium cyclopentadienylide (1.195 ml, 2.390 mmol, 1.0M in THF) were added followed by a solution of 2,2'-bis(bromomethyl)-3,3',6,6'-tetramethoxy-1,1'-biphenyl (1000 mg, 2.173 mmol) in dry THF (20 ml). The mixture was warmed up slowly and stirred for 14 hours, then quenched carefully with a saturated solution of NH₄Cl (5.0 mL) and extracted with ether (2 X 50.0 mL). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure. The crude was purified by flash column chromatography (Biotage SNAP, Pent:EtOAc 5:95 -> 15:85, 15 CV) to afford (*Rac*)-5,8,9,12-tetramethoxy-4,13-dihydro-2H-dibenzo[a,c]cyclopenta[f][8]annulene (87:13 double bond isomers ratio, 499 mg, 1,391 mmol, 64,0 % yield) **9** as a white solid and (*Rac*)-1',4',8',11'-tetramethoxy-5',7'-dihydrospiro[cyclopenta[2,4]diene-1,6'-dibenzo[a,c][7]annulene] **10** (269 mg, 0,749 mmol, 34,5 % yield) as a white solid.

(*Rac*)-5,8,9,12-tetramethoxy-4,13-dihydro-2H-dibenzo[a,c]cyclopenta[f][8]annulene **9:** ¹H NMR (400 MHz, C₆D₆) δ 6.67 – 6.56 (m, 4H), 6.54 (d, *J* = 5.3 Hz, 0.25H), 6.15 (s, 1.5H), 6.11 (d, *J* = 5.2 Hz, 0.25H), 4.36 (d, *J* = 13.8 Hz, 1.5H), 4.18 (d, *J* = 13.8 Hz, 0.25H), 4.03 (d, *J* = 14.1 Hz, 0.25H), 3.45 (s, 0.75H), 3.44 (s, 0.75H), 3.41 (s, 4.5H), 3.37 (s, 4.5H), 3.35 (s, 1.5H), 3.32 (s, 1.5H), 3.10 (d, *J* = 14.9 Hz, 0.75H), 2.83 (d, *J* = 22.6 Hz, 0.25H), 2.65 (s, 1.5H); ¹³C NMR (101 MHz, C₆D₆) δ 152.6, 151.8, 151.6, 151.4, 145.2, 140.5, 139.0, 138.5, 129.3, 129.2, 129.2, 129.0, 128.8, 128.5, 128.3, 128.1, 127.8, 127.4, 110.8, 110.7, 109.4, 108.7, 108.6, 55.9, 55.7, 55.7, 55.6, 47.0, 38.8, 29.1, 26.9, 26.7; IR (ATR): ν_{max} = 2937, 2832, 1481, 1463, 1433, 1255, 1073, 1033, 797 cm⁻¹; HRMS (ESI) calc'd. for [M+H]⁺ = [C₂₃H₂₅O₄]⁺: 365.1747, found: 365.1744; R_f: 0.27 (Pentane: EtOAc, 9:1); Mp = 144 °C.

(Rac)-1',4',8',11'-tetramethoxy-5',7'-dihydrospiro[cyclopenta[2,4]diene-1,6'-

dibenzo[a,c][7]annulene] 10: $^1\text{H NMR}$ (400 MHz, C_6D_6) δ 6.73 (d, J = 8.9 Hz, 2H), 6.66 (d, J = 8.9 Hz, 2H), 6.40 (d, J = 6.4 Hz, 2H), 6.22 (d, J = 6.4 Hz, 2H), 3.50 (s, 6H), 3.40 (s, 6H), 3.03 (d, J = 12.9 Hz, 2H), 2.68 (d, J = 12.8 Hz, 2H); $^{13}\text{C NMR}$ (101 MHz, C_6D_6) δ 152.1, 151.8, 144.4, 128.9, 128.3, 128.1, 127.8, 110.7, 110.2, 67.2, 56.1, 55.8, 29.1; **IR (ATR):** ν_{max} = 2936, 2831, 1486, 1464, 1436, 1256, 1227, 1148, 1077, 1031, 799, 758, 718 cm^{-1} ; **HRMS (APPI/LTQ-Orbitrap) m/z:** $[\text{M}]^+$ Calcd for $[\text{C}_{23}\text{H}_{25}\text{O}_4]^+$ 365.1747; Found 365.1748; **Rf:** 0.22 (Pentane: EtOAc, 9:1); **Mp** = 147 $^{\circ}\text{C}$;

Resolution of (Rac)-9:

Separation of the enantiomers of the two pairs of isomers by preparative chiral HPLC (8 X 50mg with Chiralpak IA, 2 x 25 cm; 2:98 *i*-PrOH / *n*-hexane, 18 mL/min, 254 nm) provided:

- (S)-9 (180 mg, mmol, 45 %, >99 % ee)
- (R)-9 (180 mg, mmol, 45 %, >99 % ee)

(R)-5,8,9,12-tetramethoxy-4,13-dihydro-1H-dibenzo[a,c]cyclopenta[f][8]annulene (R)-9:

$[\alpha]_{\text{D}}^{20}$ = 304.3 (c = 0.35, CHCl_3); **Chiral HPLC:** (Chiralpak IA, 4.6 x 250 mm; 2% *i*-PrOH / hexane, 1.0 mL/min, 254 nm, t_{r} (first isomer) = 6.4 min, t_{r} (second isomer) = 6.6 min). Purity was confirmed by $^1\text{H NMR}$.

(S)-5,8,9,12-tetramethoxy-4,13-dihydro-1H-dibenzo[a,c]cyclopenta[f][8]annulene (S)-9:

$[\alpha]_{\text{D}}^{20}$ = -317.1 (c = 0.35, CHCl_3); **Chiral HPLC:** (Chiralpak IA, 4.6 x 250 mm; 2% *i*-PrOH / hexane, 1.0 mL/min, 254 nm, t_{r} (first isomer) = 10.0 min, t_{r} (second isomer) = 12.4 min). Purity was confirmed by $^1\text{H NMR}$.

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.355	MM	0.1632	645.01465	65.85848	25.0176
2	6.546	VB	0.1510	625.29187	61.30387	24.2526
3	9.770	BB	0.2441	662.52832	39.01539	25.6969
4	12.056	BB	0.3019	645.40845	29.89223	25.0329

Figure S1: HPLC trace of Rac-9 – mixture of double bond isomers

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.395	BV	0.1325	196.10138	21.48214	31.3093
2	6.588	VB	0.1583	430.23392	39.73522	68.6907

Figure S2: HPLC trace of (R)-9 – mixture of double bond isomers

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	10.043	BB	0.2114	74.99676	4.57064	26.7234
2	12.379	BB	0.2975	205.64409	9.61416	73.2766

Figure S3: HPLC trace of (S)-**9** – mixture of double bond isomers

Absolute Configuration Determination:

The absolute configurations were determined upon complexation of fraction (S)-**9** to rhodium(I) and crystallization of the resulting **Rh4** complex.

Figure S4 : Crystal structures of **Rh4** as ORTEP drawing with 30 % probability thermal ellipsoids

Complexes synthesis

General procedure for the synthesis of $Cp^xRh(cod)$ complexes.^[5]

Without any precaution to air or to moisture, Cp^xH (1 eq.) and $[Rh(cod)OAc]_2$ (0.5 eq.) dissolved in a mixture of MeOH/Toluene (0.025M, 1:3 or 0.05M, 1:1). The mixture was stirred at 25 °C or 70 °C for the indicated time and then solvents were removed under vacuum (water bath at 30 °C). The crude material was filtered over a pad silica and eluted with toluene, and concentrated under vacuum to give $Cp^xRh(COD)$ complex as a yellow solid.

Complex Rh1:

(*S*)-**3a** (0.040 g, 0.082 mmol) and $[Rh(cod)OAc]_2$ (0.027 g, 0.049 mmol) in MeOH/Toluene (0.05M, 1:1) at 70°C for 2h to give **Rh1** (0.057 g, 0.082 mmol, >99 % yield) as a yellow solid. **¹H NMR** (400 MHz, C_6D_6) δ 8.37 (s, 1H), 8.30 (s, 1H), 7.76 (dd, J = 7.9, 4.9 Hz, 2H), 7.12 (m, 3H), 6.91 (p, J = 7.5 Hz, 3H), 5.64 (s, 1H), 4.82 (s, 1H), 4.36 (t, J = 2.5 Hz, 1H), 3.79 (d, J = 14.5 Hz, 1H), 3.69 (d, J = 14.4 Hz, 1H), 3.50 (d, J = 14.5 Hz, 1H), 3.37 (dt, J = 7.9, 4.7 Hz, 2H), 3.13 (d, J = 14.4 Hz, 1H), 3.07 (t, J = 7.6 Hz, 2H), 2.24 (ddd, J = 18.1, 10.3, 5.5 Hz, 2H), 2.10 (ddt, J = 18.6, 7.9, 4.4 Hz, 2H), 1.98 – 1.81 (m, 2H), 1.67 (dq, J = 13.2, 7.0 Hz, 2H), 0.67 (s, 9H), 0.50 (s, 9H); **¹³C NMR** (101 MHz, C_6D_6) δ 142.3, 140.7, 138.6, 137.2, 137.0, 136.6, 136.2, 136.2, 133.8, 133.3, 132.3, 132.1, 128.5, 128.5, 127.9, 127.3, 127.3, 126.5, 126.4, 125.6, 109.9 (d, J_{Rh-C} = 3.4 Hz), 97.3 (d, J_{Rh-C} = 3.5 Hz), 91.9 (d, J_{Rh-C} = 3.3 Hz), 87.2 (d, J_{Rh-C} = 3.9 Hz), 81.4 (d, J_{Rh-C} = 4.1 Hz), 65.7 (d, J_{Rh-C} = 14.3 Hz), 64.8 (d, J_{Rh-C} = 14.0 Hz), 33.8, 33.6, 32.3, 31.6, 1.7; **IR (ATR):** ν_{max} = 2952, 1249, 836, 755 cm^{-1} ; **HRMS (ESI/QTOF) m/z :** $[M+H]^+$ Calcd for $[C_{41}H_{48}RhSi_2]^+$ 698.2266; Found 698.2269; **$[\alpha]_D^{20}$** = +174.4 (c = 0.3, $CHCl_3$); **Mp** = 134°C.

Complex Rh2:

(*R*)-**3b** (0.040 g, 0.067 mmol) and $[Rh(cod)OAc]_2$ (0.022 g, 0.040 mmol) in MeOH/Toluene (0.025M, 1:3) at 70°C for 3h to give **Rh2** (0.050 g, 0.061 mmol, 91 % yield) as a yellow solid. **¹H NMR** (600 MHz, C_6D_6) δ 8.63 (s, 1H), 8.37 (s, 1H), 7.43 (d, J = 8.2 Hz, 1H), 7.35 (d, J = 8.2 Hz, 1H), 7.06 – 7.01 (m, 2H), 6.98 (d, J = 8.4 Hz, 1H), 6.91 (d, J = 8.4 Hz, 1H), 6.83 (ddd, J = 8.3, 6.8, 1.1 Hz, 1H), 6.80 (ddd, J = 8.3, 6.8, 1.1 Hz, 1H), 5.80 (s, 1H), 4.65 (s, 1H), 4.58 (t, J = 2.6 Hz, 1H), 4.10 (d, J = 14.8 Hz, 1H), 3.69 (t, J = 7.9 Hz, 2H),

3.52 (d, $J = 14.3$ Hz, 1H), 3.26 (d, $J = 14.9$ Hz, 1H), 3.14 (t, $J = 7.8$ Hz, 2H), 2.76 (d, $J = 14.2$ Hz, 1H), 2.47 (ddd, $J = 12.8, 9.3, 6.3$ Hz, 2H), 2.39 (ddt, $J = 11.3, 7.7, 5.7$ Hz, 2H), 1.93 – 1.82 (m, 4H); ^{13}C NMR (151 MHz, C_6D_6) δ 140.2, 139.9, 137.6, 137.5, 136.7, 136.3, 133.9, 133.7, 132.0, 132.0, 128.6, 128.5, 128.3, 127.4, 127.3, 127.3, 127.2, 127.0, 127.0, 126.5, 126.3, 106.0 (d, $J_{\text{Rh-C}} = 3.8$ Hz), 102.0, 98.8, 94.0 (d, $J_{\text{Rh-C}} = 3.6$ Hz), 89.7 (d, $J_{\text{Rh-C}} = 3.2$ Hz), 89.3 (d, $J_{\text{Rh-C}} = 3.5$ Hz), 83.4 (d, $J_{\text{Rh-C}} = 3.8$ Hz), 65.7 (d, $J_{\text{Rh-C}} = 14.4$ Hz), 64.8 (d, $J_{\text{Rh-C}} = 14.0$ Hz), 38.5, 36.5, 32.9 (2C), 32.6 (2C); IR (ATR): $\nu_{\text{max}} = 2924, 758, 745, 498$ cm^{-1} ; HRMS (ESI/QTOF) m/z : $[\text{M}+\text{H}]^+$ calc'd. for $[\text{C}_{35}\text{H}_{29}\text{I}_2\text{Rh}]^+$ 805.9408; Found 805.9408; $[\alpha]_{\text{D}}^{20} = -317.5$ ($c = 0.3$, CHCl_3); **Mp** = 278 °C (decomposition).

Complex Rh3:

(*R*)-**3c** (0.042 g, 0.084 mmol) and $[\text{Rh}(\text{cod})\text{OAc}]_2$ (0.023 g, 0.042 mmol) in MeOH/Toluene (0.025M, 1:3) at 70°C for 1h to give **Rh3** (0.052 g, 0.072 mmol, 86 % yield) as a yellow solid. ^1H NMR (400 MHz, C_6D_6) δ 8.29 (s, 1H), 8.03 (s, 1H), 7.47 (d, $J = 8.2$ Hz, 1H), 7.39 (d, $J = 8.2$ Hz, 1H), 7.06 (ddt, $J = 8.3, 6.9, 1.4$ Hz, 2H), 7.02 (d, $J = 8.0$ Hz, 1H), 6.96 (d, $J = 7.8$ Hz, 1H), 6.87 – 6.80 (m, 2H), 5.62 (s, 1H), 4.60 (t, $J = 2.6$ Hz, 1H), 4.58 (s, 1H), 4.23 (d, $J = 14.7$ Hz, 1H), 3.71 (ddd, $J = 10.5, 7.8, 2.6$ Hz, 2H), 3.62 (d, $J = 14.0$ Hz, 1H), 3.17 (ddd, $J = 10.4, 7.8, 2.7$ Hz, 2H), 3.11 (d, $J = 14.7$ Hz, 1H), 2.63 (d, $J = 14.0$ Hz, 1H), 2.47 (tdd, $J = 12.3, 7.2, 5.2$ Hz, 2H), 2.31 (dtd, $J = 16.4, 7.4, 4.7$ Hz, 2H), 1.90 (ddq, $J = 22.0, 13.3, 4.9, 4.4$ Hz, 4H); ^{13}C NMR (101 MHz, C_6D_6) δ 137.9, 137.0, 135.8, 133.4, 133.3, 132.7, 132.4, 131.5, 131.4, 127.4, 127.2, 127.1, 127.1, 127.0, 126.7, 126.5, 125.2, 123.1, 105.5 (d, $J_{\text{Rh-C}} = 4.0$ Hz), 94.6 (d, $J_{\text{Rh-C}} = 3.6$ Hz), 89.2 (d, $J_{\text{Rh-C}} = 3.5$ Hz), 89.1 (d, $J_{\text{Rh-C}} = 3.7$ Hz), 83.4 (d, $J_{\text{Rh-C}} = 4.0$ Hz), 65.6 (d, $J_{\text{Rh-C}} = 14.2$ Hz), 64.5 (d, $J_{\text{Rh-C}} = 14.0$ Hz), 33.3, 33.0, 32.4, 31.7; IR (ATR): $\nu_{\text{max}} = 2930, 759, 746, 498$ cm^{-1} ; HRMS (APCI/QTOF) m/z : $[\text{M} + \text{H}]^+$ Calcd for $[\text{C}_{35}\text{H}_{28}\text{Br}_2\text{Rh}]^+$ 708.9607; Found 708.9599; $[\alpha]_{\text{D}}^{20} = -221.5$ ($c = 0.3$, CHCl_3); **Mp** = 293 °C.

Complex Rh4:

(*S*)-**9** (0.020 g, 0.055 mmol) and $[\text{Rh}(\text{cod})\text{OAc}]_2$ (0.016 g, 0.030 mmol) in MeOH/Toluene (0.05M, 1:1) at 25°C for 1h to give **Rh4** (0.031 g, 0.054 mmol, 98 % yield) as a yellow solid. ^1H NMR (400 MHz, C_6D_6) δ 6.65 (q, $J = 8.9$ Hz, 2H), 6.57 (s, 2H), 5.35 (s, 1H), 4.66 (s, 1H), 4.64 (t, $J = 2.5$ Hz, 1H), 4.27 (d, $J = 14.3$ Hz, 1H), 3.90 (ddd, $J = 10.4, 7.7, 2.7$ Hz, 2H), 3.78 (d, $J = 13.3$ Hz, 1H), 3.56 (dt, $J = 6.8, 3.1$ Hz, 2H), 3.52 (s, 3H),

3.38 (s, 3H), 3.37 (s, 6H), 3.20 (d, $J = 14.2$ Hz, 1H), 2.88 (d, $J = 13.2$ Hz, 1H), 2.43 – 2.27 (m, 2H), 2.17 (ddt, $J = 12.7, 10.1, 4.9$ Hz, 2H), 2.12 – 2.01 (m, 2H), 2.01 – 1.89 (m, 2H); ^{13}C NMR (101 MHz, C_6D_6) δ 152.0, 151.7, 151.5, 151.4, 129.5, 129.4, 128.7, 128.2, 110.7, 109.7, 108.9, 108.9, 107.3 (d, $J = 3.9$ Hz), 99.4 (d, $J_{\text{Rh-C}} = 3.4$ Hz), , 88.3 (d, $J_{\text{Rh-C}} = 4.6$ Hz), 88.2 (d, $J_{\text{Rh-C}} = 4.1$ Hz), 82.0 (d, $J_{\text{Rh-C}} = 4.2$ Hz), 65.0 (d, $J = 14.2$ Hz), 64.2 (d, $J = 14.0$ Hz), 55.9, 55.8, 55.7, 55.0, 33.5, 32.4, 26.1, 25.3. **IR (ATR):** $\nu_{\text{max}} = 2988, 2931, 2870, 2828, 1480, 1433, 1321, 1255, 1224, 1175, 1141, 1074, 1034, 795, 500$ cm^{-1} ; **HRMS (ESI) m/z :** $[\text{M}+\text{H}]^+$ calc'd. for $[\text{C}_{31}\text{H}_{36}\text{O}_4\text{Rh}]^+$: 575.1668, found: 575.1661; $[\alpha]_{\text{D}}^{20} = +171$ ($c = 0.3$, CHCl_3); **Mp** = 161 °C.

Representative procedure for the synthesis of $\text{Cp}^x\text{Co}(\text{CO})\text{I}_2$ complexes.^[6]

In an N_2 -filled glovebox, a flame dried Schlenk tube was charged with $\text{Co}_2(\text{CO})_8$ (1.1 equiv.). A solution of Cp^xH ligand (1.0 equiv.) in dry DCM (0.02 M) was added and the black reaction mixture was stirred at 37 °C for 6 h. The volatiles were evaporated by Schlenk technique. A solution of iodine (1.20 equiv.) in dry degassed Et_2O (0.02 M) was added to the black residue. The reaction vessel was connected to a Schlenk line to release carbon monoxide and the dark violet mixture was stirred at room temperature for 1 h. All volatiles were evaporated and the residue was purified by column chromatography on silica gel (gradient: DCM/pentane = 0% $\rightarrow$ 50%) and further trituration in Et_2O . The dark violet fractions were collected and concentrated under reduced pressure to obtain $\text{Cp}^x\text{Co}(\text{CO})\text{I}_2$ complex as a black solid.

Complex Co1:

Following the general procedure using (*R*)-**3b** (25 mg, 42 μmol), $\text{Co}_2(\text{CO})_8$ (16 mg, 46 μmol) and iodine (13 mg, 50 μmol) gave complex **Co1** (27 mg, 29 μmol , 70%) as black solid. ^1H NMR (600 MHz, CD_2Cl_2) δ 8.72 (s, 1H), 8.58 (s, 1H), 7.89 (d, $J = 8.2$ Hz, 1H), 7.81 (d, $J = 8.3$ Hz, 1H), 7.52 (ddd, $J = 8.1, 6.8, 1.2$ Hz, 1H), 7.48 (ddd, $J = 8.2, 6.8, 1.1$ Hz, 1H), 7.29 (ddd, $J = 8.3, 6.8, 1.3$ Hz, 1H), 7.25 (ddd, $J = 8.3, 6.8, 1.3$ Hz, 1H), 6.93 (d, $J = 8.5$ Hz, 1H), 6.86 (d, $J = 8.5$ Hz, 1H), 6.30 (q, $J = 2.2$ Hz, 1H), 5.55 (t, $J = 2.7$ Hz, 1H), 5.49 (t, $J = 2.2$ Hz, 1H), 4.27 (d, $J = 14.7$ Hz, 1H), 4.03 (d, $J = 16.7$ Hz, 1H), 3.93 (d, $J = 14.7$ Hz, 1H), 3.25 (d, $J = 16.6$ Hz, 1H); ^{13}C NMR (151 MHz, CD_2Cl_2) δ 207.2, 140.9, 140.7, 138.1, 137.0, 135.7, 134.7, 134.6, 132.5, 132.0, 131.8, 128.1, 128.0, 127.8, 127.7, 127.7, 127.6, 127.1, 127.1, 104.3, 102.2, 101.9, 97.6, 97.0, 91.5, 83.5, 54.0, 38.3, 37.7; **IR (ATR):** $\nu_{\text{max}} = 3060, 2062, 1453, 760, 746, 710, 467$ cm^{-1} ; **HRMS (ESI)** calc'd. for $[\text{M}-\text{CO}-\text{I}]^+ = [\text{C}_{27}\text{H}_{17}\text{CoI}_3]^+$: 780.7791, found: 780.7775; $[\alpha]_{\text{D}}^{20} = +246.5$ ($c = 0.3$, CH_2Cl_2); **Mp** = 354 °C (decomposition).

Complex Co2:

Following the general procedure using (S)-**9** (50 mg, 137 μmol), $\text{Co}_2(\text{CO})_8$ (52 mg, 151 μmol) and iodine (42 mg, 165 μmol) gave complex **Co2** (56 mg, 79 μmol , 58%) as black solid. **^1H NMR** (400 MHz, CD_2Cl_2) δ 6.98 (d, J = 1.2 Hz, 2H), 6.85 (s, 2H), 5.79 (t, J = 2.1 Hz, 1H), 5.73 (t, J = 2.7 Hz, 1H), 5.41 (t, J = 2.2 Hz, 1H), 3.91 (s, 3H), 3.88 (d, J = 16.1 Hz, 1H), 3.77 (s, 3H), 3.74 (d, J = 13.8 Hz, 1H), 3.71 (s, 3H), 3.69 (s, 3H), 3.45 (d, J = 13.9 Hz, 1H), 2.65 (d, J = 16.0 Hz, 1H); **^{13}C NMR** (101 MHz, CD_2Cl_2) δ 152.6, 151.6, 151.5, 151.1, 128.8, 127.6, 126.1, 122.7, 111.5, 111.1, 111.0, 110.7, 105.3, 104.3, 95.1, 90.6, 86.4, 56.7 (2C), 56.3, 55.6, 26.9, 24.9; **IR (ATR):** ν_{max} = 2936, 2060, 1482, 1459, 1437, 1256, 1070, 1031 cm^{-1} ; **HRMS (ESI)** calc'd. for $[\text{M-CO-I}]^+ = [\text{C}_{23}\text{H}_{23}\text{O}_4\text{CoI}]^+$: 548.9968, found: 548.9970; **$[\alpha]_D^{20}$** = -149.5 (c = 0.29, CH_2Cl_2); **Mp** = 263 $^\circ\text{C}$ (decomposition).

General procedure for the synthesis of $[\text{Cp}^x\text{IrI}_2]_2$ complexes:

Following the reported procedure,^[7] the Cp^xH ligand (mixture of double bond isomers, 100 μmol , 1.00 eq.) was dissolved in benzene (1.00 mL, degassed by pump-freeze-thaw, 3 cycles) and thallium ethoxide (120 μmol , 1.20 eq.) was added as a solution in benzene (0.50 mL, degassed) under nitrogen atmosphere and in the absence of light at room temperature. The mixture was heated to 80 $^\circ\text{C}$ sealed tube for 2 h. In parallel, ethylene gas was bubbled through a suspension of $[\text{Ir}(\text{coe})_2\text{Cl}]_2$ (60.0 μmol , 0.60 eq.) in THF (1.50 mL) at 0 $^\circ\text{C}$ until a clear yellowish solution was obtained. This solution of $[\text{Ir}(\text{ethylene})_2\text{Cl}]_2$ was added to the above prepared reaction mixture at room temperature under nitrogen atmosphere and was stirred for 18 h. The obtained red solution containing a white precipitate was filtered over a short column of silica gel (wash with 50 mL toluene). After removal of the volatiles the corresponding $[\text{Cp}^x\text{Ir}(\text{ethylene})_2]$ complex was obtained as a yellowish gum/solid. The obtained $[\text{Cp}^x\text{Ir}(\text{ethylene})_2]$ complex (100 μmol , 1.00 eq.) was dissolved in toluene (0.50 mL) and a solution of iodine (105 μmol , 1.05 eq.) in toluene (1.00 mL) was added at 0 $^\circ\text{C}$ under nitrogen atmosphere. The orange reaction mixture was stirred in the dark for 45 min. Hexane (10.0 mL) was added and the dark precipitate was filtered off over a pipette filled with paper, washed with hexane (3 x 5 mL) and was afterwards dissolved in dichloromethane. After removal of the solvent the corresponding $[\text{Cp}^x\text{IrI}_2]_2$ was obtained as a red-brown, shining solid.

Complex Ir1:

Following the general procedure using (*R*)-**3b** (40 mg, 67 μ mol, 1.0 eq.), thallium ethoxide (20 mg, 81 μ mol, 1.2 eq.) and $[\text{Ir}(\text{coe})_2\text{Cl}]_2$ (36 mg, 40 μ mol, 0.60 eq.) gave the corresponding iridium(I) ethylene complex (29 mg, 34 μ mol, 51%) as a colorless gum. Oxidation with iodine (18 mg, 70 μ mol, 1.0 eq.) gave complex **Ir1** (30 mg, 14 μ mol, 43%) as a red shining solid. $^1\text{H NMR}$ (600 MHz, CDCl_3) δ 8.59 (s, 2H), 8.55 (s, 2H), 7.91 (d, J = 8.2 Hz, 2H), 7.78 (d, J = 8.2 Hz, 2H), 7.52 (t, J = 7.4 Hz, 2H), 7.44 (t, J = 7.2 Hz, 2H), 7.28 – 7.20 (m, 4H), 6.85 (dd, J = 14.7, 8.6 Hz, 4H), 5.89 (s, 2H), 5.73 (s, 2H), 5.49 (s, 2H), 3.81 (d, J = 14.7 Hz, 2H), 3.73 (d, J = 14.7 Hz, 2H), 3.67 (d, J = 16.1 Hz, 2H), 3.32 (d, J = 16.1 Hz, 2H); $^{13}\text{C NMR}$ (151 MHz, CDCl_3) δ 140.3, 140.1, 137.4, 136.5, 135.5, 134.1, 134.0, 131.9, 131.6, 131.5, 127.6, 127.5, 127.2, 127.2, 127.1, 126.9, 126.9, 101.5, 97.7, 93.0, 89.6, 84.6, 81.5, 69.0, 37.4, 36.8; **IR (ATR):** ν_{max} = 2923, 1558, 1444, 907, 883, 729 cm^{-1} ; **HRMS (APPI/LTQ-Orbitrap)** calc'd. for $[(\text{M-I})/2]^+ = [\text{C}_{27}\text{H}_{17}\text{I}_3\text{Ir}]^+$: 914.8088, found: 914.8074; $[\alpha]_{\text{D}}^{20} = -279.5$ (c = 0.2, CHCl_3); Mp = 250 $^{\circ}\text{C}$ (decomposition).

Complex Ir2:

Following the general procedure using (*S*)-**9** (35 mg, 96 μ mol, 1.0 eq.), thallium ethoxide (38 mg, 154 μ mol, 1.6 eq.) and $[\text{Ir}(\text{coe})_2\text{Cl}]_2$ (69 mg, 77 μ mol, 0.80 eq.) gave the corresponding iridium(I) ethylene complex (62 mg, 96 μ mol, 99%) as a colorless gum. Oxidation with iodine (27 mg, 106 μ mol, 1.1 eq.) gave complex **Ir2** (44 mg, 27 μ mol, 57%) as a purple shining solid. $^1\text{H NMR}$ (600 MHz, CDCl_3) δ 6.98 (d, J = 8.9 Hz, 2H), 6.83 (dd, J = 11.3 Hz, J = 2.4 Hz, 4H), 6.79 (d, J = 8.9 Hz, 2H), 5.70 (s, 2H), 5.54 (s, 2H), 4.95 (d, J = 2.4 Hz, 2H), 3.80 (s, 6H), 3.79 (d, J = 13.0 Hz, 2H), 3.73 (s, 6H), 3.70 (s, 6H), 3.64 (s, 6H), 3.57 (d, J = 15.3 Hz, 2H), 3.30 (d, J = 14.1 Hz, 2H), 2.78 (d, J = 15.4 Hz, 2H); $^{13}\text{C NMR}$ (151 MHz, CDCl_3) δ 152.3, 151.4, 151.2, 150.4, 128.5, 127.4, 126.6, 122.7, 110.5, 110.5, 110.3, 110.2, 97.8, 88.9, 83.2, 80.2, 68.0, 56.8, 56.6, 56.0, 55.3, 26.5, 23.9; **IR (ATR):** ν_{max} = 2933, 1483, 1256, 1072, 1032 cm^{-1} ; **HRMS (APPI/LTQ-Orbitrap)** m/z : $[\text{M}/2]^+$ Calc'd for $[\text{C}_{23}\text{H}_{23}\text{I}_2\text{IrO}_4]^+$ 809.9310; Found 809.9325; $[\alpha]_{\text{D}}^{20} = -23.5$ (c = 1.5, CHCl_3); Mp = 238 $^{\circ}\text{C}$.

Benchmark Rh^{III}-catalyzed reactions

General Procedure for the Enantioselective Rh-Catalyzed synthesis of Chiral Cyclopentenylamine **14**.^[8]

Without protection from oxygen or moisture, the mixture of Rh^I precatalyst (with (*R*)-enantiomer) (2.5 μ mol), and 2-methylbenzoic peroxyanhydride **13** (0.68 mg, 2.5 μ mol) in CF₃CH₂OH 0.5 mL) were stirred at room temperature. After stirring for 30 minutes, aryl ketoxime ether **11** (29 mg, 0.15 mmol) was added and followed by the diazabicycle **12** (18mg, 0.10 mmol). The mixture was stirred at 50°C. After 24 h, the solvent was removed under reduced pressure. The reaction mixture was purified by silica gel column chromatography (PE/EA: 10/1 to 5/1) to afford dibenzyl 1-((1*S*,2*R*)-2-(2-((*E*)-1-(butoxyimino)ethyl)phenyl)cyclopent-3-en-1-yl)hydrazine-1,2-dicarboxylate **14**. The analytical data was consistent with the literature. ¹H NMR (600 MHz, Acetonitrile-*d*₃, 70 °C) δ 7.39 – 7.25 (m, 11H), 7.23 – 7.11 (m, 3H), 5.83 (m, 1H), 5.64 – 5.57 (m, 1H), 5.15 (m, 2H), 4.99 (s, 2H), 4.94 – 4.83 (m, 1H), 4.35 (s, 1H), 4.13 – 4.05 (m, 2H), 2.69 (br, 1H), 2.59 (dd, *J* = 14.6, 4.7 Hz, 1H), 1.65 – 1.55 (m, 2H), 1.36 (dq, *J* = 13.8, 6.8 Hz, 2H), 0.93 – 0.88 (m, 3H).

HPLC trace of *Rac*-**14**:

Figure S5: HPLC trace of *Rac*-**14**

With complex Rh1. Desired product (6.5mg, 12 μ mol, 23%) was obtained as a colorless oil in 17:83 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 254 nm; t_R (minor) = 5.1 min, t_R (major) = 12.3 min, 17 : 83 er.)

Figure S6: HPLC trace of **14** prepared with **Rh1** as catalyst

With complex Rh2. Desired product (3.6mg, 6.5 μ mol, 13%) was obtained as a colorless oil in 78:22 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 254 nm; t_R (major) = 5.1 min, t_R (minor) = 12.7 min, 78 : 22 er.)

Figure S7: HPLC trace of **14** prepared with **Rh2** as catalyst

With complex Rh3. Desired product (21.0mg, 38 μ mol, 76%) was obtained as a colorless oil in 95:5 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 254 nm; t_R (major) = 5.0 min, t_R (minor) = 12.2 min, 95 : 5 er.)

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.028	BB	0.1169	1915.46997	251.36276	94.7181
2	12.242	BB	0.3268	106.81448	4.08813	5.2819

Figure S8: HPLC trace of **14** prepared with **Rh3** as catalyst

With complex Rh4. Desired product (27.5mg, 49 μ mol, 99%) was obtained as a colorless oil in 5:95 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 254 nm; t_R (minor) = 5.1 min, t_R (major) = 12.4 min, 5 : 95 er.)

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	5.090	BV	0.1221	231.98605	28.16319	5.3058
2	12.417	BB	0.4266	4140.31348	149.74406	94.6942

Figure S9: HPLC trace of **14** prepared with **Rh4** as catalyst

With complex Rh6. Desired product (4.0mg, 7.2 μ mol, 14%) was obtained as a slightly yellow oil in 95:5 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 254 nm; t_R (major) = 5.0 min, t_R (minor) = 12.0 min, 55 : 45 er.)

Figure S10: HPLC trace of **14** prepared with **Rh6** as catalyst

General Procedure for the Enantioselective Rh-Catalyzed C-H Allylations of Benzamides.^[9]

Without protection from oxygen and moisture, Rh^I precatalyst (with (*R*)-enantiomer) (5.00 μ mol), dibenzoylperoxide (1.20 mg, 5.00 μ mol) and **15** (15.0 mg, 0.10 mmol, 1.0 equiv.) were weighed into a vial equipped with a magnetic stir bar and sealed with a rubber septum. 0.5 mL of CH₂Cl₂ was added and the mixture was stirred at 23°C for 15 mins. After cooling down to -20°C, the allene **16** (20.0mg, 0.12 mmol, 1.2 equiv.) was added then the reaction mixture was kept at -20°C for 18 hours. The mixture was quickly filtered on a pad of silica gel, eluting with EtOAc, and then concentrated under reduced pressure. The residue was purified further by silica gel column (hexane:EtOAc 4:1 to 3:2), yielding (*R*)-2-(1-cyclohexylidene-4-methoxybutan-2-yl)-N-methoxybenzamide **17**. The analytical data was consistent with the literature. ¹H NMR (400 MHz, CDCl₃) δ 9.65 (s, 1H), 7.48 – 7.37 (m, 2H), 7.30 (d, *J* = 7.6 Hz, 1H), 7.22 (td, *J* = 7.5, 1.3 Hz, 1H), 5.15 (d, *J* = 9.3 Hz, 1H), 4.23 – 4.10 (m, 1H), 3.90 (s, 3H), 3.35 (dt, *J* = 10.3, 5.3 Hz, 1H), 3.28 (s, 3H), 3.23 (td, *J* = 9.4, 5.1 Hz, 1H), 2.22 (dd, *J* = 18.3, 5.0 Hz, 1H), 2.15 – 2.06 (m, 1H), 2.01 (dq, *J* = 14.8, 8.0, 6.5 Hz, 3H), 1.90 (ddt, *J* = 14.0, 9.4, 4.9 Hz, 1H), 1.56 – 1.39 (m, 6H);

Figure S11: HPLC trace of Rac-17

With complex Rh1. Desired product (18.5mg, 58 μ mol, 58%) was obtained as a slightly yellow oil in 93:7 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 220 nm; t_R (major) = 6.3 min, t_R (minor) = 7.3 min, 93 : 7 er.)

Figure S12: HPLC trace of 17 prepared with Rh1 as catalyst

With complex Rh2. Desired product (26.7mg, 84 μ mol, 84%) was obtained as a colorless oil in 2:98 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 220 nm; t_R (minor) = 6.4 min, t_R (major) = 7.3 min, 2 : 98 er.)

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.432	MM	0.2254	10.77831	7.96837e-1	1.9785
2	7.345	BB	0.2313	533.98169	33.98544	98.0215

Figure S13: HPLC trace of **17** prepared with **Rh2** as catalyst

With complex Rh3. Desired product (27.1mg, 85 μ mol, 85%) was obtained as a colorless oil in 6:94 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 220 nm; t_R (minor) = 6.4 min, t_R (major) = 7.2 min, 6 : 94 er.)

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.366	BB	0.2191	59.56707	3.96942	6.0647
2	7.171	BB	0.2383	922.62793	56.56597	93.9353

Figure S14: HPLC trace of **17** prepared with **Rh3** as catalyst

With complex Rh4. Desired product (27.3mg, 86 μ mol, 86%) was obtained as a colorless oil in 90:10 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 220 nm; t_R (major) = 6.4 min, t_R (minor) = 7.5 min, 90 : 10 er.)

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.378	BB	0.2058	687.20880	48.93403	90.0689
2	7.463	BB	0.2203	75.77209	5.01541	9.9311

Figure S15: HPLC trace of **17** prepared with **Rh4** as catalyst

With complex Rh5. Desired product (25.0mg, 79 μ mol, 79%) was obtained as a colorless oil in 11:89 er. **Chiral HPLC:** (Chiralpak IB; hexane:i-PrOH 80:20, 1.0 mL/min, 220 nm; t_R (minor) = 6.4 min, t_R (major) = 7.3 min, 11 : 89 er.)

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	6.367	BV	0.2415	528.56543	31.86182	10.6354
2	7.300	MM	0.3462	4441.30078	213.83522	89.3646

Figure S16: HPLC trace of **17** prepared with **Rh5** as catalyst

Single crystal X-ray analysis

Crystallographic data for the structures reported in this paper have been deposited at the Cambridge Crystallographic Data Center (CCDC) as Supplementary Publication No. CCDC1918423, CCDC1919209-1919210 and CCDC191912-191918. Copies of the data can be obtained free of charge on application to the CCDC via <https://www.ccdc.cam.ac.uk/structures>.

Crystallization methods:

- X-Ray quality crystals of **Rh1** ((*R*)-enantiomer) were obtained by slow vapor diffusion of MeOH in a solution of the complex (5 mg) in Pentane (0.1 ml) at room temperature for two days.
- X-Ray quality crystals of **Rh2** were obtained by slow vapor diffusion of Pentane in a solution of the complex (5 mg) in chloroform (0.2 ml) at room temperature for several days
- X-Ray quality crystals of **Rh3** were obtained by slow vapor diffusion of Pentane in a solution of the complex (5 mg) in chloroform (0.2 ml) at room temperature for several days
- X-Ray quality crystals of **Rh4** were obtained by slow vapor diffusion of Pentane in a solution of the complex (5 mg) in chloroform (0.2 ml) at room temperature for several days.
- X-Ray quality crystals of **Rh6** were obtained by slow vapor diffusion of EtOH in a solution of the complex (5 mg) in Hexane (0.2 ml) at room temperature for several days.
- X-Ray quality crystals of **Co1** were obtained by slow diffusion from a solution of the complex (5 mg) in chloroform (0.3 ml) layered with few drops of hexane at room temperature for 4 days.
- X-Ray quality crystals of **Co2** were obtained by slow vapor diffusion of Et₂O from a solution of the complex (5 mg) in DCM (0.1 ml) at room temperature for two days.
- X-Ray quality crystals of **Ir1** were obtained by slow diffusion from a solution of the complex (10 mg) in dichloromethane (0.3 ml) layered with few drops of diethyl ether at room temperature for 4 days.
- X-Ray quality crystals of **Ir2** were obtained by slow diffusion from a solution of the complex (10 mg) in chloroform (0.3 ml) layered with few drops of diethyl ether at room temperature for 4 days.

NMR spectra

Figure S17: (S)- 3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-dicarboxylic acid, ¹H NMR

Figure S18: (S)- 3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-dicarboxylic acid, ¹³C NMR

Figure S19: (S)-(3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-diyl)dimethanol, ^1H NMR

Figure S20: (S)-(3,3'-bis(trimethylsilyl)-[1,1'-binaphthalene]-2,2'-diyl)dimethanol, ^{13}C NMR

Figure S21: (S)-2a, ^1H NMR

Figure S22: (S)-2a, ^{13}C NMR

Figure S25: (R)-diisopropyl 3,3'-diiodo-[1,1'-binaphthalene]-2,2'-dicarboxylate, ^1H NMR

Figure S26: (R)-diisopropyl 3,3'-diiodo-[1,1'-binaphthalene]-2,2'-dicarboxylate, ^{13}C NMR

Figure S27: (R)-(3,3'-diiodo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol, ^1H NMR

Figure S28: (R)-(3,3'-diiodo-[1,1'-binaphthalene]-2,2'-diyl)dimethanol, ^{13}C NMR

Figure S31: (R)-3b, ^1H NMR

Figure S32: (R)-3b, ^{13}C NMR

Figure S33: (R)-diisopropyl 3,3'-dibromo-[1,1'-binaphthalene]-2,2'-dicarboxylate, ^1H NMR

Figure S34: (R)-diisopropyl 3,3'-dibromo-[1,1'-binaphthalene]-2,2'-dicarboxylate, ^{13}C NMR

Figure S35: (R)-(3,3'-dibromo-[1,1'- binaphthalene]-2,2'-diyl)dimethanol, ¹H NMR

Figure S36: (R)-(3,3'-dibromo-[1,1'- binaphthalene]-2,2'-diyl)dimethanol, ¹³C NMR

Figure S37: (R) -**2c**, ^1H NMR

Figure S38: (R) -**2c**, ^{13}C NMR

Figure S39: (R)-3c, ¹H NMR

Figure S40: (R)-3c, ¹³C NMR

Figure S41: 2-iodo-3,6-dimethoxybenzoic acid, ¹H NMR

Figure S42: 2-iodo-3,6-dimethoxybenzoic acid, ¹³C NMR

Figure S43: **6**, ^1H NMR

Figure S44: **6**, ^{13}C NMR

Figure S45: (3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-diyl)dimethanol, ^1H NMR

Figure S46: (3,3',6,6'-tetramethoxy-[1,1'-biphenyl]-2,2'-diyl)dimethanol, ^{13}C NMR

Figure 47: **8**, ^1H NMR

Figure S48: **8**, ^{13}C NMR

Figure S49: (*Rac*)-**9**, ^1H NMR

Figure S50: (*Rac*)-**9**, ^{13}C NMR

Figure S53: (R)-9, ^1H NMR

Figure S54: (S)-9, ^1H NMR

Figure S61: **Rh4**, ^1H NMR

Figure S62: **Rh4**, ^{13}C NMR

Figure S63: **Co1**, ¹H NMR

Figure S64: **Co1**, ¹³C NMR

— 197.59

S59

Figure S67: Ir1, ¹H NMR

Figure S68: Ir1, ¹³C NMR

Figure S69: **Ir2**, ^1H NMR

Figure S70: **Ir2**, ^{13}C NMR

Figure S71: **14**, ¹H NMR

Figure S72: **17**, ¹H NMR

References

- (1) Egami, H.; Sato, K.; Asada, J.; Kawato, Y.; Hamashima, Y., Concise synthesis of binaphthol-derived chiral dicarboxylic acids. *Tetrahedron* **2015**, *71* (37), 6384-6388.
- (2) Ooi, T.; Uematsu, Y.; Maruoka, K., New, Improved Procedure for the Synthesis of Structurally Diverse N-Spiro C2-Symmetric Chiral Quaternary Ammonium Bromides. *J. Org. Chem.* **2003**, *68* (11), 4576-4578.
- (3) Widhalm, M.; Abraham, M.; Arion, V. B.; Saarsalu, S.; Maeorg, U., A modular approach to a new class of phosphinohydrazones and their use in asymmetric allylic alkylation reactions. *Tetrahedron: Asymmetry* **2010**, *21* (16), 1971-1982.
- (4) Ooi, T.; Uematsu, Y.; Maruoka, K., Asymmetric Strecker Reaction of Aldimines Using Aqueous Potassium Cyanide by Phase-Transfer Catalysis of Chiral Quaternary Ammonium Salts with a Tetranaphthyl Backbone. *J. Am. Chem. Soc.* **2006**, *128* (8), 2548-2549.
- (5) Audic, B.; Wodrich, M. D.; Cramer, N., Mild complexation protocol for chiral Cp^xRh and Ir complexes suitable for in situ catalysis. *Chem. Sci.* **2019**, *10* (3), 781-787.
- (6) Ozols, K.; Jang, Y.-S.; Cramer, N., Chiral Cyclopentadienyl Cobalt(III) Complexes Enable Highly Enantioselective 3d-Metal-Catalyzed C–H Functionalizations. *J. Am. Chem. Soc.* **2019**, *141* (14), 5675-5680.
- (7) Dieckmann, M.; Jang, Y.-S.; Cramer, N., Chiral Cyclopentadienyl Iridium(III) Complexes Promote Enantioselective Cycloisomerizations Giving Fused Cyclopropanes. *Angew. Chem. Int. Ed.* **2015**, *54* (41), 12149-12152.
- (8) Wang, S.-G.; Cramer, N., An Enantioselective Cp^xRh(III)-Catalyzed C–H Functionalization/Ring-Opening Route to Chiral Cyclopentenylamines. *Angew. Chem. Int. Ed.* **2019**, *58* (8), 2514-2518.
- (9) Ye, B.; Cramer, N., A Tunable Class of Chiral Cp Ligands for Enantioselective Rhodium(III)-Catalyzed C–H Allylations of Benzamides. *J. Am. Chem. Soc.* **2013**, *135* (2), 636-639.