

DYNAMICAL COLLAPSE OF CYLINDRICAL SYMMETRIC DIPOLAR BOSE-EINSTEIN CONDENSATES

JACOPO BELLAZZINI AND LUIGI FORCELLA

ABSTRACT. We study the formation of singularities for cylindrical symmetric solutions to the Gross-Pitaevskii equation describing a Dipolar Bose-Einstein condensate. We prove that solutions arising from initial data with energy below the energy of the Ground State and that do not scatter collapse in finite time. The main tools to prove our result is a crucial localization property for the fourth power of the Riesz transforms, that we prove by means of the decay properties of the heat kernel associated to the parabolic biharmonic equation, and pointwise estimates for the square of the Riesz transforms. Furthermore, other essential tools are the variational characterization of the Ground State energy, and suitable localized virial identities for cylindrical symmetric functions.

1. INTRODUCTION

Since the first experimental observation in 1995 of a quantum state of matter at very low temperature called Bose-Einstein condensate (BEC), see e.g. [1, 7, 11], the study of the asymptotic dynamics of nonlinear equations describing this phenomena rapidly increased, both numerically and theoretically. Since BEC exists in an ultracold and dilute regime, the most relevant interactions are the isotropic, elastic two-body collisions. After the first pioneering experimental works, other condensates have been produced with different atoms, in particular condensates made out of particles possessing a permanent electric or magnetic dipole moment. Such kind of condensates are called Dipolar Bose-Einstein condensates, see e.g. [3, 4, 24, 26, 28], and their peculiarity is given by the long-range anisotropic interaction between particles, in contrast with the short-range, isotropic character of the contact interaction of BEC.

A Dipolar quantum gases is well modelled, see [21, 29, 30] for the validity of such model, by the Gross-Pitaevskii equation (GPE)

$$ih \frac{\partial u}{\partial t} = -\frac{h^2}{2m} \nabla^2 u + W(x)u + U_0 |u|^2 u + (V_{dip} * |u|^2)u, \quad (1.1)$$

where the wave function $u = u(t, x)$. Here t is the time variable, $x = (x_1, x_2, x_3)$ is the space coordinate, h is the Planck constant, m is the mass of a dipolar particle and $W(x)$ is an external, real potential which describes the electromagnetic trap. The coefficient $U_0 = 4\pi h^2 a_s / m$ describes the local interaction between dipoles in the condensate, a_s being the s -wave scattering length

2000 *Mathematics Subject Classification.* 35Q55, 35B40, 82C10, 35J20.

Key words and phrases. Gross-Pitaevskii equation, BEC, NLS-like equation, finite time blow-up, singular integral estimates.

(positive for repulsive interactions and negative for attractive interactions). The long-range dipolar interaction potential between two dipoles is given by

$$V_{dip}(x) = \frac{\mu_0 \mu_{dip}^2}{4\pi} \frac{1 - 3 \cos^2(\theta)}{|x|^3}, \quad x \in \mathbb{R}^3,$$

where μ_0 is the vacuum magnetic permeability, μ_{dip} is the permanent magnetic dipole moment and θ is the angle between the dipole axis $n \in \mathbb{R}^3$ and the vector x . For simplicity, we fix the dipole axis as the vector $n = (0, 0, 1)$. The wave function is normalized according to

$$\int_{\mathbb{R}^3} |u(x, t)|^2 dx = N,$$

where N is the total number of dipolar particles in the Dipolar BEC. In this work we consider the case when the trapping potential W is not active, i.e. we freeze $W(x) = 0$.

As we are interested in the mathematical features of the GPE, we consider it in its dimensionless form, therefore we write the Cauchy problem associated to (1.1) as follows:

$$\begin{cases} i\partial_t u + \frac{1}{2}\Delta u = \lambda_1 |u|^2 u + \lambda_2 (K * |u|^2) u, & (t, x) \in \mathbb{R} \times \mathbb{R}^3 \\ u(0, x) = u_0(x) \end{cases}, \quad (1.2)$$

where the dipolar kernel K acting in the convolution on the mass density $|u|^2$ is given by

$$K(x) = \frac{x_1^2 + x_2^2 - 2x_3^2}{|x|^5}.$$

The two coefficients $\lambda_{1,2}$ involved in the equation are two physical, real parameters defined by

$$\lambda_1 = 4\pi a_s N \gamma, \quad \lambda_2 = \frac{mN\mu_0\mu_{dip}^2}{4\pi h^2} \gamma;$$

they describe the strength of the local nonlinearity and the nonlocal nonlinearity, respectively. Following the terminology introduced by Carles, Markowich and Sparber in [8], where the authors give a first mathematical treatment concerning various aspects about local/global well-posedness of (1.2), we consider the partition of the coordinate plane (λ_1, λ_2) into the so-called *Unstable Regime*

$$\begin{cases} \lambda_1 - \frac{4\pi}{3}\lambda_2 < 0 & \text{if } \lambda_2 > 0 \\ \lambda_1 + \frac{8\pi}{3}\lambda_2 < 0 & \text{if } \lambda_2 < 0 \end{cases}, \quad (1.3)$$

and its complementary, the so-called *Stable Regime*

$$\begin{cases} \lambda_1 - \frac{4\pi}{3}\lambda_2 \geq 0 & \text{if } \lambda_2 > 0 \\ \lambda_1 + \frac{8\pi}{3}\lambda_2 \geq 0 & \text{if } \lambda_2 < 0 \end{cases}. \quad (1.4)$$

Heuristically, when comparing (1.2) to the classical cubic NLS (i.e. when $\lambda_2 = 0$), one can think to the configuration given by (1.3) as the nonlinearity were focusing, and to (1.4) as the nonlinearity were defocusing. This notation is although incorrect in the context of the GPE, as we will emphasize in some remark later on in the paper, after we introduce some basic notation.

Solutions to (1.2) conserve along the flow the mass and the energy (besides other quantities not used in this paper); more rigorously

$$M(t) = M(u(t)) := \int_{\mathbb{R}^3} |u(t)|^2 dx = M(0)$$

and

$$E(t) = E(u(t)) := \frac{1}{2} \left(\int_{\mathbb{R}^3} |\nabla u(t)|^2 + \lambda_1 |u(t)|^4 + \lambda_2 (K * |u(t)|^2) |u(t)|^2 dx \right) = E(0), \quad (1.5)$$

for any $t \in (-T_{min}, T_{max})$, where $T_{min}, T_{max} \in (0, \infty]$ are the minimal and maximal time of existence of the solution, respectively. Local existence of solutions to (1.2) was shown in [8], in both the configurations given by (1.3) and (1.4).

The Unstable Regime (1.3) is of particular relevance, since stationary solutions are allowed in this region. More precisely, we recall that stationary states are solutions of the following species:

$$u(x, t) = e^{-i\kappa t} u(x),$$

where $u(x)$ is a time-independent function solving the stationary equation

$$-\frac{1}{2} \Delta u + \lambda_1 |u|^2 u + \lambda_2 (K * |u|^2) u + \kappa u = 0 \quad (1.6)$$

constrained on the manifold $S(1)$, where

$$S(1) = \{u \in H^1(\mathbb{R}^3) \text{ s.t. } \|u\|_{L^2(\mathbb{R}^3)}^2 = 1\}, \quad (1.7)$$

and $\kappa \in \mathbb{R}$ is a real parameter usually referred as the chemical potential. We postpone the rigorous discussion about existence of solutions to (1.6) in Section 3. We introduce now some crucial quantities often used along the paper, and we proceed enunciating the main results of this work and the strategy to get them.

Let us recall some notation consistent to our previous papers [5, 6]: by means of the Plancherel identity, the energy defined in (1.5) can be rewritten as

$$E(t) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u|^2 dx + \frac{1}{2(2\pi)^3} \int_{\mathbb{R}^3} \left(\lambda_1 + \lambda_2 \hat{K}(\xi) \right) (\widehat{|u|^2})^2(\xi) d\xi$$

where the Fourier transform of the dipolar kernel K is explicitly given by

$$\hat{K}(\xi) = \frac{4\pi}{3} \frac{2\xi_3^2 - \xi_2^2 - \xi_1^2}{|\xi|^2}, \quad \xi \in \mathbb{R}^3. \quad (1.8)$$

We refer to [8] for a proof of the explicit calculation of \hat{K} , done by means of the decomposition in spherical harmonics of the Fourier character $e^{-ix \cdot \xi}$.

Remark 1.1. It is worth mentioning right now that, by using (1.8), $\hat{K}(\xi)$ is a linear combination of symbols associated to the square of the Riesz transforms \mathcal{R}_j^2 for $j = 1, 2, 3$, i.e. $\widehat{\mathcal{R}_j^2 f}(\xi) = -\frac{\xi_j^2}{|\xi|^2} \hat{f}(\xi)$. Therefore $(K * f)(x)$ is a linear combination of $\mathcal{R}_j^2 f(x)$'s.

A trivial computation provides a lower and an upper bound for \hat{K} , and more precisely

$$\hat{K} \in \left[-\frac{4}{3}\pi, \frac{8}{3}\pi \right],$$

and from the latter it is straightforward to claim that the convolution with K defines an $L^2 \mapsto L^2$ continuous operator.

We split the energy as sum of the kinetic and potential energies, respectively defined by

$$T(u) = \int_{\mathbb{R}^3} |\nabla u|^2 dx \quad (1.9)$$

and

$$P(u) = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} \left(\lambda_1 + \lambda_2 \hat{K}(\xi) \right) (|\widehat{u}|^2)^2(\xi) d\xi, \quad (1.10)$$

and we introduce the quantity

$$G(u) = T(u) + \frac{3}{2}P(u). \quad (1.11)$$

Moreover, the following useful identity holds true: $E - \frac{1}{3}G = \frac{1}{6}T$. The functional G naturally appears by means of the Pohozaev identities related to (1.6), see [2].

In spite of the fact that we are primarily interested in solutions satisfying (1.7), we consider, for a positive $c > 0$, the generic manifold

$$S(c) = \left\{ u \in H^1(\mathbb{R}^3) \text{ s.t. } \|u\|_{L^2(\mathbb{R}^3)}^2 = c \right\},$$

which will be useful for the mathematical study of existence of standing states, and their variational characterization. The case $c = 1$ clearly corresponds to the mass normalization expressed in (1.7). For a fixed $c > 0$, the energy $E(u)$ has a mountain pass geometry on $S(c)$ and we denote by $\gamma(c)$ the mountain pass energy at level c to which it corresponds a stationary state. Again, we refer to [Section 3](#) for precise definitions and rigorous results.

The energy level $\gamma(c)$ has the variational characterization below, that will be essential in the sequel; by introducing the manifold

$$V(c) = \{ u \in H^1(\mathbb{R}^3) \text{ s.t. } \|u\|_{L^2(\mathbb{R}^3)}^2 = c \text{ and } G(u) = 0 \}$$

we recall, see [6], that

$$\gamma(c) = \inf \{ E(u) \text{ s.t. } u \in V(u) \}. \quad (1.12)$$

A contradiction argument in conjunction with a continuity argument implies that provided $E(u_0) < \gamma(c)$, with $c = \|u_0\|_{L^2(\mathbb{R}^3)}^2$ and $G(u_0) > 0$, the local solution $u \in \mathcal{C}((-T_{min}, T_{max}); H^1(\mathbb{R}^3))$ to (1.2) can be extended globally in time, i.e. $T_{min} = T_{max} = \infty$, and $G(u(t)) > 0$ for any $t \in \mathbb{R}$,

see [6, Theorem 1.3].

The global existence of solutions under the hypothesis $E(u_0) < \gamma(c)$, with $c = \|u_0\|_{L^2(\mathbb{R}^3)}^2$, and $G(u_0) > 0$, suggests the possibility that all solutions arising from these initial data scatter, in analogy of what was proved by Duyckaerts, Holmer, and Roudenko in [13, 18] for the cubic focusing NLS by exploiting the original approach of concentration/compactness and rigidity method in the spirit of the Kenig and Merle road map, see [19]. The authors in fact recently proved in [5] that the conditions $E(u_0) < \gamma(c)$, for $c = \|u_0\|_{L^2(\mathbb{R}^3)}^2$, and $G(u_0) > 0$ imply scattering of solutions to the Dipolar Gross-Pitaevskii equation (1.2).

The main aim of this paper is to study the asymptotic dynamics in the complementary configuration, i.e. $E(u_0) < \gamma(c)$, with $c = \|u_0\|_{L^2(\mathbb{R}^3)}^2$, and $G(u_0) < 0$. We shall underline that even in the case when $\lambda_2 = 0$ and $\lambda_1 < 0$, namely when (1.2) reduces to the classical focusing cubic nonlinear Schrödinger equation

$$\begin{cases} i\partial_t u + \frac{1}{2}\Delta u = \lambda_1 |u|^2 u, & (t, x) \in \mathbb{R} \times \mathbb{R}^3 \\ u(0, x) = u_0(x) \end{cases}, \quad (1.13)$$

finite time blow-up for any initial data $u_0 \in H^1(\mathbb{R}^3)$ satisfying the above conditions is still an open problem. To the best of our knowledge, the less restrictive assumptions in this context are due to Martel, see [23], where the author proves finite time blow-up in the space of cylindrical symmetric functions with finite variance in the x_3 direction and negative energy. Early results of this type are due to Glassey, see [16], in case of finite variance, and Ogawa and Tsutsumi [25], in the radial symmetric case (see also Holmer and Roudenko [18]).

We are now in position to state our main results and to explain our strategy of the proofs. Let us define $\bar{x} = (x_1, x_2)$ and let us introduce the space where we study the formation of singularities:

$$\Sigma_3 = \left\{ u \in H^1(\mathbb{R}^3) \quad s.t. \quad u(x) = u(|\bar{x}|, x_3) \quad \text{and} \quad x_3 u \in L^2(\mathbb{R}^3) \right\};$$

Σ_3 is therefore the space of cylindrical symmetric functions with finite variance in the x_3 direction.

Our main result is as follows.

Theorem 1.2. *Let $u(t) \in \Sigma_3$ be a solution to (1.2) defined on $(-T_{min}, T_{max})$, with initial datum u_0 satisfying $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$ and $G(u_0) < 0$. Then T_{min} and T_{max} are finite, namely $u(t)$ blows-up in finite time.*

As a consequence of Theorem 1.2 we give a generalization of the result by Martel in [23]. We extend here that result for all positive initial energies under the energy threshold given by the Ground State associated to NLS (which corresponds to the one given in (1.6) for $\lambda_2 = 0$). Even if the following theorem can be viewed as a straightforward corollary of Theorem 1.2, we prefer to state it as an independent result, since it has its own interest.

Theorem 1.3. *Given a solution $u(t) \in \Sigma_3$ to (1.13) with $\lambda_1 < 0$ defined on $(-T_{min}, T_{max})$, with initial datum u_0 satisfying $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$ and $G(u_0) < 0$, then T_{min} and T_{max} are finite, namely $u(t)$ blows-up in finite time.*

We point out some feature of the Dipolar GPE.

Remark 1.4. Blow-up in finite time for the focusing cubic NLS in the whole generality, i.e. for infinite-variance initial data and without assuming any symmetry, is still an open problem. See the beginning of [Section 4](#) for up-to-date references.

Remark 1.5. An usual assumption that often appears in literature in order to simplify the analysis of a model which cannot be treated in a full generality, is the restriction to a radial setting. The dipolar kernel $K(x)$ is a Calderón-Zigmund operator of the form $|x|^{-3}\mathcal{O}(x)$ where \mathcal{O} is a zero-order function having zero average on the sphere. This implies that the restriction to radial symmetric solutions makes disappear the effect of the nonlocal term in (1.2), hence the equation reduces to the classical NLS equation in the radial framework, see [\[8\]](#).

Remark 1.6. We restrict the functional space to functions belonging to $H^1(\mathbb{R}^3)$ with finite variance in the x_3 direction. As it will be clear along the proof a decay estimate for the potential energy in the exterior of a cylindrical domain will be crucial. Indeed, the finite variance in the x_3 direction enables us to localize the potential energy on the exterior of a cylinder.

Remark 1.7. As remarked in [\[8\]](#), for $0 < \lambda_1 < \frac{4\pi}{3}\lambda_2$ – namely when the local nonlinearity is defocusing, and the coupling parameter λ_2 is positive, which is the physical case – finite time blow-up may arise, so that is improper to speak about “defocusing/focusing” for the Dipolar BEC. The nonlocal interaction then can yield to formation of singularities in finite time of the solutions.

Remark 1.8. From the identity (1.11) and the fact that there exists a positive constant $\delta > 0$ such that $G(u(t)) \leq -\delta$ for any $t \in (-T_{min}, T_{max})$ (see [Lemma 3.5](#) below), it is straightforward to see that the assumption $G(u_0) < 0$ implies that $P(u(t)) < 0$ for any time in the maximal interval of existence of the solution to (1.2). This is in contrast of what happens in the counterpart scenario $G(u_0) > 0$. In the latter case, working in the Unstable Regime (1.3) does not guarantee that the potential energy $P(u(t))$ preserves the sign, as we proved in [\[5\]](#).

We turn now to state the ingredients we use in order to prove our main theorems. The strategy and the main difficulties are the following.

- A variational characterization of the Ground State energy firstly permits to prove that $\|u(x, t)\|_{\dot{H}^1(\mathbb{R}^3)}$ is bounded from below uniformly in time and that $G(u(x, t)) < -\delta$ for all times in the maximal interval of existence of the solution. As a byproduct, which is crucial for what follows, it exploits the bound $G(u(x, t)) \leq -\tilde{\delta}\|u(x, t)\|_{\dot{H}^1(\mathbb{R}^3)}^2$, for some $\tilde{\delta} > 0$.
- The virial identities, valid both for (1.2) and (1.13). We define, following [Martel \[23\]](#),

$$V_\rho(t) = 2 \int_{\mathbb{R}^3} \rho_R(x) |u(t, x)|^2 dx,$$

where ρ , which is in particular a well-constructed function depending only on the two variables $\bar{x} = (x_1, x_2)$ which provides a localization in the exterior of a cylinder, parallel to the x_3 axis and with radius of size $|\bar{x}| \sim R$. Here $|\bar{x}|$ clearly denotes $|\bar{x}| := (x_1^2 + x_2^2)^{1/2}$. Moreover we consider the not-localized function x_3^2 in order to obtain a virial-like estimate of the form

$$\frac{d^2}{dt^2} V_{\rho_R + x_3^2}(t) \leq 4 \underbrace{\int_{\mathbb{R}^3} |\nabla u|^2 dx + 6\lambda_1 \int_{\mathbb{R}^3} |u|^4 dx}_{:=\mathcal{G}} + cR^{-2} + H_R(u(t)), \quad (1.14)$$

where the error H_R is defined by

$$\begin{aligned} H_R(u(t)) &= 4\lambda_1 \int_{\mathbb{R}^3} F_R(\bar{x}) |u|^4 dx + 2\lambda_2 \int_{\mathbb{R}^3} \nabla \rho_R \cdot \nabla (K * |u|^2) |u|^2 dx \\ &\quad - 4\lambda_2 \int_{\mathbb{R}^3} x_3 \partial_{x_3} (K * |u|^2) |u|^2 dx \end{aligned}$$

and $F_R(\bar{x})$ is a nonnegative function supported in the exterior of a cylinder of radius of order R .

- In the case $\lambda_1 < 0$, $\lambda_2 = 0$ (namely NLS), the decay property of the L^4 -norm of a function supported outside a cylinder of radius of order R , more precisely the estimate $\|u\|_{L^4(|\bar{x}| \geq R)}^4 \lesssim R^{-1} \|u\|_{\dot{H}^1(\mathbb{R}^3)}^2$, together with the localized virial identities, implies finite time blow-up by a convexity argument. We underline that everything works well since we are able to prove that $\|u\|_{\dot{H}^1(\mathbb{R}^3)}^2$ controls either $G(u)$ or the remainder term H_R . Note that in this case the quantity \mathcal{G} in the r.h.s. of (1.14) precisely defines $4G(u)$ in the context of (1.13).
- When $\lambda_2 \neq 0$ we have to deal with the effect of the dipolar interaction term - incorporated in H_R in (1.14) - that is nonlocal and that is neither always positive nor always negative. As already pointed out, see Remark 1.1, $K * \cdot$ acts as the sum (up to some constant coefficients) of square of Riesz transforms. Our strategy is to split u by separating it in the interior and in the exterior of a cylinder, namely $u = u_i + u_o$ where

$$u_i = \mathbf{1}_{\{|\bar{x}| \leq CR\}} u \quad \text{and} \quad u_o = \mathbf{1}_{\{|\bar{x}| \geq CR\}} u,$$

and computing the interaction given by the dipolar term. Here $\mathbf{1}$ denotes the indicator function on a measurable set. The problem here is that $K * |u_i|^2$ is not supported inside any cylinder. A crucial tool is given by the pointwise estimate

$$|\mathbf{1}_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \mathcal{R}_j^2[(1 - \mathbf{1}_{\{|\bar{x}| \leq \gamma_2 R\}})f](x)| \leq CR^{-3} \mathbf{1}_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \|f\|_{L^1(|\bar{x}| \geq \gamma_2 R)}$$

when $\gamma_2 - \gamma_1 > 0$, see Lemma 2.5.

- In order to control the remainder term and to generate the quantity $6\lambda_2 \int_{\mathbb{R}^3} (K * |u|^2) |u|^2 dx$ in (1.14) that will yield to the whole quantity $4G(u)$, see (1.11), we need to use the identity $2 \int_{\mathbb{R}^3} x \cdot \nabla (K * f) f dx = -3 \int_{\mathbb{R}^3} (K * f) f dx$ which follows from the relation $\xi \cdot \nabla_\xi \hat{K} = 0$. The difficulty here comes from the fact that the localization function used in the virial identities $\rho_R(\bar{x})$ satisfies

$$\rho_R(\bar{x}) = \begin{cases} |\bar{x}|^2 & \text{if } |\bar{x}| < R, \\ \text{constant} & \text{if } |\bar{x}| > 2R \end{cases},$$

while the function $\rho = x_3^2$ is unbounded. By observing that

$$\begin{aligned} & 2 \int_{\mathbb{R}^3} x_3 \partial_{x_3} \left(K * |u_i|^2 \right) |u_o|^2 dx + 2 \int_{\mathbb{R}^3} x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_i|^2 dx \\ &= -2 \int_{\mathbb{R}^3} \left(K * |u_i|^2 \right) |u_o|^2 dx - 2 \int_{\mathbb{R}^3} \xi_3 (\partial_{\xi_3} \hat{K}) |\widehat{|u_i|^2}| \widehat{|u_o|^2} d\xi \end{aligned}$$

and that

$$\xi_3 \partial_{\xi_3} \hat{K} = 8\pi \frac{\xi_3^2 (\xi_1^2 + \xi_2^2)}{|\xi|^4} = 8\pi \left(\frac{\xi_3^2}{|\xi|^2} - \frac{\xi_3^4}{|\xi|^4} \right) = 8\pi \widehat{\mathcal{R}}_3^2 - 8\pi \widehat{\mathcal{R}}_3^4,$$

we reduce the problem to the estimate of $|\langle \mathcal{R}_3^4 f, g \rangle_{L^2}|$ when f is supported in $\{|\bar{x}| \geq \gamma_2 R\}$ while g is supported in $\{|\bar{x}| \leq \gamma_1 R\}$, for some positive parameters $\gamma_{1,2}$ satisfying $d := \gamma_2 - \gamma_1 > 0$. Here \mathcal{R}_j^4 denotes the fourth power of the Riesz transform, and $\widehat{\mathcal{R}}_j^4$ its symbol in Fourier space.

- We compute $|\langle \mathcal{R}_3^4 f, g \rangle_{L^2}|$ by means of decay properties of the kernel associated to the parabolic biharmonic equation

$$\partial_t w + \Delta^2 w = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^3. \quad (1.15)$$

By denoting P_t the linear propagator associated to (1.15), for any two functions in $L^2(\mathbb{R}^3)$ we have the following identity, (see [Proposition 2.1](#)):

$$\langle \mathcal{R}_i^4 f, g \rangle_{L^2} = - \int_0^\infty \langle \partial_{x_i}^4 \frac{d}{dt} P_t f, g \rangle_{L^2} t dt.$$

Moreover, by using the explicit decay properties of the heat kernel for the parabolic biharmonic equation, that involves the Bessel function $J_{1/2}(s) = (\pi/2)^{-1/2} s^{-1/2} \sin(s)$, we prove that

$$|\langle \mathcal{R}_i^4 f, g \rangle_{L^2}| \lesssim R^{-1} \|g\|_{L^1} \|f\|_{L^1}$$

provided f and g are “well” localized as above (see [Theorem 2.2](#)). Summing up all the computations, the remainder term $H_R(u(t))$ is proved to be controlled by $o_R(1) + R^{-1} \|u\|_{\dot{H}^1(\mathbb{R}^3)}^2$.

- All the previous points will bring to the final estimate

$$\frac{d^2}{dt^2} V_{\rho_R+x_3^2}(t) \leq 4G(u(t)) + \epsilon_R(u(t)),$$

where $G(u) \lesssim -\tilde{\delta} \|u(t)\|_{\dot{H}^1(\mathbb{R}^3)}^2$ and $\epsilon_R(u(t)) \lesssim o_R(1) \|u(t)\|_{\dot{H}^1(\mathbb{R}^3)}^2$, which in turn implies the finite time blow-up via a convexity argument, provided $R \gg 1$.

1.1. Notation and structure of the paper. We collect here the notation used along the paper and we disclose how the paper is organized. We work in the three dimensional space \mathbb{R}^3 , and for a vector $x = (x_1, x_2, x_3) \in \mathbb{R}^3$ we denote by $\bar{x} \in \mathbb{R}^2$ the vector $\bar{x} = (x_1, x_2)$ given by the first two components of $x \in \mathbb{R}^3$. The differential operators ∇ and $\nabla \cdot$ are the common gradient and divergence operator in \mathbb{R}^3 . When using the subscript \bar{x} , i.e. $\nabla_{\bar{x}}$ or $\nabla_{\bar{x}} \cdot$, we mean that we are considering them as operators on \mathbb{R}^2 with respect to the variables (x_1, x_2) alone. The operator $\mathcal{F}f(\xi) = \hat{f}(\xi) = \int e^{-ix\xi} f(x) dx$ is the standard Fourier Transform, \mathcal{F}^{-1} being its inverse. \mathcal{R}_j is the j -th Riesz transform defined via the Fourier symbol $-i \frac{\xi_j}{|\xi|}$, i.e. $\mathcal{R}_j f(x) = \mathcal{F}^{-1} \left(-i \frac{\xi_j}{|\xi|} \hat{f} \right) (x)$. Powers of the Riesz transform are defined by means of powers of their symbols analogously. For $1 \leq p \leq \infty$ and $\Omega \subseteq \mathbb{R}^3$, $L^p(\Omega) = L^p(\Omega; \mathbb{C})$ are the classical Lebesgue spaces endowed with norm $\|f\|_{L^p} = (\int_{\Omega} |f(x)|^p dx)^{1/p}$ if $p \neq \infty$ or $\|f\|_{L^\infty} = \text{ess sup}_{x \in \Omega} |f(x)|$ for $p = \infty$. When $\Omega = \mathbb{R}^3$ we simply write L^p . For a function $f(x)$, $x \in \mathbb{R}^3$, we denote $\|f\|_{L^p_{\bar{x}}} = \|f(\cdot, x_3)\|_{L^p_{\bar{x}}(\mathbb{R}^2)}$ and similarly for more general domains $\Omega \subset \mathbb{R}^2$. We set $H^1 = H^1(\mathbb{R}^3; \mathbb{C}) := \{f \text{ s.t. } \int_{\mathbb{R}^3} (1 + |\xi|^2) |\hat{f}(\xi)|^2 d\xi < \infty\}$ and its homogeneous version $\dot{H}^1 = \dot{H}^1(\mathbb{R}^3; \mathbb{C}) := \{f \text{ s.t. } \int_{\mathbb{R}^3} |\xi|^2 |\hat{f}(\xi)|^2 d\xi < \infty\}$ endowed with their natural norms. Since we work on \mathbb{R}^3 , we simply denote $\int f dx = \int_{\mathbb{R}^3} f dx$ and often we write $\int f dx = \int_{\mathbb{R}} \int_{\mathbb{R}^2} f(\bar{x}, x_3) d\bar{x} dx_3$. The expression $(f * g)(x) := \int f(x-y)g(y) dy$ denotes the convolution operator between f and g . The L^2 inner product between two function f, g is denoted by $\langle f, g \rangle = \langle f, g \rangle_{L^2} := \int fg dx$. $\Re z$ and $\Im z$ are the common notations for the real and imaginary parts of a complex number z . When the bar-symbol over-lines a complex-valued function, we mean the complex conjugate. Given a measurable set $\mathcal{A} \subseteq \mathbb{R}^d$, $\mathbf{1}_{\mathcal{A}}(x)$ is the indicator function of \mathcal{A} . Finally, given two quantities A and B , we denote $A \lesssim B$ ($A \gtrsim B$, respectively) if there exists a positive constant C such that $A \leq CB$ ($A \geq CB$, respectively). If both relations hold true, we write $A \sim B$.

In [Section 2](#) we prove the essential integral and pointwise estimates for powers of the Riesz transforms for suitably localized functions. In [Section 3](#) we discuss the geometry of the energy functional and we disclose several properties leading to the control of the functional G in terms of $\|u\|_{\dot{H}^1}$. In [Section 4](#) we prove the blow-up in finite time for the focusing cubic NLS stated in [Theorem 1.3](#), then we conclude with [Section 5](#) where we prove the main result of the paper, namely the finite time blow-up for the Dipolar GPE. We collect in the [Appendix A](#) and [Appendix B](#) some useful identities and estimates used along the proofs in the paper.

2. LOCALIZATION PROPERTIES OF THE DIPOLAR KERNEL

This section provides the first technical tools we need in order to prove our main result concerning the finite time blow-up for the GPE [\(1.2\)](#). In the next Lemmas we prove some decay

estimates – pointwise and integral estimates – regarding the square and the 4-th power of the Riesz transforms for suitably localized functions. Firstly, we disclose a link between the 4-th power of the Riesz transform \mathcal{R}_j^4 and the linear propagator associated to the parabolic biharmonic equation, defined in terms of the Bessel functions. With this correspondence and some decay estimates for the parabolic biharmonic heat kernel we are able to show the decay estimate for $\langle \mathcal{R}_j^4 f, g \rangle$. Subsequently, we prove the pointwise estimates for \mathcal{R}_j^2 by using the explicit representation of \mathcal{R}_j^2 in terms of the singular integral defined in the principal value sense.

2.1. Intergal estimates for \mathcal{R}_j^4 . We start with the integral estimates for the fourth power of the Riesz transform, and, as anticipated above, we do it by means of some decay properties of the kernel associated to the parabolic biharmonic equation

$$\partial_t w + \Delta^2 w = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^3. \quad (2.1)$$

We denote by P_t the linear propagator associated to (2.1), namely $w(t, x) := P_t w_0(x)$ denotes the solution to the equation (2.1) with initial datum w_0 . We begin with the following proposition which provides a representation of \mathcal{R}_j^4 by using the functional calculus.

Proposition 2.1. *For any two functions in L^2 we have the following identity:*

$$\langle \mathcal{R}_i^4 f, g \rangle = - \int_0^\infty \langle \partial_{x_i}^4 \frac{d}{dt} P_t f, g \rangle t dt. \quad (2.2)$$

Proof. By passing to the frequencies space, it is easy to see that $\widehat{P_t f}(\xi) := e^{-t|\xi|^4} \widehat{f}(\xi)$ and we observe, by integration by parts, that

$$\xi_i^4 |\xi|^4 \int_0^\infty e^{-t|\xi|^4} t dt = \frac{\xi_i^4}{|\xi|^4}; \quad (2.3)$$

hence

$$\begin{aligned} \int_0^\infty \langle \partial_{x_i}^4 \frac{d}{dt} P_t f, g \rangle t dt &= \langle \int_0^\infty \partial_{x_i}^4 \frac{d}{dt} (P_t f) t dt, g \rangle = \langle \int_0^\infty \xi_i^4 \frac{d}{dt} (e^{-t|\xi|^4} \widehat{f}) t dt, \widehat{g} \rangle \\ &= - \langle \xi_i^4 |\xi|^4 \widehat{f} \int_0^\infty e^{-t|\xi|^4} t dt, \widehat{g} \rangle = - \langle \frac{\xi_i^4}{|\xi|^4} \widehat{f}, \widehat{g} \rangle = - \langle \mathcal{R}_i^4 f, g \rangle, \end{aligned}$$

where the change of order of integration (in time and in space) is justified by means of the Fubini-Tonelli's theorem, and we used the Plancherel identity when passing from the frequencies space to the physical space, and vice versa. \square

We are now in position to prove a decay estimate for functions supported outside a cylinder of radius $\gtrsim R$. In order to do that, we explicitly write the heat kernel of P_t . We introduce, for $t > 0$ and $x \in \mathbb{R}^3$

$$p_t(x) = \alpha \frac{k(\mu)}{t^{3/4}}, \quad \mu = \frac{|x|}{t^{1/4}}$$

and

$$k(\mu) = \mu^{-2} \int_0^\infty e^{-s^4} (\mu s)^{3/2} J_{1/2}(\mu s) ds,$$

where $J_{1/2}$ is the $\frac{1}{2}$ -th Bessel function, and $\alpha^{-1} := \frac{4\pi}{3} \int_0^\infty s^2 k(s) ds$ is a positive normalization constant. We refer to [14] for these definitions and further discussions about the heat kernel of the parabolic biharmonic equation. We recall that the $\frac{1}{2}$ -th Bessel function is given by

$$J_{1/2}(s) = (\pi/2)^{-1/2} s^{-1/2} \sin(s),$$

then

$$P_t f(x) = (p_t * f)(x) = c \int_{\mathbb{R}^3} f(x-y) \int_0^\infty \frac{1}{|y|^3} e^{-ts^4/|y|^4} s \sin(s) ds dy,$$

and therefore

$$\frac{d}{dt} P_t f(x) = -c \int_{\mathbb{R}^3} f(x-y) \int_0^\infty \frac{1}{|y|^3} e^{-ts^4/|y|^4} \frac{s^5}{|y|^4} \sin(s) ds dy.$$

We are ready to prove the following result.

Theorem 2.2. *Assume that $f, g \in L^1 \cap L^2$, and that f is supported in $\{|\bar{x}| \geq \gamma_2 R\}$ while g is supported in $\{|\bar{x}| \leq \gamma_1 R\}$, for some positive parameters $\gamma_{1,2}$ satisfying $d := \gamma_2 - \gamma_1 > 0$. Then*

$$|\langle \mathcal{R}_i^4 f, g \rangle| \lesssim R^{-1} \|g\|_{L^1} \|f\|_{L^1}. \quad (2.4)$$

Proof. With the change of variable $s^4 |y|^{-4} = \tau$ we get

$$\frac{d}{dt} P_t f = -\frac{c}{4} \int_{\mathbb{R}^3} \int_0^\infty \frac{1}{|y|} e^{-t\tau} \tau^{1/2} \sin(\tau^{1/4} |y|) f(x-y) d\tau dy$$

and hence, by a change of variable in space,

$$\frac{d}{dt} P_t f = -\frac{c}{4} \int_{\mathbb{R}^3} \int_0^\infty \frac{1}{|x-y|} e^{-t\tau} \tau^{1/2} \sin(\tau^{1/4} |x-y|) f(y) d\tau dy.$$

An explicit computation, see [Appendix B](#), gives that there exist an integer $M \geq 1$ and M pairs of polynomials $(\tilde{q}_k, q_k)_{k \in \{1, \dots, M\}}$ with nonnegative coefficients, such that $\min_{k \in \{1, \dots, M\}} \{deg(q_k)\} \geq 1$ – here deg stands for the degree of the polynomial – and satisfying

$$\left| \partial_{x_i}^4 \left(\frac{1}{|x-y|} \sin(\tau^{1/4} |x-y|) \right) \right| \lesssim \sum_{k=1}^M \frac{\tilde{q}_k(\tau^{1/4})}{q_k(|x-y|)}. \quad (2.5)$$

At this point, by using the identity (2.2) we infer the following:

$$\begin{aligned}
|\langle \mathcal{R}_i^4 f, g \rangle| &= \left| \int_0^\infty \langle \partial_{x_i}^4 \frac{d}{dt} P_t f, g \rangle t dt \right| \\
&= c \left| \int_0^\infty t \int g(x) \left(\int \int_0^\infty \partial_{x_i}^4 \left(\frac{1}{|x-y|} \sin(\tau^{1/4}|x-y|) \right) e^{-t\tau} \tau^{1/2} f(y) d\tau dy \right) dx dt \right| \\
&\lesssim \int_0^\infty t \int |g(x)| \left(\int \int_0^\infty \sum_{k=1}^M \frac{\tilde{q}_k(\tau^{1/4})}{q_k(|x-y|)} e^{-t\tau} \tau^{1/2} |f(y)| d\tau dy \right) dx dt \\
&= \int_0^\infty t \int |g(x)| \left(\int \int_0^\infty \sum_{k=1}^M \frac{\tilde{q}_k(\tau^{1/4})}{q_k(|y|)} e^{-t\tau} \tau^{1/2} |f(x-y)| d\tau dy \right) dx dt \\
&\leq \int_0^\infty t \int_{\{|\bar{x}| \leq \gamma_1 R\}} |g(x)| \left(\int_{\{|\bar{x}-\bar{y}| \geq \gamma_2 R\}} \int_0^\infty \sum_{k=1}^M \frac{\tilde{q}_k(\tau^{1/4})}{q_k(|\bar{y}|)} e^{-t\tau} \tau^{1/2} |f(x-y)| d\tau dy \right) dx dt.
\end{aligned}$$

Therefore, as the support of $f(x-y)$ is contained in $|\bar{x}-\bar{y}| \geq \gamma_2 R$ and the one of g is contained in $|\bar{x}| \leq \gamma_1 R$, we get that $|\bar{y}| \geq dR$. Hence, by defining $\beta = \frac{1}{4} \max_{k \in \{1, \dots, M\}} \{deg(\tilde{q}_k)\}$, we can bound

$$\begin{aligned}
|\langle \mathcal{R}_i^4 f, g \rangle| &\lesssim R^{-1} \|f\|_{L^1} \|g\|_{L^1} \int_0^\infty \int_0^\infty e^{-t\tau} \tau^{1/2} \tilde{q}_k(\tau^{1/4}) t d\tau dt \\
&\sim R^{-1} \|f\|_{L^1} \|g\|_{L^1} \int_1^\infty \int_1^\infty e^{-t\tau} \tau^{1/2} \tau^{(\max_{k \in \{1, \dots, M\}} deg(\tilde{q}_k))/4} t d\tau dt \\
&= R^{-1} \|f\|_{L^1} \|g\|_{L^1} \int_1^\infty \int_1^\infty e^{-t\tau} \tau^{\beta+1/2} t d\tau dt \\
&\lesssim R^{-1} \|f\|_{L^1} \|g\|_{L^1}
\end{aligned}$$

where we used the Fubini-Tonelli's theorem. The proof of (2.4) is concluded. \square

Remark 2.3. It is straightforward to observe that in (2.3) we can replace the symbol ξ_j^4 with $\xi_k^2 \xi_h^2$, for $k \neq h$, to get

$$\frac{\xi_k^2 \xi_h^2}{|\xi|^4} = \xi_k^2 \xi_h^2 |\xi|^{-4} \int_0^\infty e^{-t|\xi|^4} t dt, \quad (2.6)$$

and consequently

$$\langle \mathcal{R}_k^2 \mathcal{R}_h^2 f, g \rangle = - \int_0^\infty \langle \partial_{x_k}^2 \partial_{x_h}^2 \frac{d}{dt} P_t f, g \rangle t dt \quad (2.7)$$

The identities (2.6) and (2.7) of the remark above easily imply an analogous of **Theorem 2.2** (by repeating its proof with the obvious modifications) for the operator $\mathcal{R}_k^2 \mathcal{R}_h^2$ replacing \mathcal{R}_j^4 . More precisely:

Lemma 2.4. *Assume that $f, g \in L^1 \cap L^2$, and that f is supported in $\{|\bar{x}| \geq \gamma_2 R\}$ while g is supported in $\{|\bar{x}| \leq \gamma_1 R\}$, for some $\gamma_{1,2} > 0$ satisfying $d := \gamma_2 - \gamma_1 > 0$. Then*

$$|\langle \mathcal{R}_k^2 \mathcal{R}_h^2 f, g \rangle| \lesssim R^{-1} \|g\|_{L^1} \|f\|_{L^1}.$$

2.2. Pointwise estimate for \mathcal{R}_j^2 . We turn now the attention to the square of the Riesz transform. In the subsequent results, we will use a cut-off function χ satisfying the following: $\chi(x)$ is a localization function supported in the cylinder $\{|\bar{x}| \leq 1\}$ which is nonnegative and bounded, with $\|\chi\|_{L^\infty} \leq 1$. For a positive parameter γ , we define by $\chi_{\{|\bar{x}| \leq \gamma R\}}$ the rescaled function $\chi(x/\gamma R)$ (hence $\chi_{\{|\bar{x}| \leq \gamma R\}}$ is bounded, positive and supported in the cylinder of radius γR). The proof of the next lemmas is inspired by [22].

Lemma 2.5. *For any (regular) function f the following pointwise estimate is satisfied: provided $d := \gamma_2 - \gamma_1 > 0$, there exists an universal constant $C = C(d) > 0$ such that*

$$|\chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \mathcal{R}_j^2[(1 - \chi_{\{|\bar{x}| \leq \gamma_2 R\}})f](x)| \leq CR^{-3} \chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \|f\|_{L^1(|\bar{x}| \geq \gamma_2 R)}. \quad (2.8)$$

Proof. In the principal value sense, the square of the Riesz transform acts on a function g as

$$\mathcal{R}_j^2 g(x) = \iint \frac{x_j - y_j}{|x - y|^{3+1}} \frac{y_j - z_j}{|y - z|^{3+1}} g(z) dz dy.$$

Let $g(x) = \chi_{\{|\bar{x}| \geq \gamma_2 R\}}(x) f(x)$. Then

$$\chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \mathcal{R}_j^2 g(x) = \iint \left(\frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \right) g(x - z) dz.$$

Since g is supported in the exterior of a cylinder of radius $\gamma_2 R$, we can assume $|\bar{x} - \bar{z}| \geq \gamma_2 R$, and for the function $\chi_{\{|\bar{x}| \leq \gamma_1 R\}}$ is supported by definition in the cylinder of radius $\gamma_1 R$, we can assume $|\bar{x}| \leq \gamma_1 R$: therefore we have that $|\bar{z}| \geq dR$. This implies that $\{|\bar{y}| \leq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|\} = \emptyset$. Indeed,

$$\frac{1}{2}|\bar{z}| \geq |\bar{z} - \bar{y}| \geq |\bar{z}| - |\bar{y}| \implies |\bar{y}| \geq \frac{1}{2}|\bar{z}| \geq \frac{d}{2}R, \quad (2.9)$$

hence we have the following splitting:

$$\begin{aligned} I &= \int \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy = \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \\ &\quad + \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \\ &\quad + \int_{\{|\bar{y}| \geq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \geq \frac{1}{2}|\bar{z}|\}} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \\ &= \mathcal{I} + \mathcal{II} + \mathcal{III}. \end{aligned} \quad (2.10)$$

Estimate for the term \mathcal{I} . Let us focus on the first integral \mathcal{I} . The domain of integration of this integral is the cylinder of radius R parallel to the y_3 axis. Therefore

$$\mathcal{I} = \mathcal{I}_1 + \mathcal{I}_2 = \int_{|y_3| \leq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} d\bar{y} dy_3 + \int_{|y_3| \geq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} d\bar{y} dy_3.$$

For the term \mathcal{I}_1 we first notice that

$$\int_{|y_3| \leq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} d\bar{y} dy_3 = 0$$

since the domain is invariant under the change of variables $y \mapsto -y$. Therefore

$$A_1 = \int_{|y_3| \leq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} \left(\frac{z_j - y_j}{|z - y|^4} - \frac{z_j}{|z|^4} \right) d\bar{y} dy_3$$

and we write

$$\frac{z_j - y_j}{|z - y|^4} - \frac{z_j}{|z|^4} = \int_0^1 h'(s) ds$$

where

$$h(s) = \frac{z_j - sy_j}{|z - sy|^4}, \quad s \in [0, 1].$$

We have, by a straightforward calculation, that

$$h'(s) = -\frac{y_j}{|z - sy|^4} + 4 \frac{z_j - sy_j}{|z - sy|^4} (z - sy) \cdot y$$

and

$$|h'(s)| \lesssim \frac{|y|}{|z - sy|^4}.$$

Hence, by observing that $|z - sy| \geq |\bar{z} - s\bar{y}| \geq |\bar{z}| - s|\bar{y}| \geq \frac{3d}{4}R$ as $s|\bar{y}| \leq \frac{d}{4}R$, we get that

$$\max_{s \in [0, 1]} |h'(s)| \lesssim R^{-4}|y|,$$

then

$$\begin{aligned} \mathcal{I}_1 &= \int_{|y_3| \leq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} \left(\int_0^1 h'(s) ds \right) d\bar{y} dy_3 \lesssim R^{-4} \int_{|y_3| \leq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{1}{|y|^2} d\bar{y} dy_3 \\ &\lesssim R^{-4} \int_{|y| \leq \frac{\sqrt{2}d}{4}R} \frac{1}{|y|^2} dy \lesssim R^{-3}. \end{aligned} \quad (2.11)$$

The term \mathcal{I}_2 can be estimated as follows:

$$\begin{aligned} \mathcal{I}_2 &= \int_{|y_3| \geq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} d\bar{y} dy_3 \leq \int_{|y_3| \geq \frac{d}{4}R} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{1}{|y|^3} \frac{1}{|z - y|^3} d\bar{y} dy_3 \\ &\leq \int_{|y_3| \geq \frac{d}{4}R} \frac{1}{|y_3|^3} \int_{|\bar{y}| \leq \frac{d}{4}R} \frac{1}{|\bar{z} - \bar{y}|^3} d\bar{y} dy_3 \lesssim \int_{|y_3| \geq \frac{d}{4}R} \frac{1}{|y_3|^3} dy_3 \left(\frac{1}{R^3} \int_{|\bar{y}| \leq \frac{d}{4}R} d\bar{y} \right) \end{aligned}$$

where we used again the fact that if $|\bar{y}| \leq \frac{d}{4}R$ then $|\bar{z} - \bar{y}| \geq \frac{3d}{4}R$, hence we conclude with

$$\mathcal{I}_2 \lesssim R^{-2}R^{-1} = R^{-3} \quad (2.12)$$

In conclusion, by summing up the two estimates (2.11) and (2.12) we get

$$\mathcal{I} \lesssim R^{-3}. \quad (2.13)$$

Estimate for the term \mathcal{II} . We adopt a similar approach for the term \mathcal{II} that we split in two further terms:

$$\begin{aligned}\mathcal{II} &= \int_{|z_3 - y_3| \leq |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy + \int_{|z_3 - y_3| \geq |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \\ &= \mathcal{II}_1 + \mathcal{II}_2\end{aligned}$$

We first notice that

$$\int_{|z_3 - y_3| \leq |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{z_j - y_j}{|z - y|^4} d\bar{y} dy_3 = 0$$

since the domain is invariant under the change of variable $y \mapsto 2z - y$.

Estimate for the term \mathcal{II}_1 . Therefore

$$\begin{aligned}\mathcal{II}_1 &= \int_{|z_3 - y_3| \leq |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{z_j - y_j}{|z - y|^4} \left(\frac{y_j}{|y|^4} - \frac{z_j}{|z|^4} \right) d\bar{y} dy_3 \\ &= \int_{|z_3 - y_3| \leq |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{z_j - y_j}{|z - y|^4} \left(\int_0^1 h'(s) ds \right) d\bar{y} dy_3\end{aligned}$$

where

$$h(s) = \frac{sy_j + (1-s)z_j}{|sy_j + (1-s)z_j|^4}, \quad s \in [0, 1].$$

We compute

$$h'(s) = \frac{y_j - z_j}{|sy_j + (1-s)z_j|^4} + 4 \frac{sy_j + (1-s)z_j}{|sy_j + (1-s)z_j|^5} \frac{sy_j + (1-s)z_j}{|sy_j + (1-s)z_j|} \cdot (y_j - z_j)$$

and hence

$$|h'(s)| \lesssim \frac{|y_j - z_j|}{|sy_j + (1-s)z_j|^4}.$$

Now we observe, as $s|\bar{y} - \bar{z}| \leq \frac{1}{2}|\bar{z}|$, that $|sy_j + (1-s)z_j| = |s(y_j - z_j) + z_j| \geq |s(\bar{y} - \bar{z}) + \bar{z}| \geq |\bar{z}| - s|\bar{y} - \bar{z}| \geq \frac{1}{2}|\bar{z}|$ and then

$$\max_{s \in [0, 1]} |h'(s)| \lesssim \frac{|y_j - z_j|}{|\bar{z}|^4}$$

which allows us to continue the estimate for \mathcal{II}_1 as follows:

$$\begin{aligned}\mathcal{II}_1 &\lesssim \frac{1}{|\bar{z}|^4} \int_{|z_3 - y_3| \leq |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{1}{|z - y|^2} d\bar{y} dy_3 \\ &\lesssim \frac{1}{|\bar{z}|^4} \int_{|z - y| \lesssim |\bar{z}|} \frac{1}{|z - y|^2} dy \lesssim |\bar{z}|^{-3} \lesssim R^{-3}.\end{aligned}\tag{2.14}$$

Estimate for the term \mathcal{II}_2 . It remains to prove a suitable estimate for the remaining term \mathcal{II}_2 . We use (2.9) and we estimate

$$\begin{aligned}
\mathcal{II}_2 &= \int_{|z_3 - y_3| > |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} d\bar{y} dy_3 \\
&\leq \int_{|z_3 - y_3| > |\bar{z}|} \int_{|\bar{z} - \bar{y}| \leq \frac{1}{2}|\bar{z}|} \frac{1}{|y|^3} \frac{1}{|z - y|^3} d\bar{y} dy_3 \\
&\leq \int_{|z_3 - y_3| > |\bar{z}|} \int_{|\bar{y}| \geq \frac{1}{2}|\bar{z}|} \frac{1}{|y|^3} \frac{1}{|z - y|^3} d\bar{y} dy_3 \\
&\leq \int_{|z_3 - y_3| > |\bar{z}|} \frac{1}{|z_3 - y_3|^3} dy_3 \int_{|\bar{y}| \geq \frac{1}{2}|\bar{z}|} \frac{1}{|\bar{y}|^3} d\bar{y} \leq |\bar{z}|^{-2} |\bar{z}|^{-1} \lesssim R^{-3}.
\end{aligned} \tag{2.15}$$

We conclude, by summing up (2.14) and (2.15), that

$$\mathcal{II} \lesssim R^{-3}. \tag{2.16}$$

Estimate for the term \mathcal{III} . It is left to estimate the integral

$$\mathcal{III} = \int_{\{|\bar{y}| \geq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \geq \frac{1}{2}|\bar{z}|\}} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy.$$

By the Cauchy-Schwarz's inequality

$$\begin{aligned}
\mathcal{III} &= \int_{\{|\bar{y}| \geq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \geq \frac{1}{2}|\bar{z}|\}} \frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \leq \int_{\{|\bar{y}| \geq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \geq \frac{1}{2}|\bar{z}|\}} \frac{1}{|y|^3} \frac{1}{|z - y|^3} dy \\
&\leq \left(\int_{\{|\bar{y}| \geq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \geq \frac{1}{2}|\bar{z}|\}} \frac{1}{|y|^6} dy \right)^{1/2} \left(\int_{\{|\bar{y}| \geq \frac{d}{4}R\} \cap \{|\bar{z} - \bar{y}| \geq \frac{1}{2}|\bar{z}|\}} \frac{1}{|z - y|^6} dy \right)^{1/2} \\
&\leq \left(\int_{\{|y| \geq \frac{d}{4}R\}} \frac{1}{|y|^6} dy \right)^{1/2} \left(\int_{\{|z - y| \geq \frac{1}{2}|\bar{z}|\}} \frac{1}{|z - y|^6} dy \right)^{1/2} \\
&\lesssim R^{-3/2} |\bar{z}|^{-3/2} \lesssim R^{-3}.
\end{aligned} \tag{2.17}$$

The proof of the lemma is therefore concluded by observing that the integral I defined in (2.10) can be bounded, by using (2.13), (2.16) and (2.17), by

$$I := \mathcal{I} + \mathcal{II} + \mathcal{III} \lesssim R^{-3},$$

and hence

$$\begin{aligned}
|\chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \mathcal{R}_j^2 g(x)| &= \chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \left| \iint \left(\frac{y_j}{|y|^4} \frac{z_j - y_j}{|z - y|^4} dy \right) g(x - z) dz \right| \\
&\lesssim R^{-3} \chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \int |g(x - z)| dz \\
&\lesssim R^{-3} \chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x) \|f\|_{L^1(|\bar{x}| \geq \gamma_2 R)}
\end{aligned}$$

which is the estimate stated in (2.8). \square

We have an estimate similar to (2.8) if we localize inside a cylinder the function on which R_j^2 acts, and we then truncate everything with a function supported in the exterior of another cylinder.

Lemma 2.6. *For any (regular) function f the following pointwise estimate is satisfied: provided $d := \gamma_1 - \gamma_2 > 0$, there exists an universal constant $C = C(d) > 0$ such that*

$$|(1 - \chi_{|\bar{x}| \leq \gamma_1 R})(x) \mathcal{R}_j^2[(\chi_{\{|\bar{x}| \leq \gamma_2 R\}})f](x)| \leq CR^{-3} |1 - \chi_{\{|\bar{x}| \leq \gamma_1 R\}}(x)| \|f\|_{L^1(|\bar{x}| \leq \gamma_2 R)}. \quad (2.18)$$

Proof. The proof is analogous to the one for Lemma 2.5, so we skip the details. \square

3. VARIATIONAL STRUCTURE OF THE ENERGY FUNCTIONAL AND CONSEQUENCES

We pass now to the discussion on the variational structure of the energy functional and its relation to the existence of standing waves for (1.2). The following arguments are valid in the same fashion for the NLS equation (1.13), when the parameter $\lambda_2 = 0$, even if it is worth mentioning that for NLS the existence of standing waves is nowadays classical. We recall the two different approaches to prove existence of Ground States for the GPE.

The first strategy is due to Antonelli and Sparber, see [2], where existence is proved by means of minimization of the Weinstein functional

$$J(v) := \frac{\|\nabla v\|_{L^2}^3 \|v\|_{L^2}}{-\lambda_1 \|v\|_{L^4}^4 - \lambda_2 \int (K * |v|^2) |v|^2 dx}.$$

The alternative way, see the work of Jeanjean and the first author [6], is based on topological methods, where the existence of Ground States is shown by means of the existence of critical points of the energy functional under the mass constraint (1.7). In the latter approach the parameter κ which appears in (1.6) is found as Lagrange multiplier. Even if the energy functional is unbounded from below on $S(1)$, when restricting to states which are stationary for the evolution equation, i.e. they satisfy (1.6), then the energy is bounded from below by a positive constant. The latter constant, which corresponds to the mountain pass level, is reached. The mountain pass solutions therefore correspond to the least energy states (which are called Ground States, precisely). We pass now to the analysis of the geometry of the functional $E(u)$ on $S(c)$, and to this aim we introduce the L^2 -preserving scaling:

$$u^\mu(x) = \mu^{3/2} u(\mu x), \quad \mu > 0.$$

We report the next crucial lemma from [6]. We recall the definition of $V(c)$ given in (1.12):

$$V(c) = \{u \in H^1(\mathbb{R}^3) \quad \text{s.t.} \quad \|u\|_{L^2(\mathbb{R}^3)}^2 = c \text{ and } G(u) = 0\}.$$

Lemma 3.1. [6, Lemma 3.3] *Suppose that u belongs to the manifold $S(c)$ and moreover that it satisfies $\int (\lambda_1 + \lambda_2 \widehat{K}(\xi)) (|u|^2)^2 d\xi < 0$. Then the following properties hold true:*

- *there exists a unique $\tilde{\mu}(u) > 0$, such that $u^{\tilde{\mu}} \in V(c)$;*

- the map $\mu \mapsto E(u^\mu)$ is concave on $[\tilde{\mu}, \infty)$;
- $\tilde{\mu}(u) < 1$ if and only if $G(u) < 0$;
- $\tilde{\mu}(u) = 1$ if and only if $G(u) = 0$;
- the functional G satisfies

$$G(u^\mu) \begin{cases} > 0, & \forall \mu \in (0, \tilde{\mu}(u)) \\ < 0, & \forall \mu \in (\tilde{\mu}(u), +\infty) \end{cases} ;$$

- $E(u^\mu) < E(u^{\tilde{\mu}})$, for any $\mu > 0$ and $\mu \neq \tilde{\mu}$;
- $\frac{d}{d\mu} E(u^\mu) = \frac{1}{\mu} G(u^\mu)$, $\forall \mu > 0$.

With [Lemma 3.1](#) at hand, we can prove the next proposition which basically shows the dichotomy between the scattering and blow-up for [\(1.2\)](#) in terms of the quantities $\gamma(\|u_0\|_{L^2}^2)$ and $G(u_0)$.

Proposition 3.2. *Suppose that the initial datum u_0 satisfies $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$ and $G(u_0) > 0$, then*

$$M(u_0)E(u_0) < M(Q)E(Q) \tag{3.1}$$

and

$$\|u_0\|_{L^2} \|\nabla u_0\|_{L^2} < \|Q\|_{L^2} \|\nabla Q\|_{L^2}. \tag{3.2}$$

Conversely, if the conditions expressed in [\(3.1\)](#) and [\(3.2\)](#) hold true, then the initial datum u_0 satisfies $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$ and $G(u_0) > 0$.

Remark 3.3. We point out that we gave the proof of the first implication in our previous work [\[5\]](#), but we repeat it below as in that paper some steps were not rigorously justified (it is worth mentioning that the claim was however correct, and the validity of the result was not affected by that carelessness).

Proof. We start with the first implication. From the definition of the quantities in [\(1.5\)](#), [\(1.9\)](#) and [\(1.10\)](#), we straightforwardly obtain the identity

$$E(u_0) - \frac{1}{3}G(u_0) = \frac{1}{6}T(u_0). \tag{3.3}$$

Due to the scaling invariance properties of the Weinstein functional, we note that $Q_\mu := \mu Q(\mu x)$ is again a minimizer for the Weinstein functional with

$$\begin{aligned} \|Q_\mu\|_{L^2}^2 &= \mu^{-1} \|Q\|_{L^2}^2, \\ \|\nabla Q_\mu\|_{L^2}^2 &= \mu \|\nabla Q\|_{L^2}^2. \end{aligned}$$

We notice that $Q(x)e^{it}$ is a standing wave solution to the evolution equation and by the symmetry of the equation it is well known that $Q_\mu e^{i\mu^2 t} = \mu Q(\mu x)e^{i\mu^2 t}$ is another standing wave solution to

$$-\frac{1}{2}\Delta Q_\mu + \left(\lambda_1 |Q_\mu|^2 Q_\mu + \lambda_2 (K * |Q_\mu|^2) Q_\mu\right) + \mu^2 Q_\mu = 0,$$

that necessarily satisfies $G(Q_\mu) = 0$. Hence $E(Q_\mu) = \frac{1}{6}\|\nabla Q_\mu\|_{L^2}^2$.

Provided we choose the parameter μ such that $\|Q_\mu\|_{L^2}^2 = \|u_0\|_{L^2}^2$, i.e. Q_μ belongs to the constraint $S(c)$, $c = \|u_0\|_{L^2}^2$, we get (using the hypothesis)

$$E(u_0) < \gamma(\|u_0\|_{L^2}^2) = \gamma(\|Q_\mu\|_{L^2}^2) = E(Q_\mu). \quad (3.4)$$

From (3.4) we obtain

$$\|u_0\|_{L^2}^2 E(u_0) < \|Q\|_{L^2}^2 E(Q),$$

which corresponds to (3.1). It is worth remarking how we can claim the equality in (3.4) (this is precisely the clarification we do with respect to what we wrote in Remark 3.3). It is crucial to notice that if Q is a standing state (solving the elliptic equation with a corresponding Lagrange multiplier), then Q_μ is a standing state for any $\mu > 0$. On the other hand if Q is not a standing state (i.e. it does not solve the elliptic equation with any Lagrange multiplier), then Q_μ is not a standing state for any $\mu > 0$.

If we take two standing states with the same mass, let say w and v , with their corresponding Lagrange multipliers, and such that $E(w) < E(v)$, then $E(w_\mu) < E(v_\mu)$ for any $\mu > 0$. This is evident by the fact that $E(w_\mu) = \frac{1}{6}\|\nabla w_\mu\|_{L^2}^2 = \frac{\mu}{6}\|\nabla w\|_{L^2}^2 = \frac{\mu}{6}E(w)$ for any $\mu > 0$ (indeed $G(w_\mu)$ is always 0). Therefore if $E(w) < E(v)$ then $E(w_\mu) < \frac{\mu}{6}\|\nabla v\|_{L^2}^2 = E(v_\mu)$.

This implies that in the case of a Mountain Pass solution, if one takes a standing wave Q such that $E(Q) = \gamma(\|Q\|_{L^2}^2)$ then $E(Q_\mu) = \gamma(\|Q_\mu\|_{L^2}^2)$.

We prove now the validity of the other condition. If $G(u_0) > 0$ and $E(u_0) < \gamma(\|u_0\|_{L^2}^2) = E(Q_\mu)$, then we have

$$\frac{1}{6}\|\nabla Q_\mu\|_{L^2}^2 = E(Q_\mu) > E(u_0) > E(u_0) - \frac{1}{3}G(u_0) = \frac{1}{6}\|\nabla u_0\|_{L^2}^2$$

and hence

$$\|u_0\|_{L^2}\|\nabla u_0\|_{L^2} < \|Q\|_{L^2}\|\nabla Q\|_{L^2}.$$

Let us prove the reverse implication.

First of all we notice that if Q is a minimizer for the Weinstein functional, then $E(Q) = \gamma(\|Q\|_{L^2}^2)$. We take a rescaling Q_μ of Q such that $\|Q_\mu\|_{L^2}^2 = \|u_0\|_{L^2}^2$ and as before we can claim that $E(Q_\mu) = \gamma(\|u_0\|_{L^2}^2)$. Therefore (3.1) implies

$$M(u_0)E(u_0) < M(Q)E(Q) = M(Q_\mu)E(Q_\mu) = M(Q_\mu)\gamma(\|u_0\|_{L^2}^2) \implies E(u_0) < \gamma(\|u_0\|_{L^2}^2).$$

Let us focus on the statement “(3.2) $\implies G(u_0) > 0$ ”. Suppose that $G(u_0) \leq 0$ and consider $\tilde{\mu}$ such that $u^{\tilde{\mu}} \in V(\|u_0\|_{L^2}^2)$. By using Lemma 3.1 such $\tilde{\mu}$ do exists, $G(u^{\tilde{\mu}}) = 0$ (by the very

definition of $V(\|u_0\|_{L^2}^2)$ and $\tilde{\mu} \leq 1$. (In particular, if $G(u_0) = 0$ then $\tilde{\mu} = 1$.) Hence

$$E(u^{\tilde{\mu}}) = \frac{1}{6} \|\nabla u^{\tilde{\mu}}\|_{L^2}^2 = \frac{\tilde{\mu}}{6} \|\nabla u\|_{L^2}^2 \leq \frac{1}{6} \|\nabla u\|_{L^2}^2 < \frac{1}{6} \|\nabla Q\|_{L^2}^2 = E(Q).$$

This concludes the proof since we got a function, $u^{\tilde{\mu}}$, such that $E(u^{\tilde{\mu}}) < E(Q)$, which contradicts the minimality of $E(Q)$. \square

Remark 3.4. It is straightforward to see that in [Proposition 3.2](#), provided $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$, the condition [\(3.2\)](#) replaced by

$$\|u_0\|_{L^2} \|\nabla u_0\|_{L^2} > \|Q\|_{L^2} \|\nabla Q\|_{L^2} \quad (3.5)$$

will imply that $G(u_0) < 0$, and conversely [\(3.5\)](#) is satisfied provided we assume $G(u_0) < 0$. Hence $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$ and $G(u_0) > 0$ or $G(u_0) < 0$ give the dichotomy between scattering and blow-up for [\(1.2\)](#). It is worth mentioning that when restricting to the cubic NLS case, the latter ones are the same described by Holmer and Roudenko in [\[18\]](#), namely [\(3.1\)](#), [\(3.2\)](#), and [\(3.5\)](#).

Lemma 3.5. *If the initial datum u_0 satisfies $E(u_0) < \gamma(\|u_0\|_{L^2}^2)$ and $G(u_0) < 0$ then $G(u(t)) < 0$ for any $t \in (-T_{min}, T_{max})$. More precisely, there exists a positive constant $\delta > 0$ such that $G(u(t)) \leq -\delta$ for any $t \in (-T_{min}, T_{max})$.*

Proof. Suppose that $G(u(t)) > 0$ for some time $t \in (-T_{min}, T_{max})$; then by the continuity in time of the function $G(u(t))$ there exists \tilde{t} such that $G(u(\tilde{t})) = 0$. By definition we have therefore $\gamma(c) \leq E(u(\tilde{t})) = E(u_0)$ which is a contradiction with respect to the definition of $\gamma(c)$.

We prove now the uniform bound from below away from zero. We simply denote $u = u(t)$. By [Lemma 3.1](#) – third claim – there exists $\tilde{\mu} \in (0, 1)$ such that $G(u^{\tilde{\mu}}) = 0$. Then

$$E(u_0) - E(u^{\tilde{\mu}}) = (1 - \tilde{\mu}) \frac{d}{d\mu} E(u^\mu)|_{\mu=\tilde{\mu}}$$

for some $\bar{\mu} \in (\tilde{\mu}, 1)$, and due to the concavity of $\mu \mapsto E(u^\mu)$ – see [Lemma 3.1](#), second claim – we have that

$$E(u_0) - E(u^{\tilde{\mu}}) = (1 - \tilde{\mu}) \frac{d}{d\mu} E(u^\mu)|_{\mu=\bar{\mu}} \geq (1 - \tilde{\mu}) \frac{d}{d\mu} E(u^\mu)|_{\mu=1} = (1 - \tilde{\mu}) G(u)$$

where in the last equality we used the last claim of [Lemma 3.1](#). Hence

$$G(u(t)) \leq (1 - \tilde{\mu})^{-1} (E(u_0) - E(u^{\tilde{\mu}})) \leq (1 - \tilde{\mu})^{-1} (E(u_0) - \gamma(c)).$$

The proof is complete with $\delta = (1 - \tilde{\mu})^{-1} (\gamma(c) - E(u_0))$. \square

The previous Lemma implies the pointwise-in-time bound for the function $G(u(t))$, by means of the homogeneous Sobolev \dot{H}^1 -norm of $u(t)$.

Lemma 3.6. *There exists $\alpha > 0$ such that $G(u(t)) \leq -\alpha \|u(t)\|_{\dot{H}^1}^2$ for any $t \in (-T_{min}, T_{max})$.*

Proof. It follows from [Lemma 3.5](#) and the identity [\(3.3\)](#), that $\|u(t)\|_{\dot{H}^1}^2 > 6E(u_0)$. By exploiting again the identity [\(3.3\)](#) we write $\|u(t)\|_{\dot{H}^1}^2 = 6E - 2G(u(t))$, so we have

$$G(u(t)) + \alpha\|u(t)\|_{\dot{H}^1}^2 = (1 - 2\alpha)G(u(t)) + 6\alpha E.$$

Therefore for $\alpha \ll 1$ the claim follows. \square

We give the following simple consequence of the previous Lemma.

Corollary 3.7. *There exists a positive constant $c > 0$ such that*

$$\inf_{t \in (-T_{min}, T_{max})} \|u(t)\|_{\dot{H}^1} \geq c.$$

Proof. Suppose that there exists a sequence of times $\{t_n\}_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} \|u(t_n)\|_{\dot{H}^1} = 0$. Then by the Gagliardo-Nirenberg's inequality $\lim_{n \rightarrow \infty} \|u(t_n)\|_{L^4} = 0$ as well. But then $G(u(t_n)) \rightarrow 0$, since by the $L^2 \mapsto L^2$ property of the dipolar kernel

$$|G(u(t_n))| \lesssim \|u(t_n)\|_{\dot{H}^1}^2 + \|u(t_n)\|_{L^4}^4 \rightarrow 0,$$

which contradicts [Lemma 3.5](#) \square

4. BLOW-UP FOR THE FOCUSING CUBIC NLS

In this chapter we prove [Theorem 1.3](#) regarding the NLS [\(1.13\)](#). We first construct a suitable cut-off function localizing in the exterior of a cylinder parallel to the x_3 axis, which we then plug in the virial identities below. This cut-off function will be used also in the proof of [Theorem 1.2](#) regarding the blow-up in finite time for solutions to the GPE, but the estimates we need in order to control some remainders in the presence of the dipolar kernel are much more involved. Therefore we prefer to state the result for the focusing cubic NLS which has its own interest, passing to the analysis of the equation [\(1.2\)](#) in the next chapter. Previous papers in literature about the formation of singularities in finite time for L^2 -supercritical focusing NLS in 3D are [\[12, 15, 17, 20, 27\]](#), besides the already cited [\[16, 23, 25\]](#).

We now introduce some classical virial identities, valid both for [\(1.2\)](#) and [\(1.13\)](#). Let $u(t)$ be a solution to [\(1.2\)](#), which also corresponds to a solution to [\(1.13\)](#) for $\lambda_2 = 0$ and $\lambda_1 < 0$; then we define

$$V_\rho(t) = 2 \int \rho(x) |u(t, x)|^2 dx, \tag{4.1}$$

$\rho(x)$ being a sufficiently smooth function which justifies the following formal and standard computations:

$$\frac{d}{dt} V_\rho(t) = 2\Im \left\{ \int \nabla \rho \cdot \nabla u \bar{u} dx \right\}, \tag{4.2}$$

where we used the equation satisfied by $u(t)$. By using [\(4.2\)](#) and again the equation solved by $u(t)$, we have

$$\begin{aligned} \frac{d^2}{dt^2} V_\rho(t) &= 2 \int (\nabla^2 \rho \cdot \nabla u) \cdot \nabla \bar{u} dx - \frac{1}{2} \int \Delta^2 \rho |u|^2 dx \\ &\quad + \lambda_1 \int \Delta \rho |u|^4 dx - 2\lambda_2 \int \nabla \rho \cdot \nabla (K * |u|^2) |u|^2 dx. \end{aligned}$$

Therefore for the rescaled function $\rho_R(x) := R^2 \rho(x/R)$ we get

$$\begin{aligned} \frac{d^2}{dt^2} V_{\rho_R}(t) &= 2 \int \left(\nabla^2 \rho \left(\frac{x}{R} \right) \cdot \nabla u \right) \cdot \nabla \bar{u} \, dx - \frac{1}{2R^2} \int \Delta^2 \rho \left(\frac{x}{R} \right) |u|^2 \, dx \\ &\quad + \lambda_1 \int \Delta \rho \left(\frac{x}{R} \right) |u|^4 \, dx - 2\lambda_2 R \int \nabla \rho \left(\frac{x}{R} \right) \cdot \nabla (K * |u|^2) |u|^2 \, dx. \end{aligned} \quad (4.3)$$

Let us precisely build the (rescaled) function ρ , which is in particular a radial cut-off function depending only on the two variables $\bar{x} = (x_1, x_2)$:

$$\rho_R(x) = \rho_R(|\bar{x}|) = R^2 \rho(|\bar{x}|/R) = R^2 \psi(|\bar{x}|^2/R^2)$$

where

$$\psi(r) = r - \int_0^r (r-s)\eta(s) \, ds,$$

and the function $\eta : \mathbb{R} \mapsto \mathbb{R}_0^+$ is a function satisfying the following properties: it is a nonnegative, regular function with unitary mean, namely:

$$\begin{cases} \eta \in C_c^\infty(\mathbb{R}; \mathbb{R}_0^+) \\ \eta(s) = 0 \quad \text{for } s \leq 1 \text{ and } s \geq 2 \\ \int_{\mathbb{R}} \eta(s) \, ds = 1 \end{cases} .$$

We have, for $i, j \in \{1, 2\}$ and δ_{ij} being the usual Kronecker symbol,

$$\partial_{x_i} \rho_R(x) = 2x_i \psi'(|\bar{x}|^2/R^2)$$

and

$$\partial_{x_j, x_i}^2 \rho_R(x) = 2\delta_{ij} \psi'(|\bar{x}|^2/R^2) + \frac{4x_i x_j}{R^2} \psi''(|\bar{x}|^2/R^2),$$

hence

$$\begin{aligned} \left(\nabla^2 \rho \left(\frac{x}{R} \right) \nabla u \right) \cdot \nabla \bar{u} &= 2\psi'(|\bar{x}|^2/R^2) |\nabla_{\bar{x}} u|^2 + \frac{4}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2) |\nabla_{\bar{x}} u|^2 \\ &= 2|\nabla_{\bar{x}} u|^2 \left(\psi'(|\bar{x}|^2/R^2) + \frac{2}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2) \right) \end{aligned}$$

and

$$\Delta \rho_R = 4\psi'(|\bar{x}|^2/R^2) + \frac{4}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2).$$

We observe moreover that $\Delta^2 \rho \in L^\infty$ and therefore we can estimate $R^{-2} \int \Delta^2 \rho(x/R) |u|^2 \, dx$ as

$$R^{-2} \int \Delta^2 \rho(x/R) |u|^2 \, dx \leq C(\|\Delta \rho\|_{L^\infty}) R^{-2} \|u(t)\|_{L^2}^2 = CM R^{-2}, \quad (4.4)$$

by using the conservation of the mass.

We compute explicitly the following:

$$-F_R(\bar{x}) := \psi'(|\bar{x}|^2/R^2) + \frac{|\bar{x}|^2}{R^2} \psi''(|\bar{x}|^2/R^2) - 1 = - \int_0^{|\bar{x}|^2/R^2} \eta(s) \, ds - \frac{|\bar{x}|^2}{R^2} \eta(|\bar{x}|^2/R^2) \leq 0. \quad (4.5)$$

We observe that by its construction

$$L^\infty \ni F_R \begin{cases} = 0 & \text{for any } |\bar{x}| \leq R \\ > 0 & \text{for any } |\bar{x}| > R \end{cases}. \quad (4.6)$$

and the boundedness is uniform with respect to $R \geq 1$.

Moreover we consider the not-localized virial identity in the x_3 direction; we namely plug in (4.1) the function $\rho = Ax_3^2$, A being a positive constant, and we simply get,

$$\frac{d^2}{dt^2} V_{Ax_3^2}(t) = 4A \int |\partial_{x_3} u|^2 dx + 2A\lambda_1 \int |u|^4 dx - 4A\lambda_2 \int x_3 \partial_{x_3} (K * |u|^2) |u|^2 dx. \quad (4.7)$$

For $A = 1$ and by plugging $\lambda_2 = 0$ in (4.3), we are going to prove that

$$\begin{aligned} \frac{d^2}{dt^2} V_{\rho_R+x_3^2}(t) &\leq 4 \int |\nabla u|^2 dx + 6\lambda_1 \int |u|^4 dx + \tilde{H}_R(u(t)) \\ &= 4G(u(t)) + \tilde{H}_R(u(t)), \end{aligned} \quad (4.8)$$

where

$$\tilde{H}_R(u(t)) = o_R(1) + o_R(1) \|u(t)\|_{H^1}^2; \quad (4.9)$$

therefore [Lemma 3.6](#), [Corollary 3.7](#) coupled with a convexity argument will yield to the result stated in [Theorem 1.3](#), as long as $R \gg 1$. Note that for $\lambda_2 = 0$ $G(u)$ is reduced to $4 \int |\nabla u|^2 dx + 6\lambda_1 \int |u|^4 dx$.

4.1. Estimate of the remainder $H_R(u(t))$. Once fixed $\lambda_2 = 0$, and for $\lambda_1 < 0$, (4.3) reads

$$\begin{aligned} \frac{d^2}{dt^2} V_{\rho_R}(t) &= 4 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{2}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2) \right) |\nabla_{\bar{x}} u|^2 dx \\ &\quad + \lambda_1 \int \Delta \rho(x/R) |u|^4 dx - \frac{1}{2R^2} \int \Delta^2 \rho(x/R) |u|^2 dx \\ &= 4 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{2}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2) \pm 1 \right) |\nabla_{\bar{x}} u|^2 dx \\ &\quad \pm 4 \int |\partial_{x_3} u|^2 dx - \frac{1}{2R^2} \int \Delta^2 \rho(x/R) |u|^2 dx \\ &\quad + 4\lambda_1 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{|\bar{x}|^2}{R^2} \psi''(|\bar{x}|^2/R^2) \pm \frac{3}{2} \right) |u|^4 dx \end{aligned}$$

and by using the definition of $G(u)$, see (1.11), taking into account that the potential energy $P(u)$ is simply given by $\lambda_1 \int |u|^4 dx$, see (1.10) for $\lambda_2 = 0$, we get

$$\begin{aligned} \frac{d^2}{dt^2} V_{\rho_R}(t) &= 4G(u(t)) - \frac{1}{2R^2} \int \Delta^2 \rho(x/R) |u|^2 dx - 4 \int |\partial_{x_3} u|^2 dx \\ &\quad + 4 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{2}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2) - 1 \right) |\nabla_{\bar{x}} u|^2 dx \\ &\quad + 4\lambda_1 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{|\bar{x}|^2}{R^2} \psi''(|\bar{x}|^2/R^2) - \frac{3}{2} \right) |u|^4 dx \end{aligned}$$

and

$$\frac{d^2}{dt^2} V_{Ax_3^2}(t) = 4A \int |\partial_{x_3} u|^2 dx + 2A\lambda_1 \int |u|^4 dx,$$

therefore by summing up the two terms we get

$$\begin{aligned} \frac{d^2}{dt^2} V_{\rho_R + Ax_3^2}(t) &= 4G(u(t)) - \frac{1}{2R^2} \int \Delta^2 \rho(x/R) |u|^2 dx - 4(1-A) \int |\partial_{x_3} u|^2 dx \\ &\quad + 4 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{2}{R^2} |\bar{x}|^2 \psi''(|\bar{x}|^2/R^2) - 1 \right) |\nabla_{\bar{x}} u|^2 dx \\ &\quad + 4\lambda_1 \int \left(\psi'(|\bar{x}|^2/R^2) + \frac{|\bar{x}|^2}{R^2} \psi''(|\bar{x}|^2/R^2) - \frac{3}{2} + \frac{A}{2} \right) |u|^4 dx \\ &\leq 4G(u(t)) + cR^{-2} \\ &\quad + 4|\lambda_1| \int \left(1 - \psi'(|\bar{x}|^2/R^2) - \frac{|\bar{x}|^2}{R^2} \psi''(|\bar{x}|^2/R^2) \right) |u|^4 dx, \end{aligned} \quad (4.10)$$

where we have set $A = 1$, we have used the definition of $G(u)$, (4.4), and (4.5). Note that, even if it is a simple computation, we preferred to state the virial identity (4.7) with a general constant A to emphasize how the choice $A = 1$ is precisely done to make appear – in the last line of (4.10) – the function F_R defined above. At this point (4.10) reduces to

$$\frac{d^2}{dt^2} V_{\rho_R + x_3^2}(t) \leq 4G(u(t)) + cR^{-2} + 4|\lambda_1| \iint_{|\bar{x}| > R} F_R(\bar{x}) |u|^4 d\bar{x} dx_3.$$

We estimate, in the spirit of Martel [23],

$$\iint F_R(\bar{x}) |u|^4 d\bar{x} dx_3 \leq \int \|F_R |u|^2\|_{L_{\bar{x}}^\infty} \|u\|_{L_{\bar{x}}^2}^2 dx_3 \leq \|u\|_{L_{x_3}^\infty L_{\bar{x}}^2}^2 \int \|F_R |u|^2\|_{L_{\bar{x}}^\infty} dx_3, \quad (4.11)$$

where the norm in the \bar{x} variable are meant in the domain $\{|\bar{x}| \geq R\}$ due to (4.6). We now use the Strauss embedding for a radial function $g(\bar{x})$, $\bar{x} \in \mathbb{R}^2$ and $g \in H^1$, see [9],

$$\|g\|_{L_{\bar{x}}^\infty(|\bar{x}| > R)} \lesssim R^{-1/2} \|g\|_{L_{\bar{x}}^2}^{1/2} \|g\|_{H_{\bar{x}}^1}^{1/2}, \quad (4.12)$$

hence by recalling that F_R is bounded in space uniformly in $R \geq 1$, we obtain, by using (4.12) and the Cauchy-Schwarz's inequality, that

$$\begin{aligned} \int \|F_R |u|^2\|_{L_{\bar{x}}^\infty} dx_3 &\lesssim \int \| |u|^2 \|_{L_{\bar{x}}^\infty(|\bar{x}| \geq R)} dx_3 \lesssim \int \|u\|_{L_{\bar{x}}^\infty(|\bar{x}| \geq R)}^2 dx_3 \\ &\lesssim R^{-1} \left(M + \sqrt{M} \|\nabla_{\bar{x}} u\|_{L^2} \right) \lesssim R^{-1} + R^{-1} \|u\|_{\dot{H}^1}. \end{aligned} \quad (4.13)$$

On the other hand, by calling $g(x_3) = \int_{\mathbb{R}^2} |u|^2(\bar{x}, x_3) d\bar{x}$ we have

$$g(x_3) = \int_{-\infty}^{x_3} \partial_s g(s) ds = 2\Re \int_{-\infty}^{x_3} \left(\int_{\mathbb{R}^2} \bar{u} \partial_s u d\bar{x} \right) ds \leq 2 \int |u| |\nabla u| dx \leq 2\sqrt{M} \|u\|_{\dot{H}^1}$$

and then

$$\|u\|_{L_{x_3}^\infty L_y^2}^2 \lesssim \|u\|_{\dot{H}^1}. \quad (4.14)$$

By glueing up (4.13) and (4.14) we get that (4.11) satisfies

$$\iint F_R |u|^4 d\bar{x} dx_3 \lesssim \iint_{|\bar{x}| \geq R} |u|^4 d\bar{x} dx_3 \lesssim R^{-1} \|u\|_{\dot{H}^1} + R^{-1} \|u\|_{\dot{H}^1}^2 \lesssim R^{-1} + R^{-1} \|u\|_{\dot{H}^1}^2. \quad (4.15)$$

We conclude that for some $\alpha > 0$

$$\frac{d^2}{dt^2} V_{\rho_R + x_3^2}(t) \lesssim G(u(t)) + R^{-1} + R^{-1} \|u(t)\|_{\dot{H}^1}^2 \lesssim -\frac{\alpha}{2} \|u(t)\|_{\dot{H}^1}^2 \lesssim -1,$$

where in last step we used [Lemma 3.5](#), [Lemma 3.6](#), [Corollary 3.7](#) and we have chosen $R \gg 1$. We can eventually conclude that $T_{max} < +\infty$ by a convexity argument. Indeed, it implies that there exists a time T_0 such that

$$\int (\rho_R + x_3^2) |u(t)|^2 dx \rightarrow 0 \quad \text{as } t \rightarrow T_0.$$

We observe the following: by using the Weyl-Heisenberg's inequality $\|xf\|_{L^2} \|f\|_{\dot{H}^1} \gtrsim \|f\|_{L^2}^2$ in 1D we get

$$\|u_0\|_{L^2}^2 = \|u(t)\|_{L^2}^2 = \int_{\mathbb{R}^2} \int_{\mathbb{R}} |u(t)|^2 dx_3 d\bar{x} \lesssim \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}} |\partial_{x_3} u(t)|^2 dx_3 \right)^{1/2} \left(\int_{\mathbb{R}} x_3^2 |u(t)|^2 dx_3 \right)^{1/2} d\bar{x}$$

and by using the Cauchy-Schwarz's inequality we conclude with the estimate

$$\|x_3 u(t)\|_{L^2} \|\partial_{x_3} u(t)\|_{L^2} \gtrsim \|u_0\|_{L^2}^2 > 0.$$

Then $\|u(t)\|_{\dot{H}^1} \rightarrow \infty$ as $t \rightarrow T_0$. The proof of [Theorem 1.3](#) is therefore concluded.

5. BLOW-UP FOR THE DIPOLAR GPE

This last chapter is devoted to the proof of [Theorem 1.2](#). With respect to the NLS equation, when dealing with the equation (1.2) governing a Dipolar BEC, we get an additional term to be estimated in the sum of the virial identities (4.3) and (4.7), namely the sum of the two terms involving λ_2 . It is worth mentioning that all the terms that we have shown to be small in the NLS case will remain the same. What we are going to prove is that for the Dipolar GPE (1.2) we have a virial-type estimate of the form (cf. with (4.8), (4.9))

$$\frac{d^2}{dt^2} V_{\rho_R + x_3^2}(t) \leq 4 \int |\nabla u|^2 dx + 6\lambda_1 \int |u|^4 dx + H_R(u(t)) \quad (5.1)$$

where

$$H_R = \tilde{H}_R - 2\lambda_2 \left(\int \nabla \rho_R \cdot \nabla (K * |u|^2) |u|^2 dx + 2 \int x_3 \partial_{x_3} (K * |u|^2) |u|^2 dx \right). \quad (5.2)$$

As already proved in [Section 4](#), see [\(4.15\)](#),

$$\tilde{H}_R(u(t)) = o_R(1) + o_R(1)\|u(t)\|_{\dot{H}^1}^2, \quad (5.3)$$

so we are going to show that

$$\mathcal{V} + \mathcal{W} := \int \nabla \rho_R \cdot \nabla (K * |u|^2) |u|^2 dx + 2 \int x_3 \partial_{x_3} (K * |u|^2) |u|^2 dx \quad (5.4)$$

will contribute for a term $-3 \int (K * |u|^2) |u|^2 dx$ that will yield to the quantity $4G(u)$ (see the definition [\(1.11\)](#)) when summed to the first two terms in the r.h.s. of [\(5.1\)](#), plus a remainder term which must be proved to be small, again of the type $o_R(1) + o_R(1)\|u\|_{\dot{H}^1}^2$.

5.1. Estimate of the remainder $H_R(u(t))$. The \mathcal{V} term. By its definition, we get that the function ρ_R fulfils

$$\partial_{x_i} \rho_R(x) = 2x_i \psi'(|\bar{x}|^2/R^2) = 2\bar{x} \left(1 - \int_0^{|\bar{x}|^2/R^2} \eta(s) ds \right) = \begin{cases} 2\bar{x} & \text{if } |\bar{x}|^2/R^2 \leq 1 \\ 0 & \text{if } |\bar{x}|^2/R^2 > 2 \end{cases},$$

hence $\text{supp } \nabla_{\bar{x}} \rho_R$ – where supp stands for the support of a function – is contained in the cylinder of radius $\sqrt{2}R$.

We split the function u by partitioning the whole space in the region inside and the region outside a cylinder, namely we write $u = u_i + u_o$ where

$$u_i = \mathbf{1}_{\{|\bar{x}| \leq 4R\}} u \quad \text{and} \quad u_o = \mathbf{1}_{\{|\bar{x}| \geq 4R\}} u.$$

Since $\text{supp } \nabla \rho_R \cap \text{supp } u_o = \emptyset$ we get

$$\begin{aligned} \mathcal{V} &:= \int \nabla_{\bar{x}} \rho_R \cdot \nabla_{\bar{x}} (K * |u|^2) |u|^2 dx \\ &= \int \nabla_{\bar{x}} \rho_R \cdot \nabla_{\bar{x}} (K * |u_o|^2) |u_i|^2 dx - \int \nabla_{\bar{x}} \rho_R \cdot \nabla_{\bar{x}} (K * |u_i|^2) |u_i|^2 dx \\ &= \mathcal{V}_{oi} + \mathcal{V}_{ii}. \end{aligned}$$

Estimate for the term \mathcal{V}_{oi} . The term \mathcal{V}_{oi} can be estimated in this way: by integrating by parts,

$$\begin{aligned} \mathcal{V}_{oi} &= \int \nabla_{\bar{x}} \rho_R \cdot \nabla_{\bar{x}} (K * |u_o|^2) |u_i|^2 dx \\ &= - \int \Delta_{\bar{x}} \rho_R (K * |u_o|^2) |u_i|^2 dx - \int \nabla_{\bar{x}} \rho_R \cdot \nabla_{\bar{x}} (|u_i|^2) (K * |u_o|^2) dx = \mathcal{V}'_{oi} + \mathcal{V}''_{oi} \end{aligned}$$

and, by using [Lemma 2.5](#), in particular the pointwise estimate [\(2.8\)](#), we obtain

$$|\mathcal{V}'_{oi}| \lesssim \int \mathbf{1}_{\{|\bar{x}| \leq \sqrt{2}R\}} |K * |u_o|^2| |u_i|^2 dx \lesssim R^{-3} \|u_o\|_{L^2}^2 \|u_i\|_{L^2}^2 \lesssim R^{-3} \|u\|_{L^2}^4 \lesssim R^{-3}.$$

Similarly, by using that $|\nabla_{\bar{x}}\rho_R| \lesssim R$ on its support in conjunction with the Cauchy-Schwarz's inequality, we get

$$\begin{aligned} |\mathcal{V}'_{oi}| &\lesssim \int \mathbf{1}_{\{|\bar{x}| \leq \sqrt{2}R\}} |\nabla_{\bar{x}}\rho_R| \left| K * |u_o|^2 \right| |u_i| |\nabla_{\bar{x}}u_i| dx \\ &\lesssim R \int \mathbf{1}_{\{|\bar{x}| \leq \sqrt{2}R\}} \left| K * |u_o|^2 \right| |u_i| |\nabla_{\bar{x}}u_i| dx \lesssim R^{-2} \|u_o\|_{L^2}^2 \|u_i\|_{L^2} \|u_i\|_{\dot{H}^1} \lesssim R^{-2} \|u\|_{\dot{H}^1} \end{aligned}$$

and then the estimate for \mathcal{V}_{oi} is concluded by summing up the two bounds above:

$$|\mathcal{V}_{oi}| \lesssim R^{-2} \|u\|_{\dot{H}^1} + R^{-3}. \quad (5.5)$$

Estimate for the term \mathcal{V}_{ii} . We analyse the term \mathcal{V}_{ii} . We do a further splitting and we introduce another localization function. By setting up $\tilde{\rho}_R = \rho_R - |\bar{x}|^2$ we can write

$$\begin{aligned} \mathcal{V}_{ii} &= \int \nabla_{\bar{x}}\rho_R \cdot \nabla_{\bar{x}} \left(K * |u_i|^2 \right) |u_i|^2 dx \\ &= \int \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(K * |u_i|^2 \right) |u_i|^2 dx + 2 \int \bar{x} \cdot \nabla_{\bar{x}} \left(K * |u_i|^2 \right) |u_i|^2 dx = \mathcal{V}'_{ii} + \mathcal{V}''_{ii}. \end{aligned} \quad (5.6)$$

We further localize the function u_i by splitting again u_i as $u_i = w_{i,i} + w_{i,o}$, where

$$w_{i,i} = \mathbf{1}_{\{|\bar{x}| \leq R/10\}} u_i \quad \text{and} \quad w_{i,o} = \mathbf{1}_{\{|\bar{x}| \geq R/10\}} u_i = \mathbf{1}_{\{R/10 \leq |\bar{x}| \leq 4R\}} u.$$

By using the fact that $\text{supp } \nabla_{\bar{x}}\tilde{\rho}_R$ is contained in $\{|\bar{x}| \geq R\}$, then $\text{supp } \nabla_{\bar{x}}\tilde{\rho}_R \cap \{|\bar{x}| \leq R/10\} = \emptyset$, we write

$$\begin{aligned} \mathcal{V}'_{ii} &= \int \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(K * |w_{i,i}|^2 \right) |w_{i,o}|^2 dx \\ &\quad + \int \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(K * |w_{i,o}|^2 \right) |w_{i,o}|^2 dx = \mathcal{A} + \mathcal{B}. \end{aligned}$$

Now, similarly to the term \mathcal{V}_{oi} , by integrating by parts and by using in this case [Lemma 2.6](#), and precisely the pointwise estimate [\(2.18\)](#), we have

$$\begin{aligned} \mathcal{A} &= \int \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(K * |w_{i,i}|^2 \right) |w_{i,o}|^2 dx = - \int_{\{R \leq |\bar{x}| \leq 4R\}} \Delta_{\bar{x}}\tilde{\rho}_R \left(K * |w_{i,i}|^2 \right) |w_{i,o}|^2 dx \\ &\quad - \int_{\{R \leq |\bar{x}| \leq 4R\}} \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(|w_{i,o}|^2 \right) \left(K * |w_{i,i}|^2 \right) dx \\ &\lesssim R^{-3} \|w_{i,i}\|_{L^2}^2 \|w_{i,o}\|_{L^2}^2 + R^{-2} \|w_{i,i}\|_{L^2}^2 \|w_{i,o}\|_{L^2} \|w_{i,o}\|_{\dot{H}^1} \lesssim R^{-3} + R^{-2} \|u\|_{\dot{H}^1}. \end{aligned} \quad (5.7)$$

Note that we used the following facts: $|\Delta_{\bar{x}}\tilde{\rho}_R| \lesssim 1$, and by recalling that $u_i = \mathbf{1}_{\{|\bar{x}| \leq 4R\}} u$, we infer that $|\nabla\tilde{\rho}_R| \lesssim R$ on $\{|\bar{x}| \leq 4R\}$.

Estimate for the term \mathcal{B} . Hence it remains to prove a suitable estimate for the term

$$\mathcal{B} = \int_{\{R \leq |\bar{x}| \leq \sqrt{2}R\}} \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(K * |w_{i,o}|^2 \right) |w_{i,o}|^2 dx. \quad (5.8)$$

By setting $g = |w_{i,o}|^2$ and making use of the Plancherel identity we get

$$\int \nabla_{\bar{x}}\tilde{\rho}_R \cdot \nabla_{\bar{x}} \left(K * |w_{i,o}|^2 \right) |w_{i,o}|^2 dx = \int \widehat{g \nabla_{\bar{x}}\tilde{\rho}}(\xi) \cdot \bar{\xi} \hat{K} \bar{g} d\xi.$$

By recalling the precise expression for the Fourier transform of the dipolar kernel, see (1.8), we note that \hat{K} is a linear combination of the symbols defining the square of the j -th Riesz transform \mathcal{R}_j^2 :

$$\hat{K}(\xi) = \frac{4\pi}{3} \frac{2\xi_3^2 - \xi_2^2 - \xi_1^2}{|\xi|^2} = \sum_{j=1}^3 \alpha_j \frac{\xi_j^2}{|\xi|^2}.$$

Consider therefore the generic term $\int \widehat{\nabla_{\bar{x}} \tilde{\rho}_R g}(\xi) \cdot \bar{\xi} \frac{\xi_j^2}{|\xi|^2} \bar{g} d\xi$;

$$\begin{aligned} \int g \widehat{\nabla_{\bar{x}} \tilde{\rho}_R}(\xi) \cdot \bar{\xi} \frac{\xi_j^2}{|\xi|^2} \bar{g}(\xi) d\xi &= \int (\widehat{\nabla_{\bar{x}} \tilde{\rho}_R} * \hat{g})(\xi) \cdot \bar{\xi} \frac{\xi_j^2}{|\xi|^2} \bar{g}(\xi) d\xi \\ &= \iint \hat{g}(\eta) \widehat{\nabla_{\bar{x}} \tilde{\rho}_R}(\xi - \eta) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} \pm \frac{\eta_j \bar{\eta}}{|\eta|} \right) \frac{\xi_j}{|\xi|} \hat{g}(\xi) d\eta d\xi \\ &= \int \nabla_{\bar{x}} \tilde{\rho}_R \cdot \nabla_{\bar{x}} (\mathcal{R}_j g)(x) \mathcal{R}_j \bar{g}(x) dx \\ &+ \iint \hat{g}(\eta) \widehat{\nabla_{\bar{x}} \tilde{\rho}_R}(\xi - \eta) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} - \frac{\eta_j \bar{\eta}}{|\eta|} \right) \frac{\xi_j}{|\xi|} \hat{g}(\xi) d\eta d\xi \\ &= -\frac{1}{2} \int \Delta_{\bar{x}} \tilde{\rho}_R |\mathcal{R}_j g(x)|^2 dx \\ &+ \iint \hat{g}(\eta) \widehat{\nabla_{\bar{x}} \tilde{\rho}_R}(\xi - \eta) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} - \frac{\eta_j \bar{\eta}}{|\eta|} \right) \frac{\xi_j}{|\xi|} \hat{g}(\xi) d\eta d\xi. \end{aligned} \tag{5.9}$$

The first term in the r.h.s. of (5.9) is simply estimated by

$$\|u\|_{L^4(|\bar{x}| \geq R/10)}^4 \lesssim R^{-1} \|u\|_{\dot{H}^1}^2 \tag{5.10}$$

due to the $L^2 \mapsto L^2$ continuity property of the Riesz transform and (4.15). For the second term in the r.h.s. of (5.9) we proceed in this way. First of all suppose that $j = 1, 2$; then

$$\begin{aligned} &\iint \hat{g}(\eta) \widehat{\nabla_{\bar{x}} \tilde{\rho}_R}(\xi - \eta) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} - \frac{\eta_j \bar{\eta}}{|\eta|} \right) \frac{\xi_j}{|\xi|} \hat{g}(\xi) d\eta d\xi \\ &= \int \frac{\xi_j}{|\xi|} \hat{g}(\xi) \iint \hat{g}(\bar{\eta}, \eta_3) \delta(\xi_3 - \eta_3) \widehat{\tilde{\rho}_R}(\bar{\xi} - \bar{\eta}) (\bar{\xi} - \bar{\eta}) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} - \frac{\eta_j \bar{\eta}}{|\eta|} \right) d\bar{\eta} d\eta_3 d\xi \\ &= \int \frac{\xi_j}{|\xi|} \hat{g}(\xi) \int \hat{g}(\bar{\eta}, \xi_3) \underbrace{\widehat{\tilde{\rho}_R}(\bar{\xi} - \bar{\eta}) (\bar{\xi} - \bar{\eta}) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} - \frac{\eta_j \bar{\eta}}{\sqrt{|\bar{\eta}|^2 + \xi_3^2}} \right)}_{\vec{F}_j} d\bar{\eta} d\xi. \end{aligned}$$

If instead $j = 3$ we have

$$\begin{aligned}
& \iint \hat{g}(\eta) \widehat{\nabla_{\bar{x}} \tilde{\rho}_R}(\xi - \eta) \cdot \left(\frac{\xi_3 \bar{\xi}}{|\xi|} - \frac{\eta_3 \bar{\eta}}{|\eta|} \right) \frac{\xi_3}{|\xi|} \hat{g}(\xi) \, d\eta \, d\xi \\
&= \int \frac{\xi_3}{|\xi|} \hat{g}(\xi) \iint \hat{g}(\bar{\eta}, \eta_3) \delta(\xi_3 - \eta_3) \widehat{\tilde{\rho}_R}(\bar{\xi} - \bar{\eta}) (\bar{\xi} - \bar{\eta}) \cdot \left(\frac{\xi_3 \bar{\xi}}{|\xi|} - \frac{\eta_3 \bar{\eta}}{|\eta|} \right) \, d\bar{\eta} \, d\eta_3 \, d\xi \\
&= \int \frac{\xi_3}{|\xi|} \hat{g}(\xi) \int \hat{g}(\bar{\eta}, \xi_3) \widehat{\tilde{\rho}_R}(\bar{\xi} - \bar{\eta}) (\bar{\xi} - \bar{\eta}) \cdot \underbrace{\left(\frac{\xi_3 \bar{\xi}}{|\xi|} - \frac{\xi_3 \bar{\eta}}{\sqrt{|\bar{\eta}|^2 + \xi_3^2}} \right)}_{\vec{F}_3} \, d\bar{\eta} \, d\xi.
\end{aligned}$$

We notice that the Jacobian of both

$$\vec{F}_j(\bar{v}) = \frac{v_j \bar{v}}{\sqrt{|\bar{v}|^2 + \xi_3^2}} = \left(\frac{v_j v_1}{\sqrt{|\bar{v}|^2 + \xi_3^2}}, \frac{v_j v_2}{\sqrt{|\bar{v}|^2 + \xi_3^2}} \right), \quad \bar{v} = (v_1, v_2) \in \mathbb{R}^2, \quad j = 1, 2$$

and

$$\vec{F}_3(\bar{v}) = \frac{\xi_3 \bar{v}}{\sqrt{|\bar{v}|^2 + \xi_3^2}} = \left(\frac{\xi_3 v_1}{\sqrt{|\bar{v}|^2 + \xi_3^2}}, \frac{\xi_3 v_2}{\sqrt{|\bar{v}|^2 + \xi_3^2}} \right), \quad \bar{v} = (v_1, v_2) \in \mathbb{R}^2$$

is uniformly bounded, namely $|J_{\vec{F}_j}(\bar{v})| \lesssim 1$ for any $\bar{v} \in \mathbb{R}^2$, for $j = 1, 2, 3$; therefore we can bound the last term in the r.h.s. of (5.9) by

$$\begin{aligned}
& \int_{\mathbb{R}^3} |\hat{g}(\xi)| \int_{\mathbb{R}^2} |\hat{g}(\bar{\eta}, \xi_3)| \left| \widehat{\tilde{\rho}_R}(\bar{\xi} - \bar{\eta}) \right| |\bar{\xi} - \bar{\eta}|^2 \, d\bar{\eta} \, d\xi \\
& \leq \int_{\mathbb{R}^3} |\hat{g}(\xi)| \int_{\mathbb{R}^2} |\hat{g}(\bar{\eta}, \xi_3)| \left| \widehat{\Delta_{\bar{x}} \rho_R}(\bar{\xi} - \bar{\eta}) \right| \, d\bar{\eta} \, d\xi \\
& + \int_{\mathbb{R}^3} |\hat{g}(\xi)| \int_{\mathbb{R}^2} |\hat{g}(\bar{\eta}, \xi_3)| \left| \widehat{\Delta_{\bar{x}} |\bar{x}|^2}(\bar{\xi} - \bar{\eta}) \right| \, d\bar{\eta} \, d\xi \tag{5.11} \\
& = \int_{\mathbb{R}^3} |\hat{g}(\xi)| (|\hat{g}(\cdot, \xi_3)| * h_R)(\bar{\xi}) \, d\bar{\xi} + \int_{\mathbb{R}^3} |\hat{g}(\xi)| (|\hat{g}(\cdot, \xi_3)| * 4\delta)(\bar{\xi}) \, d\bar{\xi} \\
& = \int_{\mathbb{R}^3} |\hat{g}(\xi)| (|\hat{g}(\cdot, \xi_3)| * h_R)(\bar{\xi}) \, d\bar{\xi} + 4 \int_{\mathbb{R}^3} |\hat{g}(\xi)|^2 \, d\xi
\end{aligned}$$

where we defined $h_R = \left| \widehat{\Delta_{\bar{x}} \rho_R} \right|$. We continue in this way: in the first term in the r.h.s. of (5.11) we first apply the Cauchy-Schwarz's inequality and then the Young's inequality for convolutions with respect to $d\bar{\xi}$, and eventually the Cauchy-Schwarz's inequality with respect to $d\xi_3$ to obtain

$$\begin{aligned}
\int_{\mathbb{R}^3} |\hat{g}(\xi)| (|\hat{g}(\cdot, \xi_3)| * h_R)(\bar{\xi}) \, d\bar{\xi} &= \int_{\mathbb{R}} \int_{\mathbb{R}^2} |\hat{g}(\bar{\xi}, \xi_3)| (|\hat{g}(\cdot, \xi_3)| * h_R)(\bar{\xi}) \, d\bar{\xi} \, d\xi_3 \\
&\leq \int_{\mathbb{R}} \|\hat{g}(\cdot, \xi_3)\|_{L_{\bar{\xi}}^2} \| |\hat{g}(\cdot, \xi_3)| * h_R \|_{L_{\bar{\xi}}^2} \, d\xi_3 \\
&\leq \|h_R\|_{L_{\bar{\xi}}^1} \int_{\mathbb{R}} \|\hat{g}(\cdot, \xi_3)\|_{L_{\bar{\xi}}^2}^2 \, d\xi_3 = \|h_R\|_{L_{\bar{\xi}}^1} \|\hat{g}\|_{L^2}^2 = \|h_R\|_{L_{\bar{\xi}}^1} \|g\|_{L^2}^2,
\end{aligned}$$

where in the last step we used the Plancherel identity. Since by definition the L^1 -norm of h_R is the L^1 -norm of the Fourier transform of $\Delta_{\bar{x}}\rho_R$, and the latter is a smooth and compactly supported function by its very construction, then its Fourier transform is still in L^1_{ξ} uniformly in R . Hence the estimate for the last integral in the r.h.s. of (5.9), recalling that $g = |w_{i,o}|^2 = |\mathbf{1}_{\{|\bar{x}| \geq R/10\}}|^2$, can be concluded with

$$\iint \widehat{g}(\eta) \widehat{\nabla_{\bar{x}}\rho_R}(\xi - \eta) \cdot \left(\frac{\xi_j \bar{\xi}}{|\xi|} - \frac{\eta_j \bar{\eta}}{|\eta|} \right) \frac{\xi_j}{|\xi|} \widehat{g}(\xi) d\eta d\xi \lesssim \|u\|_{L^4(|\bar{x}| \geq R/10)}^4 \lesssim R^{-1} \|u\|_{\dot{H}^1}^2, \quad (5.12)$$

where we used (4.15) for the last inequality. By summing up (5.10) and (5.12) we give the bound for the whole \mathcal{B} term in (5.8):

$$\mathcal{B} \lesssim R^{-1} \|u\|_{\dot{H}^1}^2. \quad (5.13)$$

The estimate (5.13) in conjunction with (5.7) conclude the estimate for \mathcal{V}'_{ii} :

$$\mathcal{V}'_{ii} \lesssim R^{-1} \|u\|_{\dot{H}^1}^2. \quad (5.14)$$

To end up with the full estimate leading to the proof of the main theorem, we are left to study the remaining term \mathcal{V}''_{ii} together with the term \mathcal{W} as defined in (5.4).

5.2. Estimate of the remainder $H_R(u(t))$. The $\mathcal{V}''_{ii} + \mathcal{W}$ term. In order to conclude the estimate for the term $\mathcal{V} + \mathcal{W}$, we are left to control the last term \mathcal{V}''_{ii} coming from the splitting in (5.6), and the term \mathcal{W} , see (5.4), that we did not handle so far. A straightforward computation gives

$$\begin{aligned} \mathcal{V}''_{ii} + \mathcal{W} &= 2 \int \bar{x} \cdot \nabla_{\bar{x}} \left(K * |u_i|^2 \right) |u_i|^2 dx + 2 \int x_3 \partial_{x_3} \left(K * |u|^2 \right) |u|^2 dx \\ &= 2 \int x \cdot \nabla \left(K * |u_i|^2 \right) |u_i|^2 dx + 2 \int x_3 \partial_{x_3} \left(K * |u_i|^2 \right) |u_o|^2 dx \\ &\quad + 2 \int x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_i|^2 dx + 2 \int x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_o|^2 dx \\ &= -3 \int \left(K * |u_i|^2 \right) |u_i|^2 dx + 2 \int x_3 \partial_{x_3} \left(K * |u_i|^2 \right) |u_o|^2 dx \\ &\quad + 2 \int x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_i|^2 dx + 2 \int x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_o|^2 dx \end{aligned} \quad (5.15)$$

where we used the identity (A.2). By means of (A.4) with $f = g = |u_o|^2$ we obtain

$$2 \int x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_o|^2 dx = - \int \left(K * |u_o|^2 \right) |u_o|^2 dx - \int \xi_3 (\partial_{\xi_3} \hat{K}) |\widehat{|u_o|^2}| |\widehat{|u_o|^2}| d\xi \quad (5.16)$$

while, again by means of (A.4), we write

$$\begin{aligned} &2 \int x_3 \partial_{x_3} \left(K * |u_i|^2 \right) |u_o|^2 dx + 2 \int x_3 \partial_{x_3} \left(K * |u_o|^2 \right) |u_i|^2 dx \\ &= -2 \int \left(K * |u_i|^2 \right) |u_o|^2 dx - 2 \int \xi_3 (\partial_{\xi_3} \hat{K}) |\widehat{|u_i|^2}| |\widehat{|u_o|^2}| d\xi. \end{aligned} \quad (5.17)$$

We explicitly write $\xi_3 \partial_{\xi_3} \widehat{K}$ and we observe that is bounded:

$$\xi_3 \partial_{\xi_3} \widehat{K} = 8\pi \frac{\xi_3^2 (\xi_1^2 + \xi_2^2)}{|\xi|^4} \in L_\xi^\infty; \quad (5.18)$$

hence (5.16) is simply estimated by

$$2 \int x_3 \partial_{x_3} (K * |u_o|^2) |u_o|^2 dx \lesssim \|u_o\|_{L^4}^4 \lesssim R^{-1} \|u\|_{\dot{H}^1}^2$$

by using the $L^2 \mapsto L^2$ continuity of the dipolar kernel for the first integral, while for the second integral we used the boundedness property (5.18) together with the Plancherel identity, and (4.15).

It remains to handle (5.17). First of all we note that

$$\xi_3 \partial_{\xi_3} \widehat{K} = 8\pi \frac{\xi_3^2 (\xi_1^2 + \xi_2^2)}{|\xi|^4} = 8\pi \left(\frac{\xi_3^2}{|\xi|^2} - \frac{\xi_3^4}{|\xi|^4} \right) = 8\pi \widehat{\mathcal{R}}_3^2 - 8\pi \widehat{\mathcal{R}}_3^4,$$

where, with abuse of notation, we write $\widehat{\mathcal{R}}_3^2$ and $\widehat{\mathcal{R}}_3^4$ to denote the symbols of \mathcal{R}_3^2 and \mathcal{R}_3^4 , respectively. This in turn implies that the r.h.s. of (5.17) can be rewritten, for some constants $\alpha_{1,2,3}$ as

$$-16\pi \int (\mathcal{R}_3^4(|u_i|^2)) |u_o|^2 dx + \sum_{j=1}^3 \alpha_j \int (\mathcal{R}_j^2(|u_i|^2)) |u_o|^2 dx.$$

By splitting $u_i = w_{ii} + w_{io}$ with $w_{ii} = \mathbf{1}_{\{|\bar{x}| \leq R/10\}} u_i = \mathbf{1}_{\{|\bar{x}| \leq R/10\}} u$ and $w_{io} = \mathbf{1}_{\{|\bar{x}| \geq R/10\}} u_i = \mathbf{1}_{\{R/10 \leq |\bar{x}| \leq 4R\}} u$ we decompose each of the terms in the sum involving \mathcal{R}_j^2 's as

$$\int (\mathcal{R}_j^2(|u_i|^2)) |u_o|^2 dx = \int (\mathcal{R}_j^2(|w_{ii}^2|)) |u_o|^2 dx + \int (\mathcal{R}_j^2(|w_{io}^2|)) |u_o|^2 dx$$

and by using the Cauchy-Schwarz's inequality, the $L^2 \mapsto L^2$ continuity of the Riesz transform, the localization properties of w_{io} and u_o , and (4.15) we obtain

$$\int (\mathcal{R}_j^2(|w_{io}^2|)) |u_o|^2 dx \lesssim \|w_{io}\|_{L^4}^2 \|u_o\|_{L^4}^2 \lesssim \|u\|_{L^4(|\bar{x}| \geq R/10)}^4 \lesssim R^{-1} \|u\|_{\dot{H}^1}^2. \quad (5.19)$$

By using (2.18) we estimate

$$\int (\mathcal{R}_j^2(|w_{ii}^2|)) |u_o|^2 dx \lesssim R^{-3} \|w_{ii}\|_{L^1} \|u_o\|_{L^2}^2 \lesssim R^{-3}.$$

With the same decomposition of the function u_i we separate

$$\int (\mathcal{R}_3^4(|u_i|^2)) |u_o|^2 dx = \int (\mathcal{R}_3^4(|w_{ii}^2|)) |u_o|^2 dx + \int (\mathcal{R}_3^4(|w_{io}^2|)) |u_o|^2 dx;$$

then, similarly to (5.19) we have

$$\int (\mathcal{R}_3^4(|w_{io}^2|)) |u_o|^2 dx \lesssim R^{-1} \|u\|_{\dot{H}^1}^2,$$

while by using [Theorem 2.2](#) we can bound

$$\int \left(\mathcal{R}_3^4(|w_{ii}|^2) \right) |u_o|^2 dx \lesssim R^{-1} \|w_{ii}\|_{L^2}^2 \|u_o\|_{L^2}^2 \lesssim R^{-1}.$$

The remaining term in [\(5.15\)](#) is $-3 \int (K * |u_i|^2) |u_i|^2 dx$. By adding and subtracting u to u_i we can infer, by similar computations done before, that

$$\begin{aligned} -3 \int \left(K * |u_i|^2 \right) |u_i|^2 dx &= -3 \int \left(K * |u|^2 \right) |u|^2 dx + \epsilon(R, u) \\ \epsilon(R, u) &\lesssim o_R(1) + R^{-1} \|u\|_{\dot{H}^1}^2. \end{aligned} \tag{5.20}$$

5.3. Proof of [Theorem 1.2](#). We can now summarize all the previous contributions towards the final step of the proof. We have, starting from [\(5.1\)](#)-[\(5.2\)](#), that

$$\begin{aligned} \frac{d^2}{dt^2} V_{\rho_R + x_3^2}(t) &\leq 4 \int |\nabla u|^2 dx + 6\lambda_1 \int |u|^4 dx + \tilde{H}_R(u(t)) - 2\lambda_2(\mathcal{V} + \mathcal{W}) \\ &= 4 \int |\nabla u|^2 dx + 6\lambda_1 \int |u|^4 dx + \tilde{H}_R(u(t)) - 2\lambda_2(\mathcal{V}_{oi} + \mathcal{V}'_{ii} + \mathcal{V}''_{ii} + \mathcal{W}). \end{aligned}$$

Thanks to the estimates [\(5.5\)](#), [\(5.3\)](#), [\(5.14\)](#), and [\(5.20\)](#) which provide a control for the terms \mathcal{V}_{oi} , $\tilde{H}_R(u(t))$, \mathcal{V}'_{ii} , and $\mathcal{V}''_{ii} + \mathcal{W}$, respectively, we end up with

$$\frac{d^2}{dt^2} V_{\rho_R + x_3^2}(t) \leq 4G(u(t)) + o_R(1) + o_R(1) \|u(t)\|_{\dot{H}^1}^2.$$

Therefore by means of [Lemma 3.5](#), [Lemma 3.6](#), and [Corollary 3.7](#), provided $R \gg 1$ we infer that $T_{min} = T_{max} = +\infty$ via a convexity argument. The proof of [Theorem 1.2](#) is now complete.

APPENDIX A. IDENTITIES FOR CONVOLUTION WITH THE DIPOLAR KERNEL

We recall here some useful identities often used along the proofs contained in the paper. They are basically versions for the dipolar kernel K of some identities found by Cipolatti, see [\[10\]](#) in the context of the Davey-Stewartson system. For our equation [\(1.2\)](#) the crucial identity is

$$\xi \cdot \nabla_{\xi} \hat{K} = 0. \tag{A.1}$$

We begin with the following.

Lemma A.1. *Given a (smooth) functions f the following identity holds true:*

$$2 \int x \cdot \nabla (K * f) f dx = -3 \int (K * f) f dx \tag{A.2}$$

Proof. The identity is proved by means of some properties of the dipolar kernel and its explicit representation in the frequencies space. We start by considering two (smooth) functions f, g and we write

$$\begin{aligned} \int x \cdot \nabla (K * f) g dx &= \int x \cdot (\nabla K * f) g dx = \int x \cdot \left(\int \nabla K(x-y) f(y) dy \right) g(x) dx \\ &= \iint (x-y) \cdot \nabla K(x-y) f(y) g(x) dy dx + \int \left(\int y \cdot \nabla K(x-y) f(y) dy \right) g(x) dx, \end{aligned}$$

and since the dipolar kernel is an odd function, we continue as follows:

$$\begin{aligned}
\int x \cdot \nabla (K * f) g \, dx &= \int ((x \cdot \nabla K) * f) g - \iint y \cdot \nabla K(y-x) f(y) g(x) \, dy \, dx \\
&= \int ((x \cdot \nabla K) * f) g \, dx - \iint x \cdot \nabla K(x-y) f(x) g(y) \, dx \, dy \\
&= \int ((x \cdot \nabla K) * f) g \, dx - \int x \cdot (\nabla K * g) f \, dx
\end{aligned}$$

hence, by using the Plancherel identity and (A.1) we conclude that

$$\begin{aligned}
\int x \cdot \nabla (K * f) g \, dx + \int x \cdot \nabla (K * g) f \, dx &= \int ((x \cdot \nabla K) * f) g \, dx \\
&= - \int \operatorname{div}_\xi (\xi \hat{K}) \hat{f} \hat{g} = -3 \int (K * f) g \, dx.
\end{aligned} \tag{A.3}$$

By plugging $f = g$ in (A.3) we prove (A.2). \square

A consequence of the previous identities is the following.

Lemma A.2. *Given two (smooth) functions f, g the following identity holds true:*

$$\int x_3 \partial_{x_3} (K * f) g \, dx + \int x_3 \partial_{x_3} (K * g) f \, dx = - \int (K * f) g \, dx - \int \xi_3 (\partial_{\xi_3} \hat{K}) \hat{f} \hat{g} \, d\xi. \tag{A.4}$$

Proof. Similarly to the previous proof we have:

$$\begin{aligned}
\int x_3 \partial_{x_3} (K * f) g \, dx + \int x_3 \partial_{x_3} (K * g) f \, dx &= \int ((x_3 \partial_{x_3} K) * f) g \, dx = - \int \partial_{\xi_3} (\xi_3 \hat{K}) \hat{f} \hat{g} \, d\xi \\
&= - \int (K * f) g \, dx - \int \xi_3 (\partial_{\xi_3} \hat{K}) \hat{f} \hat{g} \, d\xi.
\end{aligned}$$

\square

APPENDIX B. PROOF OF (2.5)

We give here the proof of the fact that there exists an integer $M \geq 1$ and M pairs of polynomial with nonnegative coefficients $(\tilde{q}_k, q_k)_{k \in \{1, \dots, M\}}$ satisfying the following: $\deg(q_k) \geq 1$ and

$$\left| \partial_{x_i}^4 \left(\frac{1}{|x-y|} \sin(\tau^{1/4}|x-y|) \right) \right| \lesssim \sum_{k=1}^M \frac{\tilde{q}_k(\tau^{1/4})}{q_k(|x-y|)}. \tag{B.1}$$

As the derivative is invariant under translations and by defining $c = \tau^{1/4}$, we can reduce everything to the estimate of $\partial_{x_i}^4 (|x|^{-1} \sin(c|x|))$. By setting $f(r) = r^{-1} \sin(cr)$ and $g(x) = |x|$ we can see

$$|x|^{-1} \sin(c|x|) = (f \circ g)(x),$$

and without loss of generality we assume $i = 3$. Then we see g as a function of x_3 alone, i.e. $g(x_3) = \sqrt{x_3^2 + x_2^2 + x_1^2}$. We first collect some identities.

$$\begin{aligned} f'(r) &= cr^{-1} \cos(cr) - r^{-2} \sin(cr) \\ f''(r) &= -c^2 r^{-1} \sin(cr) - 2cr^{-2} \cos(cr) + 2r^{-3} \sin(cr) \\ f'''(r) &= -c^3 r^{-1} \cos(cr) + c^2 r^{-2} \sin(cr) + 8cr^{-3} \cos(cr) - 6r^{-4} \sin(cr) \\ f''''(r) &= c^4 r^{-1} \sin(cr) + 2c^3 r^{-2} \cos(cr) - 10c^2 r^{-3} \sin(cr) - 30cs^{-4} \cos(cr) + 24r^{-5} \sin(cr) \end{aligned}$$

and

$$\begin{aligned} g' &= \partial_{x_3} g(x_3) = \frac{x_3}{(x_3^2 + x_2^2 + x_1^2)^{1/2}} \\ g'' &= \partial_{x_3}^2 g(x_3) = \frac{1}{(x_3^2 + x_2^2 + x_1^2)^{1/2}} - \frac{x_3^2}{(x_3^2 + x_2^2 + x_1^2)^{3/2}} \\ g''' &= \partial_{x_3}^3 g(x_3) = -\frac{3}{(x_3^2 + x_2^2 + x_1^2)^{3/2}} + \frac{3x_3^3}{(x_3^2 + x_2^2 + x_1^2)^{5/2}} \\ g'''' &= \partial_{x_3}^4 g(x_3) = -\frac{3}{(x_3^2 + x_2^2 + x_1^2)^{3/2}} + \frac{18x_3^2}{(x_3^2 + x_2^2 + x_1^2)^{5/2}} - \frac{15x_3^4}{(x_3^2 + x_2^2 + x_1^2)^{7/2}} \end{aligned}$$

At this point we recall that by the Faà di Bruno's formula

$$\begin{aligned} \partial_{x_3}^4 (f \circ g)(x) &= f''''(|x|)[g'(x)]^4 + 6f'''(|x|)g''(x)[g'(x)]^2 + 3f''(|x|)[g''(x)]^2 \\ &\quad + 4f''(|x|)g'''(x)g'(x) + f'(|x|)g''''(x) \end{aligned}$$

and the claim (B.1) easily follows by replacing $c = \tau^{1/4}$ and translating back $x \mapsto x - y$.

ACKNOWLEDGEMENTS

J. B. is partially supported by Project 2016 ‘‘Dinamica di equazioni nonlineari dispersive’’ from FONDAZIONE DI SARDEGNA.

REFERENCES

- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, *Observation of Bose-Einstein Condensation in a Dilute Atomic Vapor*, *Science* **269** (1995), no. 5221, 198–201.
- [2] P. Antonelli and C. Sparber, *Existence of solitary waves in dipolar quantum gases*, *Phys. D* **240** (2011), no. 4-5, 426–431.
- [3] W. Bao and Y. Cai, *Mathematical Theory and Numerical methods for Bose-Einstein condensation*, *Kinetic and Related Models AMS* **6** (2013), no. 1, 1–135.
- [4] W. Bao, Y. Cai, and H. Wang, *Efficient numerical method for computing ground states and dynamic of dipolar Bose-Einstein condensates*, *J. Comput. Phys.* **229** (2010), 7874–7892.
- [5] J. Bellazzini and L. Forcella, *Asymptotic dynamic for dipolar quantum gases below the ground state energy threshold*, *J. Funct. Anal.* **277** (2019), no. 6, 1958–1998.
- [6] J. Bellazzini and L. Jeanjean, *On dipolar quantum gases in the Unstable Regime*, *SIAM J. Math. Anal.* **48** (2016), no. 3, 2028–2058.

- [7] C. C. Bradley, C. A. Sackett, J. J. Tolett, and R. J. Hulet, *Evidence of Bose-Einstein Condensation in an Atomic Gas with Attractive Interaction*, Phys. Rev. Lett. **75** (1995), 1687–1690.
- [8] R. Carles, P. A. Markowich, and C. Sparber, *On the Gross-Pitaevskii equation for trapped dipolar quantum gases*, Nonlinearity **21** (2008), no. 11, 2569–2590.
- [9] Y. Cho and T. Ozawa, *Sobolev inequalities with symmetry*, Commun. Contemp. Math. **11** (2009), no. 3, 355–365.
- [10] R. Cipolatti, *On the existence of standing waves for a Davey-Stewartson system*, Comm. Partial Differential Equations **17** (1992), no. 5-6, 967–988.
- [11] K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. Van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Bose-Einstein condensation in a gas of sodium atoms*, Physical Review Letters **75** (1995), no. 22, 3639.
- [12] D. Du, Y. Wu, and K. Zhang, *On blow-up criterion for the nonlinear Schrödinger equation*, Discrete Contin. Dyn. Syst. **36** (2016), no. 7, 3639–3650.
- [13] T. Duyckaerts, J. Holmer, and S. Roudenko, *Scattering for the non-radial 3D cubic nonlinear Schrödinger equation*, Math. Res. Lett. **15** (2008), no. 6, 1233–1250.
- [14] A. Ferrero, F. Gazzola, and H.-C. Grunau, *Decay and eventual local positivity for biharmonic parabolic equations*, Discrete Contin. Dyn. Syst. **21** (2008), no. 4, 1129–1157.
- [15] L. Glangetas and F. Merle, *A Geometrical Approach of Existence of Blow up Solutions in H^1 for Nonlinear Schrödinger Equations*, Laboratoire d'Analyse Numérique, Univ. Pierre and Marie Curie (1995, Rep. No. R95031).
- [16] R. T. Glassey, *On the blowing up of solutions to the Cauchy problem for nonlinear Schrödinger equations*, J. Math. Phys. **18** (1977), no. 9, 1794–1797.
- [17] J. Holmer and S. Roudenko, *On blow-up solutions to the 3D cubic nonlinear Schrödinger equation*, Appl. Math. Res. Express. AMRX (2007), Art. ID abm004, 31. [Issue information previously given as no. 1 (2007)].
- [18] ———, *A sharp condition for scattering of the radial 3D cubic nonlinear Schrödinger equation*, Comm. Math. Phys. **282** (2008), no. 2, 435–467.
- [19] C. E. Kenig and F. Merle, *Global well-posedness, scattering and blow-up for the energy-critical, focusing, nonlinear Schrödinger equation in the radial case*, Invent. Math. **166** (2006), no. 3, 645–675.
- [20] E. A. Kuznetsov, J. J. Rasmussen, K. Rypdal, and S. K. Turitsyn, *Sharper criteria for the wave collapse*, Phys. D **87** (1995), no. 1-4, 273–284. The nonlinear Schrödinger equation (Chernogolovka, 1994).
- [21] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, *The physics of dipolar bosonic quantum gases*, Reports on Progress in Physics **72** (2009), no. 12, 126401.
- [22] J. Lu and Y. Wu, *Sharp threshold for scattering of a generalized davey-Stewartson system in three dimension*, Comm. Pure Appl. Anal. **14** (2015), 1641–1670.
- [23] Y. Martel, *Blow-up for the nonlinear Schrödinger equation in nonisotropic spaces*, Nonlinear Anal. **28** (1997), no. 12, 1903–1908.
- [24] R. Nath, P. Pedri, P. Zoller, and M. Lewenstein, *Soliton-soliton scattering in dipolar Bose-Einstein condensates*, Phys. Rev. A **76** (2007), 013606–013613.
- [25] T. Ogawa and Y. Tsutsumi, *Blow-up of H^1 solution for the nonlinear Schrödinger equation*, J. Differential Equations **92** (1991), no. 2, 317–330.
- [26] L. Pitaevskii and S. Stringari, *Bose-Einstein condensation*, International Series of Monographs on Physics, vol. 116, The Clarendon Press, Oxford University Press, Oxford, 2003.
- [27] P. Raphaël and J. Szeftel, *Standing ring blow up solutions to the N -dimensional quintic nonlinear Schrödinger equation*, Comm. Math. Phys. **290** (2009), no. 3, 973–996.
- [28] L. Santos, G. Shlyapnikov, P. Zoller, and M. Lewenstein, *Bose-Einstein condensation in trapped dipolar gases*, Phys. Rev. Lett. **85** (2000), 1791–1797.

- [29] S. Yi and L. You, *Trapped atomic condensates with anisotropic interactions*, Phys. Rev. A **61** (2000), no. 4, 041604.
- [30] ———, *Trapped condensates of atoms with dipole interactions*, Phys. Rev. A **63** (2001), no. 5, 053607.

JACOPO BELLAZZINI
UNIVERSITÀ DI SASSARI, VIA PIANDANNA 4, 07100 SASSARI, ITALY
E-mail address: `jbellazzini@uniss.it`

LUIGI FORCELLA
ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE, INSTITUTE OF MATHEMATICS, STATION 8, CH-1015 LAUSANNE,
SWITZERLAND.
E-mail address: `luigi.forcella@epfl.ch`