Vinylbenziodoxol(on)es: Synthetic Methods and Applications

Nina Declas,^{a†} Guillaume Pisella^{a†} and Jerome Waser^{*,a}

^a Laboratory of Catalysis and Organic Synthesis, Institute of Chemical Sciences and Engineering, Ecole Polytechnique Fédérale de Lausanne, EPFL SB ISIC LCSO, Av. Forel 2, 1015 Lausanne, Switzerland, jerome.waser@epfl.ch [†]These authors contributed equally.

Dedicated to Prof. Antonio Togni at the occasion of his retirement and 65th birthday.

Cyclic hypervalent iodine reagents are now frequently used in synthetic organic chemistry, either as oxidants or group-transfer reagents. Vinylbenziodoxol(on)es (VBXs) bearing alkene substituents have been less investigated than the corresponding trifluoromethyl or alkynyl reagents. Nevertheless, since 2016 the development of new synthetic methods to access VBXs has awakened the interest of the synthetic community, leading to new transformations highlighting their unique reactivity as electrophilic alkene synthons. In this review, an overview of the synthesis and applications of VBX reagents will be presented. The review is organized according to the two main classes of VBX reagents reported so far –simple alkyl/aryl-substituted VBXs and heteroatom (S, O, N, X)-substituted VBXs– as they differ significantly from the point of view of synthetic access.

Keywords: Hypervalent iodine • Benziodoxolones • Alkenes • Vinylation • Umpolung

1. Introduction

The carbon-carbon double bond is one of the most versatile functional groups in synthetic organic chemistry and it is of paramount importance in applied sciences.^[1–4] Classical strategies such as elimination, carbonyl olefination^[5] or olefin metathesis^[6] enable efficient alkene synthesis. In addition, modern transition metal catalyzed cross coupling reactions and C-H bond functionalizations are routinely used to modify olefins. In this context, vinyl halides (or pseudohalides) have been applied as electrophilic vinyl partners for the creation of sp²-sp² bond. However, vinyl halides can suffer from a lack of reactivity/selectivity in reactions with classical nucleophiles, radical intermediates and/or under metal-free conditions. Hypervalent iodine(III) reagents are environmentally benign and possess enhanced reactivity in comparison to classical iodine (I) compounds.^[7] The exceptional reactivity of iodine(III) derivatives originates from the weak three-center four-electron bond between the iodine atom and the two *trans* ligands associated to it: the *hypervalent bond* (Figure 1).

Figure 1. Hypervalent iodine bond and structure of benziodoxol(on)es.

Hypervalent iodine(III) compounds have become important functional-group transfer reagents in organic synthesis.^[8] Among them, benziodoxol(on)e derivatives are exceptionally stable due to their rigid cyclic structure, and have gained in popularity over the acyclic iodonium salts.^[9] Well-known benziodoxol(on)es such as Togni's reagent^[10] and ethynylbenziodoxol(on)es (EBX)^[11–12] are designed for the transfer of trifluoromethyl and alkynyl groups respectively. They are air and moisture tolerant and convenient to handle. Other benziodoxol(on)e-based reagents for the selective transfer of (hetero)aryl,^[13–16] azide,^[17–24] cyano,^[25–29] and halogens^[30–32] are also frequently used.

Taking advantage of the hypervalent bond, vinylodonium salts have been utilized for the vinylation of various nucleophiles under mild conditions.^[33–39] They have also been used as highly reactive partners in several metal-catalyzed cross-coupling reactions.^[40–47] Nevertheless, their reactivity can be difficult to control and often lead to product mixtures.^[48] They have been studied since several decades and have been the subject of specific reviews and book chapters.^[49–54] In contrast, *vinylbenziodoxol(on)e* reagents (VBX) have emerged only recently as unique iodine(III) reagents for the selective transfer of vinyl groups (Figure 2). This review article will summarize the different syntheses of VBX reagents and then focus on their use as alkenylating reagents in new synthetic methods. Two classes of VBX will be differentiated and treated separately according to their substitution pattern: Compounds incorporating only C or H substituents on the olefin core will be referred as *VBXs* (section 2), while compounds having a heteroatom conjugated to the olefin, will be identified as *hetero-VBXs* and abbreviated *X-VBXs* (with X = N, O, S,...) (section 3).

Figure 2. VBX reagents covered in this review article.

2. VBX

Scheme 1. One-pot synthesis of VBX from 2-iodobenzoic acid (1).

The scope was essentially limited to aryl-substituted derivatives **2a-d**, and milder acidic conditions had to be employed for the more electron-rich **2c**. With a cyclohexyl substituent (**2e**), drastic diminution of the yield was observed. The structure of Phenyl-VBX (Ph-VBX, **2a**) was established by NMR, IR spectroscopy and X-ray crystallography. The VBX products are bench stable and can be kept in the dark at room temperature for prolonged time.

In 2017, an alternative route to access aryl-substituted VBX from pre-formed iodine(III) reagents was proposed by Nachtsheim (Scheme 2).^[56] The procedure follows a common ligand exchange strategy used for the synthesis of vinyl(aryl)iodonium salts.^[57] In this case, readily accessible hydroxybenziodoxolone (**3**) is first activated with TMSOTf, and then treated with vinylboronic acid. Final cyclization in presence of pyridine affords the corresponding VBX reagent **2**.

2.1. Synthesis

The first access to vinylbenziodoxolone reagents was reported in 2016 by the group of Olofsson.^[55] The authors disclosed a practical one-pot synthetic procedure from 2-iodobenzoic acid (**1**) and coined the acronym VBX for the first time (Scheme 1). Oxidation of 2-iodobenzoic acid (**1**) with *m*-CPBA in presence of triflic acid forms the putative iodane intermediate **I**. Addition of vinylboronic acid then leads to iodonium salt **II**, which is then cyclized by mild basic treatment. The obtained VBX reagents **2** can be conveniently isolated by precipitation.

Scheme 2. Synthesis of Ar-VBX **2** from hydroxybenziodoxolone (**3**).

Ph-VBX (**2a**) was efficiently synthesized employing this strategy, when only a moderate yield (39%) was obtained for tolyl-VBX **2b** using a pinacol boronic ester precursor instead of the acid. More recently, **2b** was accessed in good yield starting with the boronic acid precursor and additional examples of aryl-VBX **2c-2i** were disclosed by our group using this protocol.^[58] Furthermore, we developed a new protocol based on Lewis acid activation for the synthesis of alkyl-substituted VBX reagents (Scheme 3). The combination of acetoxybenziodoxolone (**4**) and BF₃·OEt₂ was found optimal to access alkyl-VBX reagents such as **2e**, **2j-i** in better yields. This protocol also gave access to other sensitive reagents, such as thiophene-

substituted VBX **2m**, diene reagent **2n** and enyne reagent **2o**. In addition, the efficient preparation of benziodoxole-based reagents **6a-d**, having a hexafluoroisopropyl ligand backbone, was possible using these new reaction conditions starting from benziodoxole **5**.

Scheme 3. VBX synthesis starting from acetoxybenziodoxol(on)es **4** and **5**.

Since the seminal work of Olofsson in 2016,^[55] several syntheses of vinylbenziodoxol(on)e (VBX) have been therefore developed and allow a straightforward access to these cyclic iodine(III) reagents. A large array of functionalities ((hetero)aryl, alkyl, vinyl, alkynyl) are tolerated on the olefin, and the core structure of the reagents can be also modulated. However, major limitations need to be pointed out: The difficult handling of alkenyl boronic acids as vinyl source may restrain the scope and the practicability of the procedures, but efforts to use more practical precursors (silyl reagents or directly unsubstituted alkenes) were not successful so far. Efforts to synthesize the Z-isomer of VBXs have remained elusive, and these compounds might be unattainable based on previous reports on vinyl(aryl)iodonium salts indicating that both decomposition and isomerization are potential issues.^[59-60]

2.2. Reactivity

Metal-free reactions

In addition to their synthetic protocol, Olofsson and coworkers made a comparative study of the reactivity of Ph-VBX (**2a**) and acyclic vinyl(phenyl)iodonium salt **7** for the C-alkenylation of nitrocyclohexane (**8**) under basic conditions (Scheme 4).^[55] They observed the 1,1-disubstituted alkene **10** as major product in the case of Ph-VBX (**2a**), while the 1,2-

disubstituted isomer **9** was obtained preferentially with iodonium salt **7**. No arylated product **11** was observed with VBX **2a**, which confirmed the complete chemoselectivity for vinyl transfer, whereas traces of side-product **11** were detected with **7**. The reaction was briefly optimized to increase the yield and the regioselectivity in favour of **10**.

Scheme 4. Reactivity comparison between VBX **2a** and vinyliodonium salt **7**.

The same group applied VBX reagents to the electrophilic alkenylation of thiols under mild and transition metal-free conditions (Scheme 5).^[61] After screening of several bases and solvents, *t*BuOK in THF was selected, allowing the formation of products **12** in good yield, excellent *trans* stereoselectivity and with minor formation of disulfide side products. The reaction was suitable for a large array of thiols, including aromatic (**12a-i**) and aliphatic thiols (primary (**12j**), secondary (**12k**) and tertiary (**12l**)). More challenging substrates possessing heterocycles (**12g** and **12i**) or further nucleophilic functions (**12m** and **12n**) were well tolerated, and the double functionalization of di-thiols could be achieved (**12o**). In this work, substitution on the aryl ring of reagent **2** was also investigated. An increase in yield of 20% for product **12j** was observed for a 4,5-dimethyl-substituted reagent, which also allowed easier purification of the product. The scope in respect to VBX reagents was also investigated and allowed the introduction of various aryl-substituted vinyl groups having different electronic properties (**12p-r**). However, vinylation with Cy-VBX (**2e**) gave only modest yield and the scope of alkyl-VBXs was not further explored. Few examples of vinylations with N- and O-VBX were disclosed (**12s** and **12t**). The reaction of N- and S-VBX with sulphur nucleophiles had been already reported previously by our group^[62] and the Miyake group^[63] respectively (See section 3.2).

Scheme 5. Electrophilic vinylation of free thiols with VBX reagents.

Our group reported the alkylation of sulfenate anions, generated via retro-Michael addition from sulfanyl esters **13**, using EBX reagents in presence of *t*BuOK as base.^[64] The transformation was successfully extended to the vinylation using Ar-VBX **2** without re-optimization of the experimental conditions (Scheme 6). Perfect conservation of the stereochemistry was observed in these two particular thio-vinylation cases, allowing access to valuable *E*-vinyl sulfoxides **14a** and **14b**.

Scheme 6. Vinylation of sulfenate anions generated *in situ* from sulfanyl esters **13**.

Scheme 7. Intramolecular photocatalyzed imino-vinylation of olefins with VBX **2**.

Our group has been particularly interested in merging hypervalent iodine chemistry and photoredox-catalyzed transformations. We reported the use of Ph-VBX (**2a**) as radical acceptor in a decarboxylation-fragmentation cascade of cyclic oxime **18** photocatalyzed by organic dye **20** to give nitrile **19** (Scheme 8).^[68] In this example, partial isomerization of the double bond was observed. Isomerization could be minimized by using a milder base.

Scheme 8. Fragmentation of cyclic oxime **18** to give nitrile **19**.

A three-component radical difunctionalization of unactivated alkenes was reported by Studer and Jiang (Scheme 9).^[69] In their methodology, an amidyl radical is generated by decomposition of α -amido-oxy acid precursor under organo-photoredox conditions. The authors reported the amido-alkenylation of 1-hexene (**22**), using **21** and Ph-VBX (**2a**). The expected three-component product **23** was obtained, albeit in low yield.

Scheme 9. Example of 1,2-amido-alkenylation of olefin **22** reported by Studer.

Recently, we disclosed a methodology for the C-terminus functionalization of peptides through a catalytic photoredox decarboxylation.^[70] The work was directed towards alkynylation, but Ph-VBX (**2a**) could be also used for the selective vinylation of dipeptide **24** to give **25** in 82% yield (Scheme 10).

Scheme 10. Olefination at the C-terminus of dipeptide **24** using organo-photoredox catalysis.

Nemoto and coworkers reported in 2020 the C(sp³)-H olefination of THF (**26**) by using Ar-VBX reagents **2** in presence of benzophenone (**28**) as organic triplet-sensitizer at 400 nm (Scheme 11).^[71] Good yields were

obtained. However, the irradiating conditions led to isomerization of the double bond.

Scheme 11. Benzophenone (**28**) promoted vinylation of THF (**26**).

Metal-catalyzed transformations with VBX

Transition-metal-catalyzed activation of inert C(sp²)-H bonds is a particularly active research area in organic synthesis.^[72] An iridium-catalyzed direct C-H alkenylation of 2-vinylaniline derivatives with VBX reagents **2** was reported by the Nachtsheim group (Scheme 12).^[56] The key deprotonation-metalation event was directed by the NH₂ group and VBXs showed superior reactivity compare to vinylodonium salts. This method gave direct access to useful functionalized 1,3-dienes **28a-h** in excellent yields and with high to perfect (Z,E) stereoselectivity.

Scheme 12. Ir(III)-catalyzed C-H vinylation of 2-vinylanilines.

Finally, our group has studied the use of VBX reagents in copper catalysis. In our work for the difunctionalization of thiirane and thiethanes with hypervalent iodines reagents, we reported two examples of ring-opening reaction of cyclohexene sulfide (**29**) with Ph-VBX (**2a**) and Cy-VBX (**2e**) to access vinyl-thioethers **30a** and **30b** (Scheme 13).^[73] The released benzoate **31** acts as a nucleophile, resulting in perfect atom economy, which is rare in hypervalent iodine chemistry.^[74-75] The copper (I) catalyst is believed to activate the iodine(III) reagent and accelerate the formation of a reactive episulfonium intermediate **I**.

Scheme 13. Copper-catalyzed oxyvinylation of thiirane **29**.

In 2020, we disclosed the insertion of VBX reagents into diazo compounds using copper catalysis (Scheme 14).^[58] Based on our previous experience on oxyalkynylation,^[76-78] we identified VBX reagents **2** as privileged partners to realize the 1,1-oxyvinylation of transient copper-carbene

Scheme 14. Copper-catalyzed 1,1-oxyvinylation of diazo compounds using VBX reagents **2** and **6**.

3. Hetero-VBX

3.1. Synthesis of hetero-VBX

Among VBX reagents, hetero-substituted vinylbenziodoxolones (X-VBX, **2**) have been broadly studied since 2013. The first reported vinyl-substituted cyclic iodinanones **35** were discovered by the group of Kitamura as early as 1996 (Scheme 15).^[82] Iodinanones **35** were obtained from a Michael addition reaction of sodium azide on alkynyl-benzoic-iodonium triflates **34**. The unexpected *syn*-addition, leading to the *trans* olefin, contrasted with previous reactions of alkynyl-iodonium salts with azide nucleophiles that usually led to the *cis* isomer. Different N-VBX **35** bearing alkyl substituents from butyl to decyl could be obtained in moderate yields.

intermediates. This transformation cannot be performed using existing methodologies with vinyl halides.^[79-81] The reaction was proposed to start with the nucleophilic attack of the carboxylate of VBX **2** onto the highly electrophilic copper-carbene **II** generated by reaction of the catalyst **I** with the diazo compound. Vinyl transfer from iodonium intermediate **III** would then give oxyvinylation product **32**. No isomerization of the transferred olefin was observed. Importantly, the reaction tolerated well the presence of various substituents on the alkene (products **32a-c** and **32f**), which motivated us to develop new functionalized VBX reagents (See section 2.1). Variation of the diazo compound was also possible (products **32d** and **32e**). In addition, the utilization of non-nucleophilic benziodoxole-based VBX **6** enabled the extension of the reaction to a three-component variant with external alcohol nucleophiles instead of the iodobenzoate (products **33a** and **33b**).

Scheme 15. First reported vinyl-substituted cyclic iodinanones **35**.

After this pioneering discovery, the study of X-VBX reagents remained latent for many years. In 2013, our group published a chemoselective thioalkynylation with TIPS-EBX.^[83] The reaction proceeded at room temperature within five minutes, making the study of its mechanism particularly challenging. With Me-EBX (**36a**) thiovinyl benziodoxolone **39a** was observed as a side product in the synthesis of thioalkyne **38a** from benzyl thiol (**37**) (Scheme 16).^[84] Interestingly, the use of a catalytic amount of base allowed to isolate **39a** in 20% yield as a single *Z* isomer. Its formation was first rationalized by protonation of the vinyl iodane **I**

generated by conjugate addition as proposed by Ochiai (Scheme 16A).^[85] In contrast, alkynylation product **38** would be obtained via α -elimination to give carbene **II** followed by 1,2-migration. Computational studies by our group later uncovered two further reaction pathways: Starting from iodine-sulfur interaction complex **III**; both a concerted α -addition/elimination transition state **IV** and a β -addition transition state **V**

with a sulfur-iodine interaction were located (Scheme 16B).^[86] The latter pathway is favored for alkyl substituents and leads to the same vinyl iodane **I** as proposed by Ochiai, rationalizing the formation of **39**. The synthesis of S-VBX reagents was however not further optimized at this stage.

Scheme 16. Alkynylation of thiols with EBX reagents, isolation of S-VBX intermediate **39a** and proposed mechanisms.

In 2019, our group reported a cysteine bioconjugation methodology for the introduction of hypervalent iodine compounds onto biomolecules (Scheme 17).^[87] Cysteine-containing peptides and proteins were engaged in a selective addition onto the alkynyl triple bond of EBX **36b**, resulting in stable VBX conjugates **39b**. The general cysteine labeling protocol demonstrate high efficiency, chemoselectivity, and functional-group tolerance under native conditions. The hypervalent iodine and the azide group could be then selectively modified (see section 3.2).

Scheme 17. Reaction of EBX **36b** with cysteine-containing biomolecules.

Using modified conditions, thiols contained in small organic molecules could also be used for the synthesis of S-VBX reagents (Scheme 18). Derivatives **39a** and **39c-d** were obtained in moderate yield with complete Z-stereoselectivity.^[87]

Scheme 18. Stereoselective synthesis of S-VBX reagents **39a** and **39c-d**.

The formation of O-VBXs **41** from EBX reagents **40** was firstly studied by Yoshikai and coworkers (Scheme 19). In 2016, they published the formation of *E*-alkenylbenziodoxoles **41** from the palladium-catalyzed

Scheme 19. Pd-catalyzed conversion of EBX **40** to functionalized O-VBX reagents **41** and **42** through 1,2-I(III) shift and 1,1-hydrocarboxylation or 1,1-carboxyalkynylation.

The addition of phenols on EBXs **36** was later reported by Miyake and coworkers, who developed a regio- and stereoselective synthesis of *Z*-2-iodovinyl phenyl ethers **43** under blue LED irradiation in presence of Cs_2CO_3 as a base (Scheme 20).^[90] Several vinyl iodides such as **43a-43d** were obtained in good to excellent yields. The reaction is efficient with a broad range of phenols. In addition, different phenyl-substituted EBXs **36** can be used in the transformation, and one example is described with an EBX reagent **36** bearing an alkyl group. Interestingly, control experiments in absence of light led to the observation and isolation of O-VBX **44a**. Based on a series of experiments and computations, they proposed that the transformation proceeds through an electron donor-acceptor (EDA) complex I. In I, a photoinduced electron transfer is supposed to occur from

addition of carboxylic acids to EBXs **40**.^[88] Mechanistic experiments and computations suggested that the Pd(II) carboxylate-ligated alkynylbenziodoxole **I** undergoes a 1,2-I(III) shift to form **II**, followed by a migratory insertion of the vinylidene-Pd species (1,1-hydrocarboxylation) to give **III**. Finally protonation delivers the O-VBX reagent **41** with a *trans* geometry. The use of the COD ligand was critical to prevent decomposition. By changing the ligand to octahydrophenazine and increasing the equivalents of EBX **40**, alk-1-en-3-ynylbenziodoxoles **42** were obtained via interception of **III** with **40**, leading to a 1,1-carboxyalkynylation.^[89] Both methods tolerated a variety of alkyl- alkenyl- alkynyl- and aryl(alkynyl)benziodoxoles and a series of alkyl- and arylcarboxylic acids, leading to hypervalent iodine reagents such as **41a-c** and **42a-c** in excellent yields. The reagents could be efficiently used as precursors in cross-coupling reactions (see section 3.2).

the phenoxide donor to the O-VBX acceptor. DFT calculations support that this one-electron reduction of the VBX leads to spontaneous phenyl-iodine bond cleavage to provide the *Z*-2-iodovinyl phenyl ethers **43**. Using Cs_2CO_3 , light was shown to be essential for the transformation. However, when the strong organic base 1,5,7-triazabicyclo[4.4.0]dec-5-ene (TBD) was used, no light was needed to promote the reaction. In this case, a thermal activation of the electron transfer was proposed.^[91]

Scheme 20. Light-driven synthesis of Z-2-iodovinyl phenyl ethers **43** and isolation of O-VBX **44a**.

This work was an important breakthrough for the metal free addition of nucleophiles on EBX, however only two examples of VBX reagents were isolated. At the same time, our group was working on the synthesis of new O- and N-VBX reagents with focus on a protocol allowing to isolate them in good yields.^[62] Based on the insights gained from the isolation of S-VBX **39a** in our previous work on the alkynylation of thiols,^[84] optimized reaction conditions were developed (Scheme 21). Z-Enol ether and enamide-substituted benziodoxolones reagents **44**, and **45** were obtained by stereoselective addition of phenols and sulfonamides to EBXs **36** using a catalytic amount of Cs₂CO₃. The mild conditions developed tolerated numerous functional groups and the methodology could be used for modification of natural products and drugs. O- and N-VBXs such as **44b-c**, **45a-c** were obtained in good to excellent yields.

Scheme 21. Stereoselective synthesis of O- and N-VBX reagents **44b-c** and **45a-c**.

Nucleophilic addition of sulfonamides onto EBXs **36** had been previously reported by the group of Cossy, but this methodology, which involved a stoichiometric amount of base, led to the formation of the corresponding ynamides **46**.^[92] To better understand the different reaction outcome when using a catalytic amount of base, computational studies were performed and an α and a β -addition transition state could be located (**I** and **II** in Scheme 22). The β -addition was shown to be favoured by about 10 kcal mol⁻¹ to give **III**. Protonation of **III** was favoured by 15 kcal mol⁻¹ compared to carbon-iodine bond breaking leading to **46**, giving **45** as the only product. However, in a presence of an excess of base, larger amounts of **III** are present in the reaction mixture, allowing for irreversible carbon-iodine cleavage.

Scheme 22. Proposed mechanism for the selective formation of N-VBX reagent **45** over alkylation product **46**.

During this work, one example of unsubstituted N-VBX **45** could be obtained from trimethylsilyl-EBX (**36c**), with *in situ* desilylation (Scheme 23). In 2019, Itoh, Tada and coworkers reported the synthesis of *cis*- β -amidevinyl benziodoxolones from an ethynyl-1,2-benziodoxol-3(1*H*)-one-chloroform complex (**36d**).^[93] The reaction proceeded under mild conditions, in presence of a catalytic amount of base and sulfonamides derived from various amines, including amino acids, could be used as nucleophiles for the transformation. Hypervalent iodine reagents **45d-f** could be accessed in excellent yields. Furthermore, a synthetic approach for the synthesis of selectively deuterated N-VBX **45** was also developed.

Benziodoxolones are the starting material of choice for the stereoselective synthesis of hetero-VBX, by addition of phenol, sulphonamide and thiol nucleophiles. However, with the previously described methodologies, the addition of aliphatic alcohols could not be achieved. Yoshikai and coworkers answered this challenge with a regio- and stereoselective synthesis of highly substituted β - λ^3 -iodanyl vinyl ethers **48** achieved through *trans*-1,2-difunctionalization of alkynes with benziodoxole triflate (BXT, **47**) and alcohols (Scheme 24).^[94] No base was required for the reaction to proceed and the mild conditions developed tolerated various internal and terminal alkynes, as well as a large range of alcohols, furnishing derivatives such as **48a-e** in high yields. In contrast to previously established methods, this approach did not require the preparation of a different EBX **40** for each O-VBX **48**. Supported by DFT calculations, the reaction was proposed to proceed through a stereoselective addition of alcohol via transition state **I**. Upon dissociation of triflate from BXT (**47**), a BX cation activates the alkyne, which triggers the nucleophilic addition of alcohol onto the triple bond with a Markovnikov-type selectivity. In the case of unsymmetrical dialkyl alkynes, the attack of alcohol was shown to occur on the sterically less hindered alkyne carbon. Such mechanism gives a *trans*- β -addition, contrary to the usual *cis*- β -addition observed for addition on EBX reagents.

A combination of fluorobenziodoxole (FBX, **49**) and $\text{BF}_3 \cdot \text{OEt}_2$ in ethereal solvent was proposed as an alternative to generate the BX cation (Scheme 25).^[95] The reaction proceeded with the same regio- and stereoselectivity as previously described, and after nucleophilic attack of cyclopentyl methyl ether (CPME), the oxonium intermediate undergo β -elimination to afford product **48**, HBF_4 and cyclopentene. Compounds such as **48f-i** could be synthesized in comparable to better yields than with the BXT (**47**)/alcohol system. A broad range of internal and terminal alkynes is again tolerated, but, compared to the previous methodology, the scope of alkoxy groups is limited to methoxy derived from the solvent CPME.

Scheme 25. Stereoselective synthesis of O-VBX **48** from FBX (**49**) and CPME.

Li, Wen and coworkers reported a substituent-controlled synthesis of oxadiazine derivatives from EBX **36** and amidoximes **50** (Scheme 26).^[96] Under basic conditions, the deprotonated amidoximes **50** attacks the β -carbon of EBX **36** to form a *Z*-vinylbenziodoxolone intermediate **I** that could be isolated at room temperature in the case of **51**. At higher temperature, VBX intermediate **I** can undergo either intramolecular cyclization to give **52** or a more complex addition/condensation cascade with a second equivalent of **36** and **50** resulting in heterocycle **53** depending on the substrate.

Scheme 26. Synthesis of oxadiazine derivatives **52** and **53** from EBXs **36** and amidoximes **50**.

Halovinylbenziodoxoles **54** and **55** were also synthesized by the group of Yoshikai who reported the stereoselective hydrochlorination and iodochlorination of EBX **40** (Scheme 27).^[97] The reactions were achieved

under mild, open-air conditions. Highly substituted 2-chlorinated VBX **54a-c** were obtained by *anti*-hydrochlorination using pyridine hydrochloride as an HCl source. The procedure tolerates a variety of EBXs **40** derived from (hetero)aryl- and alkyl-acetylenes and the hydrochlorination could even be extended to ethynylbenziodoxolone and ethynyl(phenyl)iodonium tosylate. A narrower scope was observed for the iodochlorination and alkyl-EBXs **40** were not tolerated in the transformation. The reaction, which involved iodine monochloride, surprisingly afford the *syn*-iodochlorination products **55a-b**, contrasting with the *anti*-selectivity usually observed. Whereas the hydrochlorination is supposed to proceed through β -addition of a chloride ion to EBX **40**, followed by protonation of the resulting vinyl anion **I**, the iodochlorination follows a different mechanism. Based on DFT calculations, the authors proposed an electrophilic activation of EBX **40** by an iodine cation, followed by structural reorganization with formation of the new C-I bond and a new iodonium bridge with the BX cation (**II**). For steric reasons, the chloride anion would then approach from the opposite side of the BX group, leading to the *syn*-iodochlorination product **55**.

Scheme 27. Stereoselective hydrochlorination and iodochlorination of EBX **40**.

Metal-VBX complexes have also been proposed as intermediates of reactions in palladium-catalyzed condensation of *N*-aryl imines^[98] and carboxyalkynylation of olefins^[99] with hypervalent iodine reagent. However, these intermediates were not isolated and could not be characterized.

3.2 Reactivity of hetero-VBX

Hypervalent iodine reduction

Hypervalent iodine derivatives can be easily reduced to their monovalent form to deliver vinyl iodides. In the work of Miyake, the hypervalent iodine

species **2** is reduced *in situ* by a photoinduced electron transfer to give **43** (Scheme 20 and 28).^[90] Photoredox conditions were also used by our group to cleave the hypervalent bond of N-VBX **45**, delivering the monovalent compound **56**.^[62] Copper-catalyzed reduction is another effective method used by the groups of Yoshikai and Itoh and Tada, to reduce the benziodoxol(on)es moiety without affecting the olefin stereochemistry.^[93-94] Using a copper(I) reductant, iodine **57** and **58** were obtained in good yields at 50 °C.

Scheme 28. Reduction of the hypervalent iodine bond in VBX.

Metal catalyzed C-C bond formation

Transition-metal-catalyzed reactions often require high temperature for substrate activation. Hypervalent iodine reagents, with their enhanced reactivity, are ideal coupling partners for metal-catalyzed transformations at lower temperature. Therefore, X-VBXs have been broadly used in reactions such as the palladium-catalyzed Stille, Sonogashira and Suzuki-Miyaura cross-couplings. The reactions usually proceed at room temperature, with complete retention of the stereoselectivity.

Yoshikai and coworkers reported that O-VBX reagent **41c** could be used in a palladium-catalyzed Stille cross-coupling with aryl stannanes at room temperature, leading to stereodefined tri-substituted alkene **59** (Scheme 29).^[88] The C-O bond on the alkene could be further functionalized by iron-catalyzed cross-coupling with butylmagnesium bromide to give **60**. Access to functionalized tetra-substituted enynes was possible using the products obtained from 1,1-carboxyalkynylation (Scheme 19).^[89]

Scheme 29. Stille followed by iron-catalyzed cross-coupling of O-VBX reagent **41c**.

Similarly, our group used palladium-catalysis for the modification of O- and N-VBX **44** and **45**.^[62] Coupling of aryl, vinyl and alkylstannyl reagents were performed at room temperature with complete stereoselectivity to give the *Z* products **61** (Scheme 30). Compounds **61a-c** were synthesized in 52-88% yields. A Sonogashira cross-coupling of **45a** with trimethylsilyl acetylene could be also performed, with however a slight isomerization of the double bond, leading to **62** in 44% yield.

Scheme 30. Stille and Sonogashira cross-coupling of N-VBX reagent **45**.

A direct comparison of the reactivity of monovalent versus hypervalent iodine toward cross-coupling was performed by our group (Scheme 31).^[62] N-VBX **45b** was synthesized from EBX **36** using the previously described conditions, and its monovalent analogue **56** could be obtained by reduction of **45b** using photoredox conditions.^[90] Both reagents were engaged in a Stille cross-coupling with vinyl stannane. At room temperature, hypervalent iodine **45b** could be transformed into the vinyl product **61d** in 80% yield, whereas no conversion occurred with simple iodine **56**. When heating up to 75 °C, 10% of the desired product could be observed with **56**. This experiment highlighted the efficiency and synthetic utility of hypervalent iodine reagents in cross-coupling reactions.

Scheme 31. Stille cross-coupling of hypervalent iodine **45b** versus monovalent iodine **56**.

Room temperature metal-catalyzed transformations were applicable to the highly substituted β - λ^3 -iodanyl vinyl ethers **48** developed by Yoshikai and coworkers, for Sonogashira and Suzuki reactions, whereas higher temperature was required for Stille coupling of the reagents.^[94] Their reagents could take part also in Rosenmund-von Braun cyanation, as well as palladium-catalyzed hydrodehalogenation of the BX moiety to give **63** (Scheme 32). Product **63**, generated *in situ*, could undergo cyclopropanation, inverse electron-demand Diels-Alder and ytterbium triflate catalyzed Mannich reaction, leading to products **64–66**.

Scheme 32. O-VBX palladium-catalyzed hydrodehalogenation of **48j** and further functionalization of enol ether **63**.

Halovinylbenziodoxoles **54** displayed similar reactivity regarding metal-catalyzed reactions, and chloro-VBX **54a** was used in Stille and Sonogashira couplings with alkynes and stannanes at room temperature (Scheme 33). Cross-coupling of the hypervalent bond of iodochlorinated-VBXs **55** was not successful.^[97]

Scheme 33. Stille and Sonogashira cross-coupling of Cl-VBX reagent **54a**.

Our group developed a Suzuki-Miyaura cross coupling reaction of S-VBX reagent **36e** installed on glutathione with boronic acids to give alkenes **69a–d** (Scheme 34).^[87] The reaction is selective for the vinyl hypervalent iodine bond and the biocompatible conditions could be applied to larger peptides and even proteins. In addition, the azide could be selectively functionalized by cycloaddition. The hypervalent iodine and the azide are therefore orthogonal in reactivity to each other and to existing natural functional groups in peptides and proteins, allowing applications for peptide and protein functionalization. For example, a triplet-state quencher and a fluorophore were introduced selectively, leading to extended fluorescence lifetime by suppressing photobleaching.

Scheme 34. Aqueous Suzuki-Miyaura cross coupling reaction of S-VBX **39e**.

Metal-free C-X bond formation

Iodonium salts are known to react as vinyl cation equivalents in presence of nucleophiles.^[100] Hetero-vinylbenziodoxol(on)es have been shown to display similar reactivity. Our group was able to transform N-VBX reagent **45g** into thioenamide **71**, using thiophenol (**70**) in presence of an excess of base (Scheme 35).^[62] The mechanism of the reaction was not studied but the slight isomerization observed suggest an addition-elimination pathway.

Scheme 35. Addition of thiophenol (**70**) onto N-VBX **45g**.

Thiol nucleophiles were also used by the group of Miyake for the synthesis of 1,2-dithio-1-alkenes **72** (Scheme 36A).^[63] In presence of an excess of nucleophile and base, the addition of two thiol equivalents to the C-C triple bond of EBX **36** was observed, leading to Z-1,2-dithio-1-alkene **72** with loss of the hypervalent bond. A number of aliphatic and aromatic thiols, as well as thioglycosides are compatible with the reaction conditions. Compounds **72a-c** were obtained in excellent yield and high stereoselectivity. Experimental and computational mechanistic studies revealed that the *cis* regioselectivity observed in the Z-1,2-bisthiolated alkene product is explained by the combination of two steps: *cis*-selective nucleophilic thiol addition to the β -position of EBX **36**, leading to an S-VBX intermediate, followed by a *cis*-specific radical addition of a second equivalent of thiol (**1**). Using a catalytic amount of base, S-VBX intermediate **39f** could be isolated and then converted to the dithiolated product **72d** in presence of an excess of Cs_2CO_3 (Scheme 36B). Phenyl-VBX (**2a**) could also be transformed to vinyl sulfides under the reported conditions, which is consistent with the work later published by the group of Olofsson on electrophilic vinylation of thiols with VBX (See section 2.2).^[61]

Scheme 36. Reaction of EBX **36** and S-VBX **39f** with thiol nucleophiles.

Scheme 36. Reaction of EBX **36** and S-VBX **39f** with thiol nucleophiles.

Nucleophilic addition onto O-VBX reagents **44** was studied by our group. Surprisingly, the O-VBXs **44** displayed an oxy-allyl cation-like behaviour, rather than the expected vinyl cation reactivity (Scheme 37).^[101] This new mode of reactivity was observed in presence of an excess of cesium carbonate and is most efficient with phenols as nucleophiles, affording aryl enol ethers products **74** with *E* stereoselectivity. Few examples were also presented with carbon and nitrogen nucleophiles. The obtained ether derivatives **74** could be transformed into α -difunctionalized ketones **75** under oxidative conditions in presence of acid. In absence of external nucleophiles, the 2-iodobenzoate group of the O-VBX reagent is transferred, leading to aryl enol esters substrates **76**. For example, both the ether products **74a-c** and the ester product **76a** could be obtained in good to excellent yields. The reaction was proposed to proceed through isomerization of VBX **44** to an allyl iodonane intermediate, followed by either a $\text{S}_{\text{N}}1$ pathway, with α -elimination of the aryl iodide to deliver an oxy-allyl cation species, or a $\text{S}_{\text{N}}2$ mechanism, with direct reaction of the nucleophile with the allylic hypervalent iodine intermediate.

Scheme 37. O-VBX reagents **44** as oxy-allyl cation synthetic equivalents.

Summary and Outlook

The quest for new vinyl cation surrogates for enabling effective olefination methods has drawn important research efforts in organic chemistry. Benziodoxolone-based transfer reagents have recently emerged as privileged reagents to reach this goal due to their high reactivity combined with sufficient stability. Both VBX and hetero-VBX reagents have attracted significant attention since 2016.

Convergent methods for the preparation of VBXs, essentially from boron precursors, have been explored and accommodate well various kind of substitutions ((hetero)aryl, alkyl, alkenyl, alkynyl) on the double bond. Although important progress has been achieved, the development of synthetic routes from more readily available starting materials remain desirable to popularize the use of VBXs within the organic chemist community. Since their introduction by Olofsson in 2016, VBXs as electrophilic alkene synthons have enabled unprecedented transformations in different domains such as transition-metal catalysis, radical chemistry or classical ionic chemistry. Different outcome when compared to vinylhalides and acyclic vinyl iodonium salts have been often observed.

Hetero-VBX possess an extra hetero-atom (N, O, S or halogen) conjugated to the alkene, which influences considerably the electronics of the later. X-VBX reagents have been obtained mostly through the addition of soft nucleophiles onto well-established EBX reagents, which need nevertheless to be first synthesized. In this respect, the new difunctionalization of alkynes from simple iodane precursors and alcohols developed by Yoshikai

and coworkers constitutes an important progress. The formation of X-VBX proceeds usually with high stereoselectivity and gives access to highly functionalized reagents with defined geometry, a challenging task for enols and enamines. The high reactivity of the hypervalent bond serves as basis for cross-coupling reactions under mild conditions to further diversify the products.

Vinylbenziodoxolones have been surprisingly nonexistent in the hypervalent iodine transfer reagent tool box until 2016 and have emerged only very recently. This may be due to the well-developed chemistry of electrophilic alkene synthons, especially halides. In contrast, electrophilic alkyne synthons were less developed, which led to a faster rise of EBX reagents. Nevertheless, the results obtained in the last five years with VBX reagents demonstrated their unique reactivity, especially for the Umpolung of nucleophilic olefins such as enol ethers and enamides. It is clear that their synthetic potential has just started to be investigated. We hope that our overview of the current state of the art will stimulate and inspire the synthetic community to further develop the synthetic access and new applications of VBX reagents.

Acknowledgements

We thank the Swiss National Science Foundation (SNSF, grant no. 200020-182798) and EPFL for financial support.

References

- [1] *The Chemistry of Alkenes*, (Ed.: S. Patai), Wiley, New York, **1964**, vol. 1.
- [2] *The Chemistry of Alkenes*, (Ed.: J. Zabicky), Wiley, New York, **1970**, vol. 2.
- [3] *Supplement A: The Chemistry of Double Bonded Functional Groups*, (Ed.: S. Patai), Wiley, New York, **1989**, vol. 2, Parts 1 and 2.
- [4] *The Chemistry of Double-Bonded Functional Groups*, (Ed.: S. Patai), Wiley, New York, **1997**, vol. 2, Parts 1 and 2.
- [5] *Modern Carbonyl Olefination*, (Ed.: T. Takeda), Wiley, Weinheim, **2003**.
- [6] *Handbook of Metathesis*, (Ed.: R. H. Grubbs), Wiley, Weinheim, **2003**.
- [7] V. V. Zhdankin, *Hypervalent Iodine Chemistry: Preparation, Structure, and Synthetic Applications of Polyvalent Iodine Compounds*, Wiley, New York, **2014**.
- [8] A. Yoshimura, V. V. Zhdankin, 'Advances in Synthetic Applications of Hypervalent Iodine Compounds' *Chem. Rev.* **2016**, 116, 3328-3435.
- [9] Y. Li, D. P. Hari, M. V. Vita, J. Waser, 'Cyclic Hypervalent Iodine Reagents for Atom-Transfer Reactions: Beyond Trifluoromethylation' *Angew. Chem., Int. Ed.* **2016**, 55, 4436-4454.
- [10] J. Charpentier, N. Fröh, A. Togni, 'Electrophilic Trifluoromethylation by Use of Hypervalent Iodine Reagents' *Chem. Rev.* **2015**, 115, 650-682.
- [11] J. Kaschel, D. B. Werz, 'Ethyne Benziodoxolone (EBX): Installing Alkynes the Reversed Way' *Angew. Chem., Int. Ed.* **2015**, 54, 8876-8878.
- [12] Waser J. Alkynylation with Hypervalent Iodine Reagents. In: Wirth T. (eds) *Hypervalent Iodine Chemistry*. Top. Curr. Chem. **2015**, vol 373. p. 187-222, Springer, Cham.
- [13] P. Caramenti, S. Nicolai, J. Waser, 'Indole- and Pyrrole-BX: Bench-Stable Hypervalent Iodine Reagents for Heterocycle Umpolung' *Chem. Eur. J.* **2017**, 23, 14702-14706.

- [14] E. Grenet, J. Waser, 'Iridium- and Rhodium-Catalyzed Directed C–H Heteroarylation of Benzaldehydes with Benziodoxolone Hypervalent Iodine Reagents' *Org. Lett.* **2018**, *20*, 1473-1476.
- [15] B. Wu, J. Wu, N. Yoshikai, 'Benziodoxole Triflate as a Versatile Reagent for Iodo(III)cyclization of Alkynes' *Chem. Asian J.* **2017**, *12*, 3123-3127.
- [16] A. A. Almasalma, E. Mejia, '1-Phenyl-1,2-benziodoxol-3-(1H)-one as Synthon for Phthalide Synthesis through Pd-Free, Base-Free, Sonogashira-Type Coupling Cyclization Reaction' *Eur. J. Org. Chem.* **2018**, 188-195.
- [17] V. V. Zhdankin, A. P. Krasutsky, C. J. Kuehl, A. J. Simonsen, J. K. Woodward, B. Mismash, J. T. Bolz, 'Preparation, X-ray Crystal Structure, and Chemistry of Stable Azidiodinanes Derivatives of Benziodoxole' *J. Am. Chem. Soc.* **1996**, *118*, 5192-5197.
- [18] Q.-H. Deng, T. Bleith, H. Wadepohl, L. H. Gade, 'Enantioselective Iron-Catalyzed Azidation of β -Keto Esters and Oxindoles' *J. Am. Chem. Soc.* **2013**, *135*, 5356-5359.
- [19] M. V. Vita, J. Waser, 'Azidation of β -Keto Esters and Silyl Enol Ethers with a Benziodoxole Reagent' *Org. Lett.* **2013**, *15*, 3246-3249.
- [20] A. Sharma, J. F. Hartwig, 'Metal-Catalysed Azidation of Tertiary C–H Bonds Suitable for Late-Stage Functionalization' *Nature* **2015**, *517*, 600-604.
- [21] S. Alazet, F. Le Vaillant, S. Nicolai, T. Courant, J. Waser, 'Divergent Access to (1,1) and (1,2)-Azidolactones from Alkenes using Hypervalent Iodine Reagents' *Chem. Eur. J.* **2017**, *23*, 9501-9504.
- [22] S. Bertho, R. Rey-Rodriguez, C. Colas, P. Retailleau, I. Gillaizeau, 'Regio- and Stereoselective Iron-Catalyzed Oxyazidation of Enamides Using a Hypervalent Iodine Reagent' *Chem. Eur. J.* **2017**, *23*, 17674-17677.
- [23] Z.-L. Li, X.-H. Li, N. Wang, N.-Y. Yang, X.-Y. Liu, 'Radical-Mediated 1,2-Formyl/Carbonyl Functionalization of Alkenes and Application to the Construction of Medium-Sized Rings' *Angew. Chem., Int. Ed.* **2016**, *55*, 15100-15104.
- [24] S. Alazet, J. Preindl, R. Simonet-Davin, S. Nicolai, A. Nanchen, T. Meyer, J. Waser, 'Cyclic Hypervalent Iodine Reagents for Azidation: Safer Reagents and Photoredox-Catalyzed Ring Expansion' *J. Org. Chem.* **2018**, *83*, 12334-12356.
- [25] V. V. Zhdankin, M. C. Scheuller, P. J. Stang, 'A General Approach to Aryl(Cyano)Iodonium Triflates - Versatile Iodonium Transfer Reagents' *Tetrahedron Lett.* **1993**, *34*, 6853-6856.
- [26] R. Frei, T. Courant, M. D. Wodrich, J. Waser, 'General and Practical Formation of Thiocyanates from Thiols' *Chem. Eur. J.* **2015**, *21*, 2662-2668.
- [27] Y.-S. Feng, Y.-J. Shu, P. Cao, T. Xu, H.-J. Xu, 'Copper(I)-Catalyzed Ring-Opening Cyanation of Cyclopropanols' *Org. Biomol. Chem.* **2017**, *15*, 3590-3593.
- [28] F. Le Vaillant, M. D. Wodrich, J. Waser, 'Room Temperature Decarboxylative Cyanation of Carboxylic Acids Using Photoredox Catalysis and Cyanobenziodoxolones: A Divergent Mechanism Compared to Alkynylation' *Chem. Sci.* **2017**, *8*, 1790-1800.
- [29] B. Ma, X. Lin, L. Lin, X. Feng, X. Liu, 'Chiral *N,N'*-Dioxide Organocatalyzed Asymmetric Electrophilic α -Cyanation of β -Keto Esters and β -Keto Amides' *J. Org. Chem.* **2017**, *82*, 701-708.
- [30] M. Wang, Y. Zhang, T. Wang, C. Wang, D. Xue, J. Xiao, 'Story of an Age-Old Reagent: An Electrophilic Chlorination of Arenes and Heterocycles by 1-Chloro-1,2-benziodoxol-3-one' *Org. Lett.* **2016**, *18*, 1976-1979.
- [31] H. Egami, T. Yoneda, M. Uku, T. Ide, Y. Kawato, Y. Hamashima, 'Difunctionalization of Alkenes Using 1-Chloro-1,2-benziodoxol-3-(1H)-one' *J. Org. Chem.* **2016**, *81*, 4020-4030.
- [32] N. O. Ilchenko, M. Hedberg, K. J. Szabó, 'Fluorinative Ring-Opening of Cyclopropanes by Hypervalent Iodine Reagents. An Efficient Method for 1,3-Oxyfluorination and 1,3-Difluorination' *Chem. Sci.* **2017**, *8*, 1056-1061.
- [33] M. Ochiai, K. Sumi, Y. Takaoka, M. Kunishima, Y. Nagao, M. Shiro, E. Fujita, 'Reactions of Vinylsilanes with Lewis Acid-Activated Iodosylbenzene: Stereospecific Syntheses of Vinylodonium Tetrafluoroborates and their Reactions as Highly Activated Vinyl Halides' *Tetrahedron* **1988**, *44*, 4095-4112.
- [34] F. M. Beringer, S. A. Galton, 'Acetylenic and Ethylenic Iodonium Salts and Their Reactions with a Carbanion' *J. Org. Chem.* **1965**, *30*, 1930-1934.
- [35] J. Guo, L. Lin, Y. Liu, X. Li, X. Liu, X. Feng, 'Nickel(II)-Catalyzed Enantioselective α -Vinylolation of β -Keto Amides/Esters with Hypervalent Iodine Salts' *Org. Lett.* **2016**, *18*, 5540-5543.
- [36] I. Papoutsis, S. Spyroudis, A. Varvoglis, 'Reactivity of a New Alkenyl Phenylodonium Tosylate Derived from Methyl 3-Aminocrotonate' *Tetrahedron* **1998**, *54*, 1005-1012.
- [37] T. Kitamura, P. J. Stang, 'Generation, Trapping and Fate of Alkylidenecarbene-Iodonium Ylides from the Addition of NaN_3 to Alkynylphenylodonium Tosylates' *Tetrahedron Lett.* **1988**, *29*, 1887-1889.
- [38] M. Ochiai, S. Yamamoto, T. Suefuji, D.-W. Chen, 'Stereoselective Synthesis of (Z)-Enethiols and Their Derivatives: Vinylic SN_2 Reaction of (E)-Alkenyl(phenyl)- λ^3 -iodanes with Thioamides' *Org. Lett.* **2001**, *3*, 2753-2756.
- [39] S. Thielges, P. Bissere, J. Eustache, 'Copper-Mediated Cross-Coupling of *H*-Phosphonates with Vinylodonium Salts: A Novel Very Mild Synthesis of 2-Arylvinylphosphonates' *Org. Lett.* **2005**, *7*, 681-684.
- [40] R. M. Moriarty, W. R. Epa, A. K. Awasthi, 'Palladium-Catalyzed Coupling of Alkenyl Iodonium Salts with Olefins: A Mild and Stereoselective Heck-Type Reaction Using Hypervalent Iodine' *J. Am. Chem. Soc.* **1991**, *113*, 6315-6317.
- [41] R. J. Hinkle, G. T. Poulter, P. J. Stang, 'Palladium(II) and Copper(I) Cocatalyzed Coupling of Stereodefined Alkenyl(phenyl)Iodonium Triflates and Unsaturated Tri-*n*-butylstannanes' *J. Am. Chem. Soc.* **1993**, *115*, 11626-11627.
- [42] S.-K. Kang, H.-W. Lee, S.-B. Jang, P.-S. Ho, 'Palladium-Catalyzed Cross-Coupling of Organoboron Compounds with Iodonium Salts and Iodanes' *J. Org. Chem.* **1996**, *61*, 4720-4724.
- [43] S.-K. Kang, T. Yamaguchi, R.-K. Hong, T.-H. Kim, S.-J. Pyun, 'Cross-Coupling and Carbonylative Cross-Coupling of Organofluorosilanes with Hypervalent Iodonium Tetrafluoroborates' *Tetrahedron* **1997**, *53*, 3027-3034.
- [44] S.-K. Kang, K.-H. Lim, P.-S. Ho, W.-Y. Kim, 'Synthesis of Alkynyl Ketones via Palladium- and Copper-Catalyzed Carbonylative Cross Coupling of Iodonium Salts with Alk-1-ynes' *Synthesis* **1997**, *1997*, 874-876.
- [45] N. Sh. Pirgulyev, V. K. Brel, N. S. Zefirov, P. J. Stang, 'Stereoselective Synthesis of Conjugated Alkynes via Palladium-Catalyzed Coupling of Alkenyl Iodonium Salts with Terminal Alkynes' *Tetrahedron* **1999**, *55*, 12377-12386.
- [46] M. Yoshida, A. Komata, S. Hara, 'Stereoselective Synthesis of (Z)- β -Fluoro- α,β -unsaturated Esters from (Z)-2-Fluoro-1-alkenylodonium Salts' *J. Fluor. Chem.* **2004**, *125*, 527-529.
- [47] E. Skucas, D. W. C. MacMillan, 'Enantioselective α -Vinylolation of Aldehydes via the Synergistic Combination of Copper and Amine Catalysis' *J. Am. Chem. Soc.* **2012**, *134*, 9090-9093.
- [48] M. Ochiai, 'Nucleophilic Vinylic Substitutions of λ^3 -Vinylidanes' *J. Organomet. Chem.* **2000**, *611*, 494-508.

- [49] P. J. Stang, 'Alkynyl- and Alkenyl(phenyl)iodonium Compounds. New Synthetic Methods' *Angew. Chem. Int. Ed.* **1992**, *31*, 274-285.
- [50] N. S. Pirkuliev, V. K. Brel, N. S. Zefirov, 'Alkenyl-iodonium salts' *Russ. Chem. Rev.* **2000**, *69*, 105-120.
- [51] T. Okuyama, 'Solvolysis of Vinyl Iodonium Salts. New Insights into Vinyl Cation Intermediates' *Acc. Chem. Res.* **2002**, *35*, 12-18.
- [52] T. Okuyama, M. Fujita, in *Recent Developments in Carbocation and Onium Ion Chemistry* (Ed.: K.K. Laali), ACS, Washington, **2007**, 68.
- [53] A. J. Canty, T. Rodemann, J. H. Ryan, Transition Metal Organometallic Synthesis Utilising Diorganoiodine(III) Reagents. *Adv. Organomet. Chem.* **2007**, *55*, 279.
- [54] D. P. Hari, S. Nicolai, J. Waser, in *PATAI'S Chemistry of Functional Groups* (Ed.: Z. Rappoport), John Wiley & Sons, Ltd, Chichester, UK, **2018**, 36-58.
- [55] E. Stridfeldt, A. Seemann, M. J. Bouma, C. Dey, A. Ertan, B. Olofsson, 'Synthesis, Characterization and Unusual Reactivity of Vinylbenziodoxolones—Novel Hypervalent Iodine Reagents' *Chem. Eur. J.* **2016**, *22*, 16066-16070.
- [56] A. Boelke, L. D. Caspers, B. J. Nachtsheim, 'NH₂-Directed C-H Alkenylation of 2-Vinylanilines with Vinylbenziodoxolones' *Org. Lett.* **2017**, *19*, 5344-5347.
- [57] M. Ochiai, M. Toyonari, T. Nagaoka, D.-W. Chen, M. Kida, 'Stereospecific Synthesis of Vinyl(phenyl)iodonium Tetrafluoroborates via Boron-Iodane Exchange of Vinylboronic Acids and Esters with Hypervalent Phenylidanes' *Tetrahedron Lett.* **1997**, *38*, 6709-6712.
- [58] G. Pisella, A. Gagnebin, J. Waser, 'Copper-Catalyzed Oxyvinylation of Diazo Compounds' *Org. Lett.* **2020**, *22*, 3884-3889.
- [59] M. Ochiai, K. Oshima, Y. Masaki, 'Stereoselective Synthesis of Highly Labile (Z)-β-Alkylvinyl(phenyl)iodonium Perchlorates' *J. Chem. Soc., Chem. Commun.* **1991**, 869-870.
- [60] M. Fujita, H. J. Lee, T. Okuyama, 'Stereochemical Inversion in the Vinylic Substitution of Boronic Esters To Give Iodonium Salts: Participation of the Internal Oxy Group' *Org. Lett.* **2006**, *8*, 1399-1401.
- [61] L. Castoldi, E. M. Di Tommaso, M. Reitti, B. Gräfen, B. Olofsson, 'Electrophilic Vinylation of Thiols under Mild and Transition Metal-Free Conditions' *Angew. Chem., Int. Ed.* **2020**, *59*, 15512-15516.
- [62] P. Caramenti, N. Declas, R. Tessier, M. D. Wodrich, J. Waser, 'Stereoselective Synthesis of Alkyl-, Aryl-, Vinyl- and Alkynyl-Substituted Z-Enamides and Enol Ethers' *Chem. Sci.* **2019**, *10*, 3223-3230.
- [63] B. Liu, J. V. Alegre-Requena, R. S. Paton, G. M. Miyake, 'Unconventional Reactivity of Ethynylbenziodoxolone Reagents and Thiols: Scope and Mechanism' *Chem. Eur. J.* **2020**, *26*, 2386-2394.
- [64] S. G. E. Amos, S. Nicolai, A. Gagnebin, F. Le Vaillant, J. Waser, 'Metal-Free Electrophilic Alkynylation of Sulfenate Anions with Ethynylbenziodoxolone Reagents' *J. Org. Chem.* **2019**, *84*, 3687-3701.
- [65] X. Wang, A. Studer, 'Iodine(III) Reagents in Radical Chemistry' *Acc. Chem. Res.* **2017**, *50*, 1712-1724.
- [66] F. Le Vaillant, J. Waser, 'Alkynylation of Radicals: Spotlight on the "Third Way" to Transfer Triple Bonds' *Chem. Sci.* **2019**, *10*, 8909-8923.
- [67] J. Davies, N. S. Sheikh, D. Leonori, 'Photoredox Imino Functionalizations of Olefins' *Angew. Chem., Int. Ed.* **2017**, *56*, 13361-13365.
- [68] F. Le Vaillant, M. Garreau, S. Nicolai, G. Gryn'ova, C. Corminboeuf, J. Waser, 'Fine-Tuned Organic Photoredox Catalysts for Fragmentation-Alkynylation Cascades of Cyclic Oxime Ethers' *Chem. Sci.* **2018**, *9*, 5883-5889.
- [69] H. Jiang, A. Studer, 'Transition-Metal-Free Three-Component Radical 1,2-Amidoalkynylation of Unactivated Alkenes' *Chem. Eur. J.* **2019**, *25*, 516-520.
- [70] M. Garreau, F. Le Vaillant, J. Waser, 'C-Terminal Bioconjugation of Peptides through Photoredox Catalyzed Decarboxylative Alkynylation' *Angew. Chem., Int. Ed.* **2019**, *58*, 8182-8186.
- [71] K. Matsumoto, M. Nakajima, T. Nemoto, 'Visible Light-Induced Direct S₀ → T₁ Transition of Benzophenone Promotes C(sp³)-H Alkynylation of Ethers and Amides' *J. Org. Chem.* **2020**, *85*, 11802-11811.
- [72] T. Gensch, M. N. Hopkinson, F. Glorius, J. Wencel-Delord, 'Mild Metal-Catalyzed C-H Activation: Examples and Concepts' *Chem. Soc. Rev.* **2016**, *45*, 2900-2936.
- [73] J. Borrel, G. Pisella, J. Waser, 'Copper-Catalyzed Oxyalkynylation of C-S Bonds in Thiiranes and Thiethanes with Hypervalent Iodine Reagents' *Org. Lett.* **2020**, *22*, 422-427.
- [74] A. Boelke, P. Finkbeiner, B. J. Nachtsheim, 'Atom-Economical Group-Transfer Reactions with Hypervalent Iodine Compounds' *Beilstein J. Org. Chem.* **2018**, *14*, 1263-1280.
- [75] G. Grelier, B. Darses, P. Dauban, 'Hypervalent Organoiodine Compounds: from Reagents to Valuable Building Blocks in Synthesis' *Beilstein J. Org. Chem.* **2018**, *14*, 1508-1528.
- [76] D. P. Hari, J. Waser, 'Copper-Catalyzed Oxy-Alkynylation of Diazo Compounds with Hypervalent Iodine Reagents' *J. Am. Chem. Soc.* **2016**, *138*, 2190-2193.
- [77] D. P. Hari, J. Waser, 'Enantioselective Copper-Catalyzed Oxy-Alkynylation of Diazo Compounds' *J. Am. Chem. Soc.* **2017**, *139*, 8420-8423.
- [78] G. Pisella, A. Gagnebin, J. Waser, 'Three-Component Reaction for the Synthesis of Highly Functionalized Propargyl Ethers' *Chem. Eur. J.* **2020**, *26*, 10199-10204.
- [79] S. K. J. Devine, D. L. Van Vranken, 'Palladium-Catalyzed Carbene Insertion into Vinyl Halides and Trapping with Amines' *Org. Lett.* **2007**, *9*, 2047-2049.
- [80] S. K. J. Devine, D. L. Van Vranken, 'Palladium-Catalyzed Carbene Insertion and Trapping with Carbon Nucleophiles' *Org. Lett.* **2008**, *10*, 1909-1911.
- [81] R. Kudirka, S. K. J. Devine, C. S. Adams, D. L. Van Vranken, 'Palladium-Catalyzed Insertion of α-Diazoesters into Vinyl Halides To Generate α,β-Unsaturated γ-Amino Esters' *Angew. Chem., Int. Ed.* **2009**, *48*, 3677-3680.
- [82] T. Kitamura, T. Fukuoka, Y. Fujiwara, 'Synthesis of Novel Vinylbenziodoxolones from Alkynyl(o-carboxyphenyl)iodonium Salts' *Synlett* **1996**, *1996*, 659-660.
- [83] R. Frei, J. Waser, 'A Highly Chemoselective and Practical Alkynylation of Thiols' *J. Am. Chem. Soc.* **2013**, *135*, 9620-9623.
- [84] R. Frei, M. D. Wodrich, D. P. Hari, P.-A. Borin, C. Chauvier, J. Waser, 'Fast and Highly Chemoselective Alkynylation of Thiols with Hypervalent Iodine Reagents Enabled through a Low Energy Barrier Concerted Mechanism' *J. Am. Chem. Soc.* **2014**, *136*, 16563-16573.
- [85] M. Ochiai, M. Kunishima, Y. Nagao, K. Fuji, M. Shiro, E. Fujita, 'Tandem Michael-Carbene Insertion Reactions of Alkynyl-iodonium Salts. Extremely Efficient Cyclopentene Annulations' *J. Am. Chem. Soc.* **1986**, *108*, 8281-8283.
- [86] M. D. Wodrich, P. Caramenti, J. Waser, 'Alkynylation of Thiols with Ethynylbenziodoxolone (EBX) Reagents: α- or β-π-Addition?' *Org. Lett.* **2016**, *18*, 60-63.
- [87] R. Tessier, J. Ceballos, N. Guidotti, R. Simonet-Davin, B. Fierz, J. Waser, '"Doubly Orthogonal" Labeling of Peptides and Proteins' *Chem* **2019**, *5*, 2243-2263.

- [88] J. Wu, X. Deng, H. Hirao, N. Yoshikai, 'Pd-Catalyzed Conversion of Alkynyl- λ^3 -iodanes to Alkenyl- λ^3 -iodanes via Stereoselective 1,2-Iodine(III) Shift/1,1-Hydrocarboxylation' *J. Am. Chem. Soc.* **2016**, *138*, 9105-9108.
- [89] J. Wu, K. Xu, H. Hirao, N. Yoshikai, 'Pd-Catalyzed, Ligand-Enabled Stereoselective 1,2-Iodine(III) Shift/1,1-Carboxyalkynylation of Alkynylbenziodoxoles' *Chem. Eur. J.* **2017**, *23*, 1521-1525.
- [90] B. Liu, C.-H. Lim, G. M. Miyake, 'Light-Driven Intermolecular Charge Transfer Induced Reactivity of Ethynylbenziodoxol(on)e and Phenols' *J. Am. Chem. Soc.* **2018**, *140*, 12829-12835.
- [91] Y. Cheng, S. Yu, 'Hydrotrifluoromethylation of Unactivated Alkenes and Alkynes Enabled by an Electron-Donor-Acceptor Complex of Togni's Reagent with a Tertiary Amine' *Org. Lett.* **2016**, *18*, 2962-2965.
- [92] T. Aubineau, J. Cossy, 'Chemoselective alkynylation of *N*-sulfonylamides versus amides and carbamates – Synthesis of tetrahydropyrazines' *Chem. Commun.* **2013**, *49*, 3303-3305.
- [93] D. Shimbo, A. Shibata, M. Yudasaka, T. Maruyama, N. Tada, B. Uno, A. Itoh, 'Synthesis of *cis*- β -Amidevinyl Benziodoxolones from the Ethynyl Benziodoxolone-Chloroform Complex and Sulfonamides' *Org. Lett.* **2019**, *21*, 9769-9773.
- [94] W. Ding, J. Chai, C. Wang, J. Wu, N. Yoshikai, 'Stereoselective Access to Highly Substituted Vinyl Ethers via *trans*-Difunctionalization of Alkynes with Alcohols and Iodine(III) Electrophile' *J. Am. Chem. Soc.* **2020**, *142*, 8619-8324.
- [95] J. Chai, W. Ding, J. Wu, N. Yoshikai, 'Fluorobenziodoxole-BF₃ Reagent for Iodo(III)etherification of Alkynes in Ethereal Solvent' *Chem. Asian J.* **2020**, *15*, 2166-2169.
- [96] M. Li, W. Li, C.-D. Lin, J.-H. Wang, L.-R. Wen, 'One Base for Two Shots: Metal-Free Substituent-Controlled Synthesis of Two Kinds of Oxadiazine Derivatives from Alkynylbenziodoxolones and Amidoximes' *J. Org. Chem.* **2019**, *84*, 6904-6915.
- [97] J. Wu, X. Deng, N. Yoshikai, 'Stereocontrolled Synthesis of Halovinylbenziodoxoles by Hydro- and Iodochlorination of Ethynylbenziodoxoles' *Chem. Eur. J.* **2019**, *25*, 7839-1842.
- [98] B. Lu, J. Wu, N. Yoshikai, 'Palladium-Catalyzed Condensation of *N*-Aryl Imines and Alkynylbenziodoxolones To Form Multisubstituted Furans' *J. Am. Chem. Soc.* **2014**, *136*, 11598-11601.
- [99] A. Ariafard, 'Computational Mechanistic Study of Palladium(II)-Catalyzed Carboxyalkynylation of an Olefin Using an Iodine(III) Oxidant Reagent' *Organometallics* **2014**, *33*, 7318-7324.
- [100] T. Okuyama, T. Takino, T. Sueda, M. Ochiai, 'Solvolysis of Cyclohexenylidonium Salt, a New Precursor for the Vinyl Cation: Remarkable Nucleofugality of the Phenylidonium Group and Evidence for Internal Return from an Intimate Ion-Molecule Pair' *J. Am. Chem. Soc.* **1995**, *117*, 3360-3367.
- [101] N. Declas, J. Waser, 'Access to Vinyl Ethers and Ketones with Hypervalent Iodine Reagents as Oxy-Allyl Cation Synthetic Equivalents' *Angew. Chem., Int. Ed.* **2020**, *59*, 18256-18260.

Entry for the Table of Contents

Twitter

This review from the Waser group @LcsoLab @EPFL_CHEM_Tweet describes the recent rise of vinylbenziodoxolone (VBX) reagents as useful tools as electrophilic alkene synthons in organic chemistry.

Biosketches:

Guillaume Pisella studied chemistry at Université Paris-Saclay and at Chimie ParisTech. During his studies, he moved twice in the Basel area to join Syngenta as a trainee in process research and the Roche pRED team as a medicinal chemistry intern. In 2016, he graduated with a Master Degree in Chemistry, having worked in the Lautens group at the University of Toronto. He then moved back to Switzerland for his doctoral studies in the Waser group in January 2017, working on the development and applications of hypervalent iodine reagents in copper-catalyzed transformations.

Jérôme Waser was born in Sierre, Valais, Switzerland in 1977. He obtained his chemistry Diploma at ETH Zurich in 2001. From 2002 to 2006, he was a PhD student at ETH Zurich with Prof. Erick M. Carreira. He then joined Prof. Barry M. Trost at Stanford University as a SNF postdoctoral fellow. From 2007 to 2014, he was an assistant professor at EPF Lausanne (EPFL). Since June 2014, he has been associate professor at EPFL. He is a recipient of the ERC Starting Grant 2013 and Consolidator Grant 2017, the Werner prize of the Swiss Chemical Society 2014 and the Springer Heterocyclic Chemistry Award 2016.

Nina Declas was born in Lyon (France) where she studied chemistry at the Ecole d'Ingénieurs en Chimie et Sciences du Numérique CPE Lyon. In 2016, she joined the Broad Institute of MIT and Harvard in Boston (Massachusetts), to perform a one-year internship as a medicinal chemist. After receiving her Master degree in 2018, she joined the group of Prof. Jérôme Waser for her doctoral studies. Her research is focus on the development and applications of hypervalent iodine reagents, and more precisely vinylbenziodoxolone reagents.