



Towards understanding the characteristics of new particle formation in the Eastern Mediterranean

Rima Baalbaki¹, Michael Pikridas², Tuija Jokinen¹, Tiia Laurila¹, Lubna Dada^{1,a,b}, Spyros Bezantakos², Lauri Ahonen¹, Kimmo Neitola^{1,2}, Anne Maisser², Elie Bimenyimana², Aliko Christodoulou^{2,3}, Florin Ungu², Chrysanthos Savvides⁴, Katrianne Lehtipalo^{1,5}, Juha Kangasluoma¹, George Biskos², Tuukka Petäjä¹, Veli-Matti Kerminen¹, Jean Sciare², and Markku Kulmala¹

¹Institute for Atmospheric and Earth System Research (INAR)/Physics, Faculty of Science, University of Helsinki, P.O. Box 64, Helsinki, 00014, Finland

²Climate & Atmosphere Research Centre (CARE-C), The Cyprus Institute, P.O. Box 27456, Nicosia, 1645, Cyprus

³SAGE – Département Sciences de L'Atmosphère et Génie de L'Environnement, IMT Lille Douai, Université de Lille, 59000, Lille, France

⁴Ministry of Labour, Welfare and Social Insurance, Department of Labour Inspection (DLI), Nicosia, Cyprus

⁵Atmospheric Composition Research Unit, Finnish Meteorological Institute, P.O. Box 503, Helsinki, 00101, Finland

^anow at: EPFL, School of Architecture, Civil and Environmental Engineering, Sion, 1951, Switzerland

^bnow at: Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, Villigen, 5232, Switzerland

Correspondence: Rima Baalbaki (rima.baalbaki@helsinki.fi)

Received: 12 October 2020 – Discussion started: 18 November 2020

Revised: 23 April 2021 – Accepted: 4 May 2021 – Published: 16 June 2021

Abstract. To quantify the contribution of new particle formation (NPF) to ultrafine particle number and cloud condensation nuclei (CCN) budgets, one has to understand the mechanisms that govern NPF in different environments and its temporal extent. Here, we study NPF in Cyprus, an Eastern Mediterranean country located at the crossroads of three continents and affected by diverse air masses originating from continental, maritime, and desert-dust source areas. We performed 1-year continuous measurements of aerosol particles down to ~ 1 nm in diameter for the first time in the Eastern Mediterranean and Middle East (EMME) region. These measurements were complemented with trace gas data, meteorological variables, and retroplume analysis. We show that NPF is a very frequent phenomenon at this site and has higher frequencies of occurrence during spring and autumn. NPF events were both of local and regional origin, and the local events occurred frequently during the month with the lowest NPF frequency. Some NPF events exhibited multiple onsets, while others exhibited apparent particle shrinkage in size. Additionally, NPF events were observed during the nighttime and during episodes of high desert-dust loadings. Particle formation rates and growth rates were comparable to those in

urban environments, although our site is a rural one. Meteorological variables and trace gases played a role in explaining the intra-monthly variability of NPF events, but they did not explain why summer months had the least NPF frequency. Similarly, pre-existing aerosol loading did not explain the observed seasonality. The months with the least NPF frequency were associated with higher H_2SO_4 concentrations but lower NO_2 concentrations, which is an indicator of anthropogenic influence. Air masses arriving from the Middle East were not observed during these months, which could suggest that precursor vapors important for nucleation and growth are transported to our site from the Middle East. Further comprehensive measurements of precursor vapors are required to prove this hypothesis.

1 Introduction

Atmospheric new particle formation (NPF) is the process by which oxidized precursor gases initially form molecular clusters that then further grow in size by multi-component condensation (Kulmala et al., 2014). A multitude of research

studies have focused on this phenomenon over the past 2 decades, because it is a large source of the global aerosol particle number and cloud condensation nuclei (CCN) load (Gordon et al., 2017; Merikanto et al., 2009; Pierce and Adams, 2009; Wang and Penner, 2009; Yu and Luo, 2009; Kerminen et al., 2012; Spracklen et al., 2006, 2008). Owing to the complex nature and nonlinearity of atmospheric processes, studies on NPF in the literature include atmospheric observations (e.g., Kulmala et al., 2013; Ehn et al., 2014; Bianchi et al., 2016; Yao et al., 2018; Williamson et al., 2019; Baccarini et al., 2020; Dall'Osto et al., 2018), chamber experiments (e.g., Sipilä et al., 2010; Tröstl et al., 2016; Wang et al., 2020; Lehtipalo et al., 2016; Kirkby et al., 2011), and theoretical computational studies (e.g., Kurtén et al., 2008; Riipinen et al., 2011; Olenius and Riipinen, 2017). The collective scientific outcome from these studies is essential to understand the mechanisms and characteristics of NPF (Kerminen et al., 2018; Lee et al., 2019; Chu et al., 2019) and how it affects the global climate (e.g., Spracklen et al., 2006; Gordon et al., 2017).

The frequency, strength, and spatiotemporal extent of NPF are mainly governed by three factors: the prevailing meteorological conditions, the availability of gaseous precursors, and the pre-existing concentrations of aerosol particles (Kerminen et al., 2018; Lee et al., 2019; Nieminen et al., 2018). These atmospheric conditions differ in space and time. Atmospheric conditions are distinct over the Mediterranean basin, especially over the Eastern Mediterranean and Middle East (EMME). This region has been identified as a hotspot for atmospheric and climate change research (Lelieveld et al., 2016; Giorgi and Lionello, 2008). It is surrounded by three continents and is affected by continental, maritime, and desert-dust pollution sources (Lelieveld et al., 2002). The surrounding complex orography of the Mediterranean affects atmospheric dynamics and boundary layer processes on different scales (Kostopoulou and Jones, 2007a, b). Further, the dry and hot weather throughout most of the year, with strongly increasing heat extremes, enables intense photochemistry (Lelieveld et al., 2016).

NPF studies over the Mediterranean have focused on the northwestern basin (Petäjä et al., 2007; Cusack et al., 2013; Berland et al., 2017; Carnerero et al., 2018; Rose et al., 2015; Brines et al., 2015; Hamed et al., 2007; Laaksonen et al., 2005; Casquero-Vera et al., 2020), whereas NPF studies in the eastern basin have been conducted mainly in Greece (Petäjä et al., 2007; Berland et al., 2017; Kalivitis et al., 2015, 2019; Pikridas et al., 2012; Kalkavouras et al., 2019, 2020; Kopanakis et al., 2013; Siakavaras et al., 2016; Kalkavouras et al., 2017) and very recently in Cyprus (Brilke et al., 2020; Debevec et al., 2018) and Jordan (Hussein et al., 2020). These studies include both short-term campaigns and long-term observation.

Based on long-term measurements, the annual frequency of NPF over the Mediterranean varies between 10 % and 36 % (Hussein et al., 2020; Kalivitis et al., 2019; Kalkavouras

et al., 2020; Kopanakis et al., 2013). The seasonal cycle has a typical maximum during spring (Kalkavouras et al., 2020; Kopanakis et al., 2013; Kalivitis et al., 2019; Pikridas et al., 2012), even though in some urban background sites the highest frequency was observed during summer (Hussein et al., 2020; Hamed et al., 2007). NPF was associated with a high increase in nucleation mode particles in most of the studies. For instance, Carnerero et al. (2018) showed that the impact of NPF on ultrafine particles is much higher than that of traffic near the highly polluted city center of Madrid. The condensation sink, which is a measure of the pre-existing aerosol surface area, was reported to be lower during NPF events in Po Valley, Corsica, and Crete (Hamed et al., 2007; Berland et al., 2017; Pikridas et al., 2012), while NPF proceeded under both clean and polluted conditions in Barcelona (Cusack et al., 2013), Marseille, and Athens (Petäjä et al., 2007). The effect of meteorological conditions on NPF occurrence varied among studies. Simultaneous NPF events were observed in several stations, illustrating that the spatial extent of NPF events can vary from tens of kilometers (Carnerero et al., 2018) to several hundred kilometers (Kalkavouras et al., 2017, 2020; Berland et al., 2017; Rose et al., 2015; Casquero-Vera et al., 2020). In the Po Valley, the production of CCN from NPF was estimated to be comparable to that originating from primary sources (Laaksonen et al., 2005). Similarly, NPF was associated with a strong increase in CCN concentrations in Finokalia and Santorini (Kalkavouras et al., 2017, 2019; Kalivitis et al., 2015). However, the impact of the increased CCN concentrations on cloud droplet number was shown to be limited by water availability (Kalkavouras et al., 2017). In Cyprus, mainly in Paphos, Gong et al. (2019) observed several NPF events where newly formed particles grew into the CCN size range, with NPF events being observed on 9 out of 27 measurement days during April 2017 (Brilke et al., 2020). At a more inland site, NPF was observed on 14 out of 20 d of measurements in March 2015 (Debevec et al., 2018). Since these studies were less than a month long, further comprehensive measurements are required to unveil the role of NPF in the atmospheric processes taking place in the EMME region.

The aim of this study is to characterize the seasonal cycle of new particle formation events in the less represented area of the EMME region. Our measurements were conducted at a rural background site on the island of Cyprus, which lies at the crossroads of three continents in the Eastern Mediterranean. We report the first long-term analysis of particle number size distribution in the area down to sizes where the initial formation occurs. We further explore the role of sulfuric acid, which is one of the key gas-phase precursors for cluster formation, and other atmospheric variables in initiating NPF at this site.

2 Materials and methods

2.1 Measurement site

Cyprus is an island country in the Eastern Mediterranean. It is the third most populous island in the Mediterranean Sea and the third largest in size with an area of 9251 km² (Fig. 1). The measurements reported in this work were carried out at the Agia Marina Xyliatos station of the Cyprus Atmospheric Observatory (CAO-AMX; Sciare, 2016), which is a rural background station that operates under the co-operative program for monitoring and evaluation of the long-range transmission of air pollutants in Europe (EMEP) and the European Research Infrastructure for the observation of Aerosols, Clouds and Trace gases Research Infrastructure (ACTRIS) networks, while at the same time it is a designated regional Global Atmospheric Watch (GAW) station. The station (35.038692° N, 33.057850° E) is located close to the villages of Agia Marina (~ 630 inhabitants) and Xyliatos (~ 150 inhabitants) and has an elevation of 532 m above sea level. The site is surrounded by vegetation, mainly oak and pine trees and Maquis shrubland, as it lies at the northeastern foothills of the Troodos Mountains. Agriculture areas surround the site from the north direction and are approximately 4 km away. The nearest main urban agglomeration is at least 35 km away. Therefore, it is not directly affected by any major local pollution source, excluding some limited traffic to reach the nearby Forestry Department premises.

The weather at CAO-AMX is characterized by hot, dry summers and mild, rainy winters. The daily mean temperature is ~ 19 °C and ranges between 1 and 36 °C, the daily mean relative humidity is ~ 55 % and ranges between 13 % and 82 %, and the daily mean ozone level is ~ 48 ppb and ranges between 26 and 77 ppb (Kleanthous et al., 2014). The most common (> 65 % occurrence) wind pattern reaching the site is the northerly “Etesian” winds transporting pollutants from both Europe and Turkey, but winds are more frequently arriving from the mainland of Turkey (Pikridas et al., 2018). The remaining air masses originate from northern Africa, the Middle East, and westerly air masses that spend several days above the sea before reaching Cyprus. The variable air mass origins at CAO-AMX from three different continents allow for a representative description of NPF processes for the EMME region as a whole.

2.2 Instrumentation

2.2.1 Aerosol particle number size distribution

The particle number size distribution between 1 and 700 nm was determined by combining data from three instruments: an Airmodus A11 Nano condensation nucleus counter (nCNC) system (Vanhanen et al., 2011), a neutral cluster and air ion spectrometer (NAIS, model 1; Manninen et al., 2016; Mirme and Mirme, 2013), and a scanning mobility parti-

cle sizer (SMPS, model TSI 3080; Wang and Flagan, 1990). The first two instruments were operated at the site for a period of 1 year from 27 January 2018 to 26 January 2019, while the SMPS measurement period was from 27 January to 1 November 2018. The monthly availability of data from each instrument is shown in Table S1.

The A11 nCNC is composed of a particle size magnifier (PSM; Airmodus A10) and a condensation particle counter (CPC; Airmodus A20). The overall length of the inlet sampling tube was 60 cm. The PSM was operated in a scanning saturator flow mode between 0.1 and 1.3 L min⁻¹, corresponding to a cutoff diameter range of approximately 1.1 to 2.5 nm. It was equipped with an inlet system that performs background (zero) measurements three times a day at random time intervals and a core sampling piece for minimizing line losses of sub-3 nm particles (Fig. S1). The duration of the background measurements was set to 12 min, which is equivalent to three full size scans. From June 2018 onwards, the nCNC was additionally equipped with a diluter to reduce the humidity of the sampled air. This procedure was necessary because the water content of the air at the measurement site was too high. The water present in the sample air was mixed with butanol inside the CPC of the nCNC and rendered it measuring zeros. Further information about the diluter design, its operation, and effect on the data can be found in the Supplement Sect. S2.2.

The NAIS is a mobility spectrometer designed to determine the number size distribution of ions in the mobility diameter range of 0.8–42 nm, as well as total (naturally charged and neutral) aerosol particles in the mobility diameter range of ~ 2–42 nm. The instrument operates at a flow rate of ~ 54 L min⁻¹. The length of the NAIS sampling tube was 65 cm, with an inner diameter of 30 mm.

The SMPS used in this study was composed of a TSI 3081 long differential mobility analyzer (DMA) and a TSI 3025a CPC. It was operated to measure the aerosol particle size distribution between 15 and 740 nm. The aerosol and sheath flows were checked weekly and were set to 0.3 and 3 L min⁻¹, respectively. The SMPS was sampling using an 80 cm long vertical inlet. Drying was achieved using a short Nafion dryer, and charge neutralization was achieved by a GRIMM 5522-A, americium-241, bipolar neutralizer.

2.2.2 Ancillary measurements

Complementary meteorological data (temperature, relative humidity, solar radiation, rainfall, pressure, wind speed, and wind direction) were measured with a time resolution of 5 min at an elevation of 10 m from the ground in the nearby village of Xyliatos (35.0140917° N, 33.0492028° E), located 2.85 km from the measurement site. Air pollutants (ozone, carbon monoxide, nitrogen oxides, sulfur oxide, PM₁₀, and PM_{2.5}) were measured at the colocated EMEP station ~ 20 m from the main measurement container, and these data had a time resolution of 1 h. Additional details about the set-




Figure 1. Maps of the Mediterranean region and Cyprus. (a) Location of Cyprus in the Mediterranean region. (b) Location of the measurement site (CAO-AMX) in Cyprus. (c) Location of the measurement site (CAO-AMX) pointed out by the blue location marker with respect to the villages of Agia Marina and Xyliatos. The geographic borders of the villages are marked by the yellow lines. The maps were retrieved from Google (© 2020 Google, TerraMetrics).

ups and the instruments used can be found in Kleanthous et al. (2014) and Pikridas et al. (2018).

2.3 Data handling

nCNC. The scanning *nCNC* data were inverted into a size distribution with the kernel inversion method presented by Lehtipalo et al. (2014), but using customized kernels which follow the instrument-specific detection efficiency calibration curves. The following diameters were used in the inversion: 1.1, 1.3, 1.5, and 2.4 nm. The choice of the inversion method was made after a comprehensive comparison between the kernel method and the expectation and minimization (EM) method (Cai et al., 2018; Chan et al., 2020). Additional details about the comparability of the two methods and the utilized inversion parameters are presented in Sect. S2.3. After inversion, the data were further corrected for line losses using the method suggested by Fu et al. (2019) for the sampling line downstream of the core sampling inlet and using the Gormley and Kennedy equation for the line losses inside the 6 cm long core sampling piece (Gormley and Kennedy, 1948).

NAIS. The NAIS data were inverted with the instrument-specific algorithm (done by the NAIS SPECTOPS software). The data were later corrected for line losses using the Gorm-

ley and Kennedy equation for laminar flow (Gormley and Kennedy, 1948). It is essential to note that the flow through the sampling inlet of the NAIS actually lies in the transient regime ($Re = 2376$); however, the penetration efficiency using this inlet was comparable for laminar flow and turbulent flow (calculated using the equation of turbulent inertial deposition from Brockmann, 2011); thus, we used the correction based on laminar flow (Fig. S4).

SMPS. The data from the SMPS were inverted using TSI's Aerosol Instrument Manager software (AIM, version 9.0). Afterwards, line loss correction was applied using the Gormley and Kennedy equation. Additional corrections based on laboratory calibrations were also applied to account for the CPC detection efficiency curves.

Full particle size distribution (PSD). The data from the three particle sizing instruments were used to reconstruct the full particle size distribution with a temporal resolution of 5 min between 1.1 and 736 nm (*nCNC*: 1.1 to 2.4 nm; NAIS particle mode: 2.4 to 30 nm; SMPS: 30 to 736 nm). However, the SMPS measured dry aerosol particle number distributions, which can differ from the ambient aerosol particle number size distribution. Thus, we back-calculated the distribution of the SMPS at ambient conditions from the dry distribution using the hygroscopicity model of Petters and Krei-

denweis (2007) and mean κ values. Additional information about these calculations and its effect on sink calculations are presented in Sect. S4. The SMPS distribution at ambient conditions was reconstructed up to 1500 nm. This does not imply that the measurement range was extended to 1500 nm but rather that now we account for particles that were originally of sizes up to 1500 nm but were dried to sizes below 736 nm in the SMPS sampling line. Additionally, since the NAIS is known to overestimate concentrations in the particle mode, the overlapping measurement range with the SMPS was used to further correct the NAIS data, assuming that the NAIS overestimate concentrations uniformly over the whole measurement range, which is a reasonable assumption for old NAIS models based on calibration results (Gagné et al., 2011; Kangasluoma et al., 2020). Finally, the PSD data were run through a 2D median filtering algorithm with a 3-by-3 neighborhood window. Moreover, the data were manually checked for the success of the outlier and noise removal techniques.

Complementary data. Gas and meteorology data sets were run through an outlier removal algorithm and filtered for erroneous samples. The outlier detection method was based on removing data points that are more than 3 standard deviations from a moving median (Davies and Gather, 1993; Pearson et al., 2016).

2.4 Event classification

The reconstructed full particle size distribution daily plots were used to categorize measurement days into NPF event days, non-event days, and undefined days based on a classification that combines the schemes reported in literature (Dal Maso et al., 2005; Hirsikko et al., 2007; Manninen et al., 2010; Kulmala et al., 2012). The classification of events utilizing PSD data that extends below 10 nm, which is a typical measurement limit for most SMPS systems, improves the event classification and allows for better identification of event days that would otherwise be classified as undefined or non-events if only PSDs above 10 nm were used (Leino et al., 2016; Dada et al., 2018; Brilke et al., 2020). In addition, spectra of total particles (both neutral and charged) are usually easier to visually classify than those corresponding to charged particles (measured by the ion mode of NAIS), because atmospheric nucleation is dominated by neutral processes (Kontkanen et al., 2013; Kulmala et al., 2013; Wagner et al., 2017). Furthermore, the concentration of the growing mode in the charged spectra is lower for the smaller particle sizes and increases with diameter as the probability of cluster ions attaching to the growing neutral particles increases (Gonser et al., 2014). Thus, it could be visually difficult to determine if particle nucleation starts from the smallest sizes when looking at the charged spectra only. In contrast, one should not neglect to look at the charged spectra, because these might show sign preference or ion-induced nucleation events (Rose et al., 2018).

2.5 NPF-specific parameters

Condensation sink (CS) is a loss term for condensable vapors used to describe their loss rate by condensation to pre-existing aerosol surface. This term was first introduced by Kulmala et al. (2001), and it is derived based on condensing vapor mass flux to the particles in the continuum regime and applying the transitional correction factor (β_m) proposed by Fuchs and Sutugin (1971):

$$CS = 4\pi D \sum_i \beta_{m_i} r_i N_i = 2\pi D \sum_i \beta_{m_i} d_{p_i} N_i, \quad (1)$$

where r , d_p , and N are the particle radius, diameter, and number concentration, respectively, in the size class i , and D is the diffusion coefficient of the condensing vapor calculated as recommended by Fuller et al. (1966):

$$D(\text{H}_2\text{SO}_4, \text{air}) = \frac{0.001T^{1.75} \sqrt{\frac{1}{M_{\text{H}_2\text{SO}_4}} + \frac{1}{M_{\text{air}}}}}{P (\sqrt[3]{V_{\text{H}_2\text{SO}_4}} + \sqrt[3]{V_{\text{air}}})^2}, \quad (2)$$

where T is the temperature, M is the molar mass, P is the atmospheric pressure, and V is the diffusion volume. Here, CS was calculated assuming that sulfuric acid is the main condensing vapor.

Coagulation sink (CoagS) is a loss term for freshly formed particles used to describe their loss rate by Brownian coagulation to pre-existing aerosol surface (Kulmala et al., 2001). It is calculated as

$$\text{CoagS}(d_p) = \sum_j K_{ij} N_j, \quad (3)$$

where K_{ij} is the Fuchs form of the Brownian coagulation coefficient (Fuchs, 1964; Seinfeld and Pandis, 2012).

Apparent growth rate (GR) is the rate of change in the diameter, d_p , that represents the growing-particle population. It was calculated here using the NAIS data for negatively charged ions, positively charged ions, and total particles (charged + neutral) by the appearance time method (Lehtipalo et al., 2014). First, the time to reach 50% of the maximum concentration is determined, and then the growth rate is derived as the slope of the linear fit between the diameters and time:

$$\text{GR} = \frac{dd_p}{dt} = \frac{\Delta d_p}{\Delta t}. \quad (4)$$

We calculated GR at three different size ranges: between 1.5 and 3 nm ($\text{GR}_{1.5-3}$), between 3 and 7 nm (GR_{3-7}), and between 7 and 20 nm (GR_{7-20}).

Event start and end times were determined based on the time evolution of the 2–4 nm particles, which is the size range suggested by Dada et al. (2018). Using this size range, we are able to capture the early stages of the event, which is unachievable if the measured PSD starts from bigger sizes. Thus, computed event start and end times might differ across

studies depending on the instrument used. An event start is determined by an increase in the 2–4 nm particle concentration above the nighttime level which lasts for at least an hour. An event end time is determined when the 2–4 nm particle concentration decreases to background levels. In case of multiple events within a 1 d window, the event start and end times were taken from the start of the first event and the end of the last event, respectively.

Particle formation rate (J) is the rate at which aerosol particles of certain size are formed in the atmosphere. It quantifies the intensity of the NPF events, and it is calculated by rearranging the equation describing the time evolution of the particle number concentration (Kulmala et al., 2012). d_p in this equation refers to the smaller limit of the size bin used in the calculation of the formation rate. We calculated J at three sizes: 1.5 nm ($J_{1.5}$), 3 nm (J_3), and 7 nm (J_7); the upper size limits used were 3, 7, and 20 nm, respectively. GR was calculated as the mean of the three GR measurements (negative ions, positive ions, and total particles) and was considered constant within the event start and end times. Outside the event times and during non-events, the GR term was considered equal to zero.

$$J_{d_p} = \frac{dN_{d_p}}{dt} + \text{CoagSN}_{d_p} + \frac{\text{GR}}{\Delta d_p} N_{d_p} \quad (5)$$

The first term on the right-hand side of Eq. (5) represents the time evolution of particle number concentration N_{d_p} , the second term represents the coagulation losses due to larger aerosol particles, and the third term represents the condensational growth to sizes bigger than the considered size range.

2.6 Sulfuric acid proxy

Sulfuric acid is one of the key gas-phase compounds identified to contribute to new particle formation (e.g., Weber et al., 1996; Sipilä et al., 2010). As direct measurements of sulfuric acid are challenging, a suite of proxies for the sulfuric acid concentrations are derived that facilitate the calculation of gas-phase sulfuric acid from ancillary observations (Dada et al., 2020; Mikkonen et al., 2011; Petäjä et al., 2009; Lu et al., 2019; Weber et al., 1997). In this study, the sulfuric acid proxy was calculated using the new method by Dada et al. (2020) for a rural site, which was developed based on observations from the same site of this study:

$$[\text{H}_2\text{SO}_4]_{\text{rural}} = -\frac{\text{CS}}{2 \times (2 \times 10^{-9})} + \left[\left(\frac{\text{CS}}{2 \times (2 \times 10^{-9})} \right)^2 + \frac{[\text{SO}_2]}{(2 \times 10^{-9})} (9 \times 10^{-9} \times \text{GlobRad}) \right]^{\frac{1}{2}}. \quad (6)$$

This proxy not only considers the formation of H_2SO_4 from SO_2 via OH oxidation and the loss of H_2SO_4 onto pre-

existing particles (condensation sink) but also includes loss of H_2SO_4 via atmospheric clustering starting from H_2SO_4 dimer formation.

2.7 Air mass origin analysis

Air mass origins for the entire measurement period were modeled using the Lagrangian particle dispersion model FLEXPART (FLEXible PARTicle dispersion model, version 8.23) in a backward mode (Stohl et al., 2005), with meteorological ($0.5^\circ \times 0.5^\circ$, 6 h starting from midnight UTC) National Center for Atmospheric Research (NCAR, ds 0.94) data as input. We used “species 1” (tracer), which does not include wet or dry deposition and assumes an infinite lifetime for the particles, as the tracer released to model the retroplumes. Retroplumes replace simple back-trajectory calculations in the interpretation of atmospheric trace substance measurements, and were traced back in time for 5 d using CAO-AMX as the receptor site. Air masses were categorized into source regions based on the potential emission sensitivity (PES) for the lowest 1 km above ground level (a.g.l.), following the classification method of Pikridas et al. (2010). In general, a retroplume was attributed to a region in the case that this had a PES value above 0.9 ns kg^{-1} . The classification scheme of the source regions took into consideration dominant air mass paths shown by Pikridas et al. (2018) and the different sources of PM with characteristic chemical fingerprint. As a result, the predominant northerly air masses were categorized into “Europe” and “NW Asia” (namely, Turkey), assuming different emissions related to SO_2 . “N. Africa” and “SW Asia” are both source areas of dust particles but with distinct emission levels, with the former being associated with more elevated concentrations. The “Asia sector” was distinguished to point out that air masses from this specific source region scarcely reach the receptor site, while the source region “Local” refers to stagnant conditions. Last, the “Marine” sector is associated with the lowest levels of ambient PM. In total, seven source regions were identified similar to the ones presented by Pikridas et al. (2018), except that in our analysis the “West Turkey” sector was merged to the NW Asia sector.

2.8 Identification of days with high dust loading

Measurement dates with high dust loading were identified using the VI-PM1 online method proposed by Drinovec et al. (2020). This method couples a high-flow virtual impactor (VI) sampler, which concentrates coarse particles, with an aerosol absorption photometer. More details about the calculations and a list of the identified dust days can be found in Sect. S5.

3 Results and discussion

In the course of identifying NPF events, the PSD spectrum is usually analyzed, mainly at sizes below 25 nm where one can detect the emergence of new aerosol particles, and then the particle growth to larger sizes is followed. Since little is known about particle number size distributions from the EMME region, we will first present the seasonal and diurnal variability of particle number concentration in different PSD modes (Sect. 3.1). Then, we will identify and characterize NPF events (Sect. 3.2). Following that, we will quantify and analyze relevant parameters that describe NPF events (Sect. 3.3) and use those parameters, together with meteorological variables, to understand why and when NPF occurs at our site (Sect. 3.4). We further present a regression and a classification analysis (Sect. 3.5). All the data in this article are presented in local time (UTC+3 from 25 March to 28 October 2018 and UTC+2 during the rest of the campaign). Unless otherwise indicated, we mainly focus on daytime data having global radiation $> 50 \text{ W m}^{-2}$ as this is the time period relevant for most NPF events, but we also briefly mention some nighttime events. For reference, the monthly range of day hours having global radiation $> 50 \text{ W m}^{-2}$ is presented in Fig. S6.

3.1 Seasonal and diurnal variability of number concentration in different modes

Figure 2 presents the monthly percentiles boxplots (25th, 50th, and 75th) and the mean averages of the cluster mode ($\sim 1\text{--}3 \text{ nm}$), nucleation mode ($3\text{--}25 \text{ nm}$), Aitken mode ($25\text{--}100 \text{ nm}$), and accumulation mode ($100\text{--}1000 \text{ nm}$) particle number concentrations computed from daily data with global radiation $> 50 \text{ W m}^{-2}$ (daytime conditions). A clear seasonal pattern is depicted which is distinct across the different particle modes. The cluster mode particles had two peaks, one in spring and another in autumn with a clear drop during the summer. The monthly boxplots also show high variability in daily concentrations throughout most months, except August and September. This variability seems highest during February. The nucleation mode particles were also highest during the spring and lowest during the summer. The autumn concentrations did not exhibit another peak but were rather similar to the summer concentrations in terms of median values, but they exhibited higher variability. The cluster and nucleation mode concentrations can be directly linked to the NPF activity, especially in sites where direct emissions of particles having these size ranges are minimal, which is the case for our site. While the high concentration of cluster mode particles during spring was associated with high concentration of nucleation mode particles, this did not hold for autumn, which might indicate that condensable vapors were not as available to grow the particles to nucleation size. The Aitken mode exhibited higher concentrations during the spring months followed by a decreasing pattern, which could

either suggest more growth from NPF to Aitken sizes or higher emission and transportation of primary particles during spring. The accumulation mode had its maximum during the warm months, except during July which did not follow the pattern of other months. Previous long-term measurements of $\text{PM}_{2.5}$ at this site have a similar pattern with higher concentrations during the warm period of the year and minimum during winter (Pikridas et al., 2018). This maximum during the summer is mainly explained by the enhanced transport of polluted air masses from the northern sector, combined with the lack of precipitation and overall dry conditions during Eastern Mediterranean summer (Pikridas et al., 2018). Lastly, it is worth mentioning that during February the concentrations of particles in all modes did not follow the overall trend. It exhibited lower concentrations of cluster, nucleation, and Aitken mode particles and higher concentration of accumulation mode particles than the before-and-after months.

The diurnal variation (at radiation $> 50 \text{ W m}^{-2}$) of the cluster and nucleation mode particles exhibited a clear cycle, with the highest values recorded between 09:00 and 15:00 LT and the maximum at 11:00 LT (Fig. 3a and b). There was a slight time difference between the appearance of the cluster mode particles and that of the nucleation mode particles which could only be seen in the 5 min data. The Aitken mode had a less distinct diurnal cycle, having the peak at later hours of the day, which might indicate that these particles have possibly grown from the cluster and nucleation modes (Fig. 3c). The accumulation mode, on the other hand, did not exhibit any clear diurnal cycle, which might suggest that these particles are not emitted or produced from any local source but are rather transported over a long range. They can be aged primary particles or particles originating from NPF taken place 1–3 d earlier in arriving air masses (Fig. 3d).

3.2 General character of the NPF events

New particle formation has been detected to occur in a variety of environments within the troposphere (Kerminen et al., 2018; Lee et al., 2019; Nieminen et al., 2018). Typically, the appearance of clusters is detected in the morning hours followed by subsequent growth. The occurrence of new particle formation events is determined by examining the time evolution of the aerosol number size distributions (e.g., Kulmala et al., 2012). Throughout the 1-year measurement campaign (365 d), we identified 207 event days (56.7 %), 119 non-event days (32.6 %), 31 undefined days (8.5 %), and 8 no-data days (2.2 %) mainly due to power cuts at the station (Fig. 4). The annual-median NPF frequency at CAO-AMX calculated without accounting for days with no data amounts to 58 %, which belongs to the high end of the global NPF frequency distribution (Nieminen et al., 2018), with the highest frequency being measured in South Africa (86 %; Hirsikko et al., 2012). High frequency of NPF occurrence is also observed for Saudi Arabia (73 %; Hakala et al., 2019).




Figure 2. Monthly variation (at radiation $> 50 \text{ W m}^{-2}$) of particle number concentration of (a) cluster mode, (b) nucleation mode, (c) Aitken mode, and (d) accumulation mode presented by boxplots. The central red marks indicate the median, the small blue squares indicate the mean, and the bottom and top edges of the boxes indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers, and the outliers are plotted individually using the “+” symbols. Data presented have daily time resolution. Month designated with “*” symbols have less than 20 d of data. Note that SMPS measurements were not available in November and December.




Figure 3. The diurnal cycle (at radiation $> 50 \text{ W m}^{-2}$) of particle number concentration of cluster mode (a), nucleation mode (b), Aitken mode (c), and accumulation mode (d). The shaded areas with dashed black boundaries represent the 25th and 75th percentile limits, while the solid line represents the median and the squares indicate the mean. Notice the difference in the y scale between the top and bottom plots.




Figure 4. Classification of NPF events presented (a) annually and (b) monthly.

NPF took place throughout the year, but it had a clear seasonal pattern with a broad spring maximum, less pronounced autumn maximum, and slightly lower frequencies during other times of the year. The months with the highest NPF frequencies were March and April, while June and August had the lowest frequencies. This seasonal pattern of NPF frequency is very similar to that recorded at the Finokalia atmospheric observation station in Crete (Kalivitis et al., 2019), which is a nearby Eastern Mediterranean site having similar synoptic conditions. Nonetheless, monthly NPF frequency at Finokalia ranged between $\sim 17\%$ and 42% , which is substantially lower than the range reported here (33% – 86%). The higher NPF frequency at CAO-AMX could partially be due to the use of PSD data that starts from the ~ 1 nm size range, which facilitates NPF classification especially during days when the particle growth does not pass the 10 nm size or does not continue for several hours. We compared the NPF classification using SMPS data only and using full PSD for time periods when SMPS data were available, and we attained 30 % less event days classified. Another factor that could contribute to the higher NPF frequency is the surrounding forest nature which emits VOCs that oxidize in the atmosphere and contribute to particle growth (Riipinen et al., 2011).

We further separated the NPF event days into class I or class II events or into the so-called “bump” events (Manninen et al., 2010). The calendar of event classification is presented in Fig. S7, and examples of event types are given in Fig. S8. Class I events differ from class II events not by the strength of the event but rather by the ability to calculate the particle growth rate for such event, meaning that the growing mode diameter or concentration does not fluctuate strongly. Bump events are NPF events where a burst of nucleation mode particles is seen, but the particles do not usually grow past the ~ 10 nm size, and the duration of these events is typically short. The calculation of growth rates for these events is sometimes problematic because the growth happens very fast (in less than 15 min), and it cannot be captured by the time resolution of the measuring instrument. In the literature, these events have been called “bursting events” (Dall’Osto et

al., 2017), “hump events” (Vakkari et al., 2011; Yli-Juuti et al., 2009), “suppressed events” (Chen et al., 2017), “stationary NPF events” (Größ et al., 2018), or “weak NPF events” (Lee et al., 2020). The fraction of these events was highest during the month with the lowest NPF frequency (mainly during summer), which could imply that during these months a lesser amount of condensing vapors was present to grow the particles to bigger sizes or extend the event duration (Fig. 5).

The NPF events started almost always from the sub-3 nm range at CAO-AMX. The apparent growth reached a diameter of 20 nm on 25 % of event days (Fig. 6a); thus, it could have been difficult to identify those days if we have relied solely on SMPS measurements which suffer from high losses and low counting statistics in the sub-10 nm size range (Brilke et al., 2020; Kangasluoma et al., 2020; Wiedensohler et al., 2012). Additionally, it was difficult to distinguish the growing mode at sizes above 50 nm, mainly because of the pre-existing aerosols and fluctuating air masses. This implies that particles growing from NPF might have been able to grow to bigger sizes, but their identification from the PSD spectrum was not possible. The median event duration was ~ 5.4 h (Fig. 6b). The events typically started 2 to 4 h after sunrise and ended 7 to 11 h after sunrise (Fig. 6c), similar to what was observed by Dada et al. (2018).

Another feature of NPF events observed at CAO-AMX was the occurrence of two or three consecutive daytime nucleation events (Fig. 7). These multiple events occurred on $\sim 20\%$ of the recorded event days. Similar observations were reported in South Africa, and they were mainly attributed to changes in air masses, interruptions by clouds, and boundary layer dynamics as well as their relation to the amount of vapors present (Hirsikko et al., 2013). Salma and Németh (2019) have also showed that NPF events with broad or multiple onsets are abundant in the urban environment of Budapest, Hungary.

We also observed events with a decreasing mode diameter, sometimes referred to as shrinkage events. These events were mainly observed in the NAIS ion mode, while some of them were also observed in both ion and total particle spectra (Fig. 8). These types of events have been observed in multi-




Figure 5. Monthly percentage of the classes of event days.




Figure 6. Percentage histograms showing the frequency distribution of (a) NPF events growing to a certain diameter, (b) NPF event duration, and (c) the event start and end times from sunrise.

ple environments and are usually attributed to particle evaporation triggered by elevated temperatures, size-dependent dilution caused by wind, or boundary layer development mixing, or changes in air masses bringing small particles to the measurement site (Alonso-Blanco et al., 2017; Backman et al., 2012; Cusack et al., 2013; Hakala et al., 2019; Kivekäs et al., 2016; Salma et al., 2016b; Skrabalova et al., 2015; Tsagkogeorgas et al., 2017; Yao et al., 2010; Young et al., 2013; Zhang et al., 2016; Carnerero et al., 2018).

We spotted a few events with nighttime clustering, which could reflect a chemistry that does not depend on photo-oxidation (Fig. 9). These events occurred mainly during the cold months associated with high cluster mode concentration. Nighttime events have been observed in other Mediterranean studies as well (Carnerero et al., 2018; Kopanakis et al., 2013; Kalivitis et al., 2012). In a boreal forest, night-

time clustering events that do not usually grow past 5 nm have been attributed to the formation of large, highly oxygenated organic molecules (HOMs), mainly from monoterpene oxidation (Lehtipalo et al., 2011; Rose et al., 2018; Bianchi et al., 2019). In the French Landes forest, nocturnal NPF events with clear growth up to 100 nm were attributed to monoterpene oxidation under stratified atmospheric conditions (Kammer et al., 2018). Monoterpene concentrations reported at the Landes forest reached up to 25 ppb, whereas those measured in the boreal forest were below 2 ppb. Concurrent measurements of biogenic volatile organic compounds (BVOCs) were not available in this study, but the average concentration of monoterpenes during March 2015, which is a month with high biogenic activity, was reported to be 0.236 ± 0.294 ppb with a maximum up to 4.5 ppb (Debevec et al., 2018).




Figure 7. Examples of days with multiple nucleation events: (a) 18 March 2018, (b) 22 March 2018, (c) 8 April 2019, (d) 30 April 2018, (e) 9 September 2018, and (f) 30 September 2018.




Figure 8. Example of days showing a decreasing mode diameter: (a) 27 March 2018, (b) 1 April 2018, (c) 18 April 2018, (d) 2 May 2018, (e) 17 September 2018, and (f) 27 October 2018 in local time. The mode diameter is plotted as black circular markers.

Lastly, the EMME region is characterized by a high loading of dust which contributes to around 34 % ($\sim 10 \mu\text{g m}^{-3}$) of the annual PM_{10} levels (Pikridas et al., 2018). In this study, 50 d with high dust loading (Table S3) was identified based on ground measurements of mineral dust concentrations (Sect. S5). Among these dates, 37 NPF event days, 9

non-events, 2 undefined days, and 2 no-data days were identified. Figure 10 shows the temporal variation of $\text{PM}_{10-2.5}$, $\text{PM}_{2.5}$, and particle number size distribution measured during three of the dust episodes with ± 5 d window before and after the dust episode. NPF took place at high dust loadings, and there is no obvious threshold for the dust loading above




Figure 9. Examples of days with nighttime clustering and growth marked with white rectangles: (a) 7 February 2018, (b) 22 February 2018, (c) 21 March 2018, (d) 20 October 2018, (e) 4 December 2018, and (f) 18 December 2018. The black line is the solar radiation (W m^{-2}), which can be read from the right axis. Note the difference in the color scale used in this figure in comparison to Figs. 7 and 8.

which NPF does not occur. In addition, the formation rates (Fig. S9) and growth rates (Fig. S10) between NPF event days not affected by high dust loading and NPF event days affected by high dust loadings seem to be comparable. J_7 was slightly higher on days affected by high dust loading, but this could be related to the lower number of dust cases compared with the non-dust cases. High dust loadings can affect NPF in opposing ways. On the one hand, it can suppress photochemical processes by scavenging reactive gases and condensable vapors (De Reus et al., 2000; Ndour et al., 2009). On the other hand, it can provide particles that can act as a site for heterogeneous photochemistry promoting the formation of gaseous OH radicals, which initiate the conversion of SO_2 to H_2SO_4 (Dupart et al., 2012; Nie et al., 2014). However, a clear association between high dust loading and NPF was not found from the data set presented here.

3.3 NPF-specific parameters

In this section we analyze two parameters that describe the strength of NPF: particle formation rates (J) and particle apparent growth rates (GR).

Particle formation rates. The particle formation rates for 1.5, 3, and 7 nm particles ($J_{1.5}$, J_3 , and J_7 , respectively) were calculated when SMPS measurements were available (until 2 November), and they are presented in Fig. 11. $J_{1.5}$ was the highest during the spring: March had the highest median $J_{1.5}$, while April had more events with extreme $J_{1.5}$ values as expressed by the higher mean. In contrast, J_3 and J_7 did not exhibit a clear seasonality, but their values were in general




Figure 10. Temporal variations of aerosols during dust episodes with 5 d before and 5 d after the dust episode. (a) Time series of particle size distribution, $\text{PM}_{10-2.5}$, and $\text{PM}_{2.5}$ between 1 and 15 February 2018 (dust episode: 6 to 10 February). (b) Time series of particle size distribution, $\text{PM}_{10-2.5}$, and $\text{PM}_{2.5}$ between 15 March and 2 April 2018 (dust episode: 20 to 28 March). (c) Time series of particle size distribution, $\text{PM}_{10-2.5}$, and $\text{PM}_{2.5}$ between 26 April and 15 May 2018 (dust episode: 26 to 27 April and 1 to 7 May).

higher during the spring. The diurnal cycle for the formation rates was more pronounced during the class I events than during the class II or bump events, and the peak median hourly value was highest during class I events (Fig. 12). The median peak of $J_{1.5}$ and J_3 during the class I events and bump events occurred between 11:00 and 12:00 LT, whereas for J_7 the peak occurred between 12:00 and 14:00 LT. For the class II events, the corresponding peaks occurred about 1 h later. To place the formation rates in a global perspective, we compare J_3 from this study to other studies (Table 1), because it is the most commonly reported value in literature. The studies in Table 1 were selected on the basis of having 1 year or more of measurement data. J_3 values determined in this study were higher than those measured at semi-pristine rural areas (Värrio, Hyytiälä, and Tomsk), lower than those measured in a megacity (Beijing), and higher than those values reported at urban and rural sites affected by urban pollution (Budapest, Helsinki, Vavihill, and Po Valley).

Apparent growth rates. We report size-segregated growth rates between 1.5 and 3 nm ($GR_{1.5-3}$), between 3 and 7 nm (GR_{3-7}), and between 7 and 20 nm (GR_{7-20}) as recommended by Kulmala et al. (2012) for negatively charged ions, positively charged ions, and total particles (charged + neutral), respectively (Fig. 13). The growth rates of total particles were higher than those of the charged fraction, which is in agreement with earlier studies showing enhanced growth rates in the neutral channel at diameters below 15 nm (Gonser et al., 2014; Manninen et al., 2009; Rohan Jayaratne et al., 2016). This behavior has been explained by Gonser et al. (2014), who provided a conceptual model of the influence of cluster ion recombination and attachment at different stages of particle nucleation and growth. The seasonal behavior of the growth rates was also distinct. In the sub-3 nm range, the growth rates of the negative ions had similar median values across the year except during July, which had higher growth rates, whereas the positive ions had a notable increase in the growth rates in the summer months. The difference in the growth rates at these cluster sizes suggests that the ion-induced NPF processes are more important in the positive channel. In the 3–7 nm size range, there was no clear seasonal pattern except that June had the highest growth rates in the negative and positive mode, while the month exhibiting the highest growth rates in the total particle mode were February and June. In the 7–20 nm size range, the growth rates exhibited a clear seasonality in all channels with a peak in February and another broad peak during the summer months. The GR increased with an increasing particle size, which is a typical feature in the sub-20 nm size range, because condensational growth is more favorable as the particle size increases and the Kelvin effect decreases (Manninen et al., 2010). The median growth rates in the three size ranges (calculated from the daily means of the three channels) were 3.7, 9.2, and 11.7 nm h⁻¹. These GRs are higher than those reported for a rural boreal environment (1.9, 3.8, and 4.3 nm h⁻¹) (Yli-Juuti et al., 2011). In comparison to

other studies, the ion mode GR reported here is on the higher range of GRs measured at 12 European sites (Manninen et al., 2010 cf. Fig. S11). The high growth rates reported here could be associated with the high fraction of bump events. As discussed in Sect. 3.3, these events are characterized by a burst of particles within a short period of time, which would translate to higher growth rates.

3.4 The driving atmospheric parameters of the NPF events

To explain the occurrence of NPF at CAO-AMX, we investigated the effect of the following variables: CS, meteorological conditions (temperature, solar radiation, pressure, relative humidity, wind speed and wind direction), trace gas concentrations (NO_x, SO₂, CO, and O₃), air mass origin, and sulfuric acid concentrations.

The median of CS at CAO-AMX, for the periods that SMPS measurements were available, was $7.9 \times 10^{-3} \text{ s}^{-1}$ (25th–75th percentiles = 5.2×10^{-3} – 13.9×10^{-3}), while the mean was $10.7 \times 10^{-3} \text{ s}^{-1} \pm 8.2 \times 10^{-3} \text{ s}^{-1}$ (computed from daily median values). These values lie within the range of coastal (Kalivitis et al., 2019) and urban environments (Salma et al., 2016a; Jun et al., 2014). They are higher than the values reported for forests and semi-pristine environments (Dal Maso et al., 2002; Dada et al., 2017) and lower than the values reported for highly polluted cities (Wu et al., 2007). The average diurnal cycle of the size-segregated CS for the whole measurement period shows that particles above 50 nm were the main contributors to the CS, even though particles down to ~ 3 nm could also exhibit a high CS (Fig. S12). Thus, nucleating aerosols can largely contribute to the available aerosol surface area. The NPF frequency typically decreases with an increasing CS (Pikridas et al., 2012; Salma et al., 2016a; Dada et al., 2017; Dai et al., 2017; Hakala et al., 2019; Hussein et al., 2020). However, NPF has been observed in polluted environments at exceptionally high values of CS, indicating that inefficient cluster scavenging or enhanced cluster growth (or a combination of both) drives NPF regardless of the high load of pre-existing particles (Yao et al., 2018; Kulmala et al., 2017). In our study, we did not find a clear relation between CS and the monthly NPF occurrence, and NPF did not necessarily occur at low values of CS (Fig. 14). To further explore the effect of CS on NPF, we checked whether the NPF event days had lower CS before the onset of nucleation (period from midnight to morning) in comparison to non-event days, but we did not find any apparent association (Fig. S13). A possible explanation for why CS is not systematically lower on NPF event days could be similar to that observed at some mountain sites where the sources of NPF precursors and their sinks (i.e., CS) share the same origin; thus, CS is not necessarily a limiting factor (Sellegrì et al., 2019).

Next, we inspected the effect of meteorological variables (Fig. 15) on the occurrence of NPF. By considering the data




Figure 11. Monthly variation of particle formation rates during NPF events: (a) $J_{1.5}$, (b) J_3 and (c) J_7 . The central marks indicate the median, the small blue squares indicate the mean, and the bottom and top edges of the boxes indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers, and the outliers are plotted individually using the “+” symbols. The numbers above the boxplot represent the number of data points within each boxplot. Data presented have daily time resolution. Daily J values were calculated by taking the mean of hourly values within event duration times.




Figure 12. Diurnal variation of $J_{1.5}$ (top), J_3 (middle), and J_7 (bottom) during class I, class II, bump events, and non-events. Shaded areas represent the 25th and 75th percentile bounds, while the solid lines represent the median.

Table 1. Comparison of formation rates from long-term measurements at various sites including this study.

Site	Type	Period	J_3			Reference
			Mean \pm SD	Median	Min–max	
CAO-AMX, Cyprus	Rural	2018–2019	6.42 \pm 8.47	4.19	0.04–82.5	This study
Vavihill, Sweden	Rural	2001–2004	4.3	1.89		Kristensson et al. (2008)
Hyytiälä, Finland	Rural	1996–2004	0.8	0.6	0.06–5.0	Dal Maso et al. (2007)
Värriö, Finland	Rural	1996–2004	0.2	0.1		Dal Maso et al. (2007)
Tomsk, Russia	Rural	2005–2006	0.4	0.4	0.04–1.1	Dal Maso et al. (2008)
Beijing, China	Urban clean	2004–2005	22.3 \pm 15.1	18.7	4.4–81.4	Wu et al. (2007)
	Urban polluted		16.2 \pm 12	9.9	3.3–51.7	
St. Louis, USA	Urban	2001–2003	17 \pm 20	9		Qian et al. (2007)
Po Valley, Italy	Rural w. urban influence	2002–2005	5.89	3.31	0.24–36.89	Hamed et al. (2007)
Budapest, Hungary	Urban	2008–2009	4.2 \pm 2.5 ^a	4.2 ^a	1.65–12.5 ^a	Salma et al. (2011)
Helsinki, Finland	Urban	1997–2006	2 ^b	1.09 ^b	0.63–2.87 ^b	Hussein et al. (2008)
Zeppelin, Norway	Arctic	2016–2018			0.02–1.62	Lee et al. (2020)

^a data represent 6–25 nm range. ^b calculated from monthly values.

from all the months together, NPF events took place over a wide range of meteorological conditions. Higher temperatures seemed to be favorable for intra-monthly NPF occurrence; however, the higher temperatures from June to September did not coincide with higher NPF frequencies (Fig. 15a). The effect of temperature on NPF has been studied extensively in chamber experiments, with a general consensus that lower temperatures favor nucleation at the kinetic regime and thus enhance NPF in inorganic systems like the sulfuric acid–ammonia system (Lee et al., 2019). However, in organic systems where highly oxygenated organic compounds (HOMs) are the main NPF species, temperature plays a double role. On the one hand, the Gibbs free-energy barrier is reduced at lower temperatures, favoring the condensation of less oxidized vapors that would not condense at higher temperatures. On the other hand, lower temperatures lead to decreased auto-oxidation reaction rates and reduced yields of HOMs. Recent studies have shown that the former effect compensates for the latter effect, having an overall increase in nucleation and growth rates at lower temperatures (Stolzenburg et al., 2018; Simon et al., 2020; Ye et al., 2019). While these mechanisms are clear in chamber studies, the situation becomes more complicated in the atmosphere because of the complexity of the atmosphere–biosphere system having simultaneous temperature-dependent processes that can enhance or suppress NPF, making current atmospheric observations inconsistent on the role of temperature on NPF (Kerminen et al., 2018). Solar radiation is regarded as one of the most important factors affecting NPF (Jokinen et al., 2017). Its intensity is relatively high in Cyprus all year round.

Intra-monthly, NPF events occurred at higher global radiation during the winter and autumn months, whereas in spring (except April) and summer months radiation did not seem to be a limiting factor for NPF (Fig. 15b). Inter-monthly, the month with the highest solar radiation did not coincide with the highest occurrence of NPF. Regarding ambient relative humidity (RH), NPF events tend to occur at lower RH in both clean and polluted environments (Kerminen et al., 2018). However, high RH values do not necessarily suppress NPF (Salma and Németh, 2019), which agrees with our observations (Fig. 15c). In terms of the surface air pressure, intra-monthly NPF was on average observed on days with higher pressures, and the inter-monthly NPF occurrence was the lowest during the month with the lowest surface pressure (Fig. 15d). With respect to wind speed, high wind speeds did not seem to prevent NPF, but event days occurred mostly under low wind speeds (Fig. S14). In terms of wind direction, NPF occurred mainly when the wind was blowing from the west to east sector but with a frequency of occurrence which is higher in north to easterly winds (Figs. S14, S15, and S16). The north-to-easterly direction is the direction where the main agglomerations and livestock farming lands are situated. These local sources could be enhancing the occurrence of NPF, but no direct relation was found between the north-to-easterly wind direction and specific event types (Fig. S16). However, NPF class I events did not occur when the wind was originating from the southeast to southwest sector.

The EMME region is among the regions with the highest background of trace gases and aerosol concentrations in the Northern Hemisphere (Lelieveld et al., 2002). Here, we




Figure 13. Monthly variation of growth rates during NPF events in three size ranges: (a) < 3 nm, (b) 3–7 nm, and (c) 7–20 nm. The central marks indicate the median, and the bottom and top edges of the boxes indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points not considered outliers, and the outliers are plotted individually using the “+” symbols. The numbers above the boxplot represent the number of data points within each boxplot. Black boxes represent the total particles (neutral+charges), blue boxes represent negative ions, and red boxes represent positive ions.




Figure 14. Monthly variation of condensation sink (s^{-1}) during event (blue) and non-event (green) days using data corresponding to when global radiation is greater than 50 W m^{-2} . The bottom and top edges of the boxplots indicate the 25th and 75th percentiles, respectively. The central mark indicates the median. The whiskers extend to the most extreme data points not considered outliers, and the outliers are plotted individually using the “+” symbols. Data presented have hourly time resolution. The shaded grey bars represent the monthly NPF percent occurrence.

investigate the relation between trace gases and the occurrence of NPF (Fig. 16). The intra-monthly SO_2 concentration was, on average, higher during NPF event days in comparison to non-event days during most of the months. The inter-monthly SO_2 concentrations during the highest NPF occurrence (March–May) were similar to the months with the lowest NPF occurrence (June–August). This indicates that SO_2 and thus sulfuric acid (as will be shown subsequently) cannot alone explain the seasonal pattern of NPF. The ozone (O_3) concentration is particularly high in Cyprus and is mainly influenced by regional and transported ozone, while local precursor emissions play a minor role in ozone formation (Kleanthous et al., 2014). Intra-monthly, the O_3 concentration was sometimes lower, similar, or higher during NPF event days compared with non-event days, with no clear seasonality. One notable remark is that in April NPF events took place at much higher O_3 concentrations than what was observed on non-event days. This could imply that higher oxidative capacity was driving NPF during April. April also had the most notable differences in global radiation and RH between NPF event and non-event days. Analogous to SO_2 , O_3 cannot explain the seasonal pattern of NPF.




Figure 15. Monthly variation of meteorological parameters during event (blue) and non-event (green) days: (a) temperature, (b) global radiation, (c) relative humidity, and (d) pressure, using data corresponding to global radiation greater than 50 W m^{-2} . The bottom and top edges of the boxplots indicate the 25th and 75th percentiles, respectively. The central mark indicates the median. The whiskers extend to the extreme data points not considered outliers. Data presented have hourly time resolution. The shaded grey bars represent the monthly NPF percent occurrence.




Figure 16. Monthly variation of trace gases during event (blue) and non-event (green) days: (a) sulfur dioxide, (b) ozone, (c) carbon monoxide, and (d) nitrogen oxides using data corresponding days when global radiation was greater than 50 W m^{-2} . The shaded grey bars represent the monthly NPF percent occurrence. The bottom and top edges of the boxplots indicate the 25th and 75th percentiles, respectively. The central mark indicates the median. The whiskers extend to the most extreme data points not considered outliers. Data presented have hourly time resolution. The shaded grey bars represent the monthly NPF percent occurrence.

CO levels are generally high over the Mediterranean (in comparison to the Pacific), with emission sources being typically from western and eastern Europe, having lower contributions from the regions surrounding the Mediterranean (Lelieveld et al., 2002). The CO–NPF relationship at CAO-AMX did not have a distinct character; however, CO concentrations were slightly lower during the summer months. Regarding NO_x , NPF event days had on average higher NO_x concentrations within the boundaries of each month, except in April. More notably, NO_x had lower concentrations during the months with lower NPF frequencies, which might indicate the role of associated anthropogenic organic vapors in triggering NPF at our site.

We examined the effect of air mass origin arriving at CAO-AMX at 08:00 LT during event and non-event days from

seven source regions: local, N Africa, marine, Europe, Asia, NW Asia, and SW Asia (Fig. 17). The last two source regions (NW Asia, and SW Asia) represent the geographic location corresponding to Cyprus. An obvious feature that pops out is that the months with the highest NPF frequency had air masses originating from southwest Asia (the Middle East), whereas during the month with the lowest NPF frequency air masses did not originate from that direction. This pattern might suggest that chemical compounds important for nucleation and subsequent growth are transported to CAO-AMX from the Middle East. Between the end of spring and late September, which are the months with the lowest NPF frequency, the circulation over the eastern Mediterranean is characterized by persistent northerly winds called the Etesians (Tyrllis and Lelieveld, 2013). The NPF events during




Figure 17. (a) The source regions of air masses reaching CAO-AMX used for the air mass sector analysis. The map was plotted using the Climate Data Toolbox for MATLAB (Greene et al., 2019). The International Hydrographic Organization (IHO) World Sea Areas v3 data were used to retrieve the boundaries of the Mediterranean Sea (Flanders Marine Institute, 2019). Note that marine areas other than the Mediterranean Sea were considered part of the continental sectors and that the NW Asia and SW Asia sectors correspond to Cyprus. (b) Monthly variation of air mass origin arriving at CAO-AMX at 08:00 LT during event (E) and non-event days (NE). There are no air masses originating from the Asia sector, because they are obscured by terrain height.

this period, as shown in Sect. 3.2, were weak or generally did not lead to particle growth into large sizes in comparison to the rest of the year. The Etesian circulation flow is caused by a sharp surface pressure difference between the westerly Azorean high-pressure regime and the Asian monsoon low-pressure regime. While the Etesians block the northward transport of desert dust, they trigger high sea levels, prevent rain over the region, and enhance marine inversions (Ulbrich et al., 2012). They favor the transport of air pollutants from central or eastern Europe and west Turkey and, together with enhanced photochemical conditions and low precipitation, contribute to high O_3 (Solomou et al., 2018) and particulate matter (PM) levels (Pikridas et al., 2018). The increase in PM levels during these months could be a limiting factor for NPF. Since we did not have particle size distribution measurements above 700 nm, bigger particles could be additionally contributing to the CS. However, from a mass concentration point of view, $PM_{2.5}$ and PM_{10} did not show a pattern that would support this hypothesis (Figs. S17 and S18). Additionally, while the southwest Asia sector might be important for NPF, it did not exhibit a clear pattern during the month with the highest NPF frequency. In fact, in April most of the air masses originated from northwest Asia. This sector

appears to be also important for NPF during June, July, and November, whereas the other sectors did not exhibit any notable pattern. Air masses did not originate from the Asia sector, because they were obscured by the terrain height. Pure marine air masses were not as frequent as other air masses. However, owing to the location of Cyprus, air masses from other continental source origins are expected to have been influenced by marine conditions as they travel to our measurement site. Thus we cannot exclude the potential marine effect on the occurrence of NPF.

Last, we investigated the role of sulfuric acid. While sulfuric acid (H_2SO_4) is considered one of the main nucleating species in the atmosphere, it is well known that H_2SO_4 binary nucleation with water requires high H_2SO_4 vapor concentrations that are not relevant within the lower parts of the troposphere (Wyslouzil et al., 1991). Additional species are required to stabilize H_2SO_4 clusters, such as ammonia, amines, or ions, while some other compounds can nucleate on their own in atmospherically relevant conditions, including iodine oxides and highly oxygenated organic compounds (HOMs) from biogenic precursors (Lee et al., 2019, and references therein). In this study, the hourly H_2SO_4 proxy concentrations ranged between 3×10^5 and 1×10^7 cm^{-3} ,

which are typical values for H_2SO_4 in the troposphere. The relationship between particle formation rates ($J_{1.5}$) and H_2SO_4 proxy concentration varied across the month of the year (Fig. 18). Lower concentrations of H_2SO_4 were required during winter and spring to achieve the same formation rates as in the other seasons. A possible explanation for this behavior is that, in the first case, stabilizing compounds are abundant in the atmosphere and thus less H_2SO_4 is required for the formation of particles. A similar hypothesis was tested by Pikridas et al. (2012) by using the accumulation mode particle acidity as an indirect measure of the availability or lack of ammonia (or any other basic species in the gas phase). The authors concluded that excess base is not available during the summer to participate in the nucleation process. In our case, however, the formation rate versus H_2SO_4 relationship is closer to those derived for the H_2SO_4 –DMA– NH_3 – H_2O system than those for the H_2SO_4 – NH_3 – H_2O system. In fact, the ternary nucleation of H_2SO_4 – NH_3 – H_2O is unlikely to be important at ground level either because of too low concentrations or too high temperatures (Kürten et al., 2018). This suggests that, in our case, the missing stabilizing base is probably not ammonia, although the role of ammonia cannot be ruled out. The distinct air mass origin during the summer could explain the decrease in the concentrations of the stabilizing base. Otherwise, the high temperature during the summer could be the factor that disfavors the occurrence of NPF. Most certainly, NPF at CAO-AMX seems to be influenced by several factors and chemical constituents. This has been also indicated by Debevec et al. (2018), who observed four types of nucleation events, within 1 month of measurements, having (1) predominant anthropogenic influence, (2) predominant biogenic influence, (3) mixed anthropogenic–biogenic influence, and (4) a marine influence with low concentrations of anthropogenic and biogenic tracers. Therefore, to reveal the main mechanisms of NPF, long-term measurements of nucleating clusters and organic precursors using state-of-the-art online mass spectrometry techniques are essential.

3.5 Regression and classification analysis

To further understand the occurrence of NPF events, we present, in Sect. S12 of the Supplement, two types of analysis: the first is a linear regression analysis of formation rate of 1.5 nm particles ($J_{1.5}$) and the second is a decision tree classification model to indicate whether each day is an NPF event day or a non-event day. Both analyses have shown that NO_2 , H_2SO_4 , and wind direction (mainly from N to E direction) are the most important parameters that are associated with NPF occurrence (Figs. S20–S23). While the role of H_2SO_4 in NPF is well known in literature, the role of NO_2 is not that clear. NO_x has been shown to play contrasting roles in NPF depending on the associated pool of gas molecules. On the one hand, when oxidized to nitric acid, it can enhance NPF in the presence of ammonia vapors (Wang et al., 2020). On the other hand, it can suppress NPF by reducing autox-




Figure 18. $J_{1.5}$ versus sulfuric acid proxy concentrations color-coded by the month of the year. Data presented have hourly time resolution.

idation and low-volatility HOM dimer formation (Wildt et al., 2014; Zhao et al., 2018). Nevertheless, Yan et al. (2020) have shown that this effect is weak when NH_3 and H_2SO_4 are additionally present and that NO is more effective than NO_2 in changing the HOM composition and volatility. Xie et al. (2015) have revealed that NO_2 can play an important role not only in surface catalytic reactions of SO_2 but also in dust-induced photochemical heterogeneous reactions of NO_2 , which produces additional sources of OH radicals and promote new particle formation and growth. However, while NPF seems to occur more frequently at higher NO_2 concentrations in our study, we cannot conclude if it plays a role in NPF or if it is a proxy of some other pollutant, especially since NO_2 concentrations were mostly lower than 4 ppb. What is evident, however, is that H_2SO_4 does not nucleate on its own at the concentrations reported in this study; thus, an unknown stabilizer and possibly other compounds participating in NPF are missing in this analysis (as explained in the regression analysis). We hypothesize that these unknown compounds (e.g., NH_3 , amine, or HOMs) are associated with the north-to-east wind directions and higher NO_2 concentrations.

4 Conclusion

Recent studies have pointed out that NPF is important in the EMME region (Brilke et al., 2020; Debevec et al., 2018; Hakala et al., 2019; Hussein et al., 2020; Kalivitis et al., 2019; Kalkavouras et al., 2019). Brilke et al. (2020) studied NPF in a coastal site in Cyprus with strong local pollution during 2017, while Debevec et al. (2018) characterized NPF at the same site of this study during 2015. While both studies were limited to 1 month of observations, we disclosed here the first long-term (1 year) characterization of NPF at a back-

ground site in Cyprus. We presented the general and seasonal characteristics of PSD and NPF, then we explored the factors that affect NPF.

Our analysis of NPF intra-monthly variability showed that, on average, NPF events occurred at higher temperatures, lower RH, and higher global radiation, except during the months of August, September, and December. On the contrary, lower pressure conditions, higher wind speeds, and local south to west wind directions seemed to be more favorable for non-events. The frequency of NPF was higher than that reported at a similar Eastern Mediterranean island site using a slightly limited measurement setup than the one applied here. This demonstrates the importance of comprehensive measurements using instruments that can measure down to cluster sizes. NPF occurred all year round, with higher frequencies during the spring and autumn and a minimum frequency during the summer. The particles did not grow significantly after nucleation during the months with the lowest NPF frequencies. These months were also characterized by lower NO_x concentrations, which is an indication of lower anthropogenic influence, and distinct air mass origin profiles from the rest of the year. Condensation sink, calculated based on a PSD up to 700 nm, had no clear relationship with NPF, but it was slightly higher during some summer months. Additionally, sulfuric acid was not the limiting factor for NPF seasonality as its estimated concentration was mostly high during the summer, up to 1×10^7 molecules cm^{-3} . The relationship between particle formation rates and sulfuric acid proxy exhibited different slopes between the months with the highest and lowest NPF frequency, suggesting that nucleation might have proceeded with varying temperatures or at different concentrations of stabilizing compounds and other aerosol precursors not measured in this study.

The analysis presented in this study is a step forward towards understanding the mechanisms of NPF in the EMME region. Future studies require long-term measurements of vapors that participate in NPF and subsequent growth. These could include, for example, ultralow volatility organic compounds (ULVOCs), extremely low-volatility organic compounds (ELVOCs), low-volatility organic compounds (LVOCs), ammonia, amines, and iodine species. Further, to understand the ubiquity of the effect of large particles, which could inhibit NPF during certain episodes but enhance NPF during episodes with high mineral dust loadings, extended PSD measurements up to coarse particles, preferably coupled with chemical speciation, are important. On a larger scale, long-term measurements of CCN particles are necessary to quantify the contribution of NPF to the CCN budget. These measurements would preferably take place not only in Cyprus but also in different locations in the Middle East and northern Africa.

Data availability. The data used in this study are available at <https://doi.org/10.5281/zenodo.4701303> (Baalbaki et al., 2021).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-21-9223-2021-supplement>.

Author contributions. The study was conceived by MK and JS. RB, TJ, TL, KN, and MP prepared and installed the instruments. RB, MP, KN, AM, EB, AC, and FU performed the regular maintenance for the instruments. RB performed the data analysis and wrote the article. LD provided support for data analysis. LA provided support for instrument troubleshooting and nCNC inversion. MP and EB performed the sector analysis. SB performed the κ measurements and provided support for hygroscopicity calculations. MP and JS provided the SMPS data. CS provided the meteorological and trace gas data. RB, MP, TJ, LD, SB, KL, JK, GB, TP, VMK, and MK participated in the scientific discussion and reviewed the article.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The authors thank Hanna Manninen, Kaspar Dällenbach, Jenni Kontkanen, Runlong Cai, and Dominik Stolzenburg for fruitful scientific discussions. Frans Korhonen, Pekka Rantala, Pasi Aalto, Erki Siivola, Sander Mirme, Joonas Vanhanen, and Aki Halonen are kindly acknowledged for their continuous and indispensable support.

Financial support. This publication has been produced within the framework of the EMME-CARE project which received funding from the European Union's Horizon 2020 Research and Innovation Programme, under grant agreement no. 856612 and from the Cyprus Government. This work has received additional funding from the European Research Council (ERC) under the European Union's Horizon 2020 Research And Innovation Programme (ERC, project no. 742206 "ATM-GTP"). The sole responsibility of this publication lies with the author. The European Union is not responsible for any use that may be made of the information contained therein. Additional support was received from the Academy of Finland (grant agreement nos. 307331, 337549, 302958, and 316114), the European Regional Development Fund, and the Republic of Cyprus through the Research and Innovation Foundation (project: INTEGRATED/0916/0016). JK acknowledges support from the Academy of Finland (project 1325656) and the University of Helsinki 3-year grant 75284132. TJ acknowledges support from the Academy of Finland (project 334514).

Open-access funding was provided by the Helsinki University Library.

Review statement. This paper was edited by Fangqun Yu and reviewed by two anonymous referees.

References

- Alonso-Blanco, E., Gómez-Moreno, F. J., Núñez, L., Pujadas, M., Cusack, M., and Artíñano, B.: Aerosol particle shrinkage event phenomenology in a South European suburban area during 2009–2015, *Atmos. Environ.*, 160, 154–164, <https://doi.org/10.1016/j.atmosenv.2017.04.013>, 2017.
- Baalbaki, R., Pikridas, M., Jokinen, T., Laurila, T., Dada, L., Bezan-takos, S., Ahonen, L., Neitola, K., Maisser, A., Bimenyimana, E., Christodoulou, A., Unga, F., Savvides, C., Lehtipalo, K., Kangasluoma, J., Biskos, G., Petäjä, T., Kerminen, V.-M., Sciare, J., and Kulmala, M.: Towards understanding the characteristics of new particle formation in the Eastern Mediterranean, Zenodo, <https://doi.org/10.5281/zenodo.4701303>, 2021.
- Baccarini, A., Karlsson, L., Dommen, J., Duplessis, P., Vüllers, J., Brooks, I. M., Saiz-Lopez, A., Salter, M., Tjernström, M., Baltensperger, U., Zieger, P., and Schmale, J.: Frequent new particle formation over the high Arctic pack ice by enhanced iodine emissions, *Nat. Commun.*, 11, 4924, <https://doi.org/10.1038/s41467-020-18551-0>, 2020.
- Backman, J., Rizzo, L. V., Hakala, J., Nieminen, T., Manninen, H. E., Morais, F., Aalto, P. P., Siivola, E., Carbone, S., Hillamo, R., Artaxo, P., Virkkula, A., Petäjä, T., and Kulmala, M.: On the diurnal cycle of urban aerosols, black carbon and the occurrence of new particle formation events in spring-time São Paulo, Brazil, *Atmos. Chem. Phys.*, 12, 11733–11751, <https://doi.org/10.5194/acp-12-11733-2012>, 2012.
- Berland, K., Rose, C., Pey, J., Culot, A., Freney, E., Kalivitis, N., Kouvarakis, G., Cerro, J. C., Mallet, M., Sartelet, K., Beckmann, M., Bourriane, T., Roberts, G., Marchand, N., Mihalopoulos, N., and Sellegri, K.: Spatial extent of new particle formation events over the Mediterranean Basin from multiple ground-based and airborne measurements, *Atmos. Chem. Phys.*, 17, 9567–9583, <https://doi.org/10.5194/acp-17-9567-2017>, 2017.
- Bianchi, F., Tröstl, J., Junninen, H., Frege, C., Henne, S., Hoyle, C. R., Molteni, U., Herrmann, E., Adamov, A., Bukowiecki, N., Chen, X., Duplissy, J., Gysel, M., Hutterli, M., Kangasluoma, J., Kontkanen, J., Kürten, A., Manninen, H. E., Münch, S., Peräkylä, O., Petäjä, T., Rondo, L., Williamson, C., Weingartner, E., Curtius, J., Worsnop, D. R., Kulmala, M., Dommen, J., and Baltensperger, U.: New particle formation in the free troposphere: A question of chemistry and timing, *Science*, 352, 1109–1112, <https://doi.org/10.1126/science.aad5456>, 2016.
- Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crouse, J. D., Wennberg, P. O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A., Donahue, N., Kjaergaard, H. G., and Ehn, M.: Highly Oxygenated Organic Molecules (HOM) from Gas-Phase Autoxidation Involving Peroxy Radicals: A Key Contributor to Atmospheric Aerosol, *Chem. Rev.*, 119, 3472–3509, <https://doi.org/10.1021/acs.chemrev.8b00395>, 2019.
- Brilke, S., Föller, N., Müller, T., Kandler, K., Gong, X., Peischl, J., Weinzierl, B., and Winkler, P. M.: New particle formation and sub-10 nm size distribution measurements during the A-LIFE field experiment in Paphos, Cyprus, *Atmos. Chem. Phys.*, 20, 5645–5656, <https://doi.org/10.5194/acp-20-5645-2020>, 2020.
- Brines, M., Dall’Osto, M., Beddows, D. C. S., Harrison, R. M., Gómez-Moreno, F., Núñez, L., Artíñano, B., Costabile, F., Gobbi, G. P., Salimi, F., Morawska, L., Sioutas, C., and Querol, X.: Traffic and nucleation events as main sources of ultrafine particles in high-insolation developed world cities, *Atmos. Chem. Phys.*, 15, 5929–5945, <https://doi.org/10.5194/acp-15-5929-2015>, 2015.
- Brockmann, J. E.: Aerosol Transport in Sampling Lines and Inlets, in: *Aerosol Measurement: Principles, Techniques and Applications*, Third edition, edited by: Kulkarni, P., Baron, P. A., and Willeke, K., John Wiley & Sons, Hoboken, New Jersey, 69–105, 2011.
- Cai, R., Yang, D., Ahonen, L. R., Shi, L., Korhonen, F., Ma, Y., Hao, J., Petäjä, T., Zheng, J., Kangasluoma, J., and Jiang, J.: Data inversion methods to determine sub-3 nm aerosol size distributions using the particle size magnifier, *Atmos. Meas. Tech.*, 11, 4477–4491, <https://doi.org/10.5194/amt-11-4477-2018>, 2018.
- Carnerero, C., Pérez, N., Reche, C., Ealo, M., Titos, G., Lee, H.-K., Eun, H.-R., Park, Y.-H., Dada, L., Paasonen, P., Kerminen, V.-M., Mantilla, E., Escudero, M., Gómez-Moreno, F. J., Alonso-Blanco, E., Coz, E., Saiz-Lopez, A., Temime-Roussel, B., Marchand, N., Beddows, D. C. S., Harrison, R. M., Petäjä, T., Kulmala, M., Ahn, K.-H., Alastuey, A., and Querol, X.: Vertical and horizontal distribution of regional new particle formation events in Madrid, *Atmos. Chem. Phys.*, 18, 16601–16618, <https://doi.org/10.5194/acp-18-16601-2018>, 2018.
- Casquero-Vera, J. A., Lyamani, H., Dada, L., Hakala, S., Paasonen, P., Román, R., Fraile, R., Petäjä, T., Olmo-Reyes, F. J., and Alados-Arboledas, L.: New particle formation at urban and high-altitude remote sites in the south-eastern Iberian Peninsula, *Atmos. Chem. Phys.*, 20, 14253–14271, <https://doi.org/10.5194/acp-20-14253-2020>, 2020.
- Chan, T., Cai, R., Ahonen, L. R., Liu, Y., Zhou, Y., Vanhanen, J., Dada, L., Chao, Y., Liu, Y., Wang, L., Kulmala, M., and Kangasluoma, J.: Assessment of particle size magnifier inversion methods to obtain the particle size distribution from atmospheric measurements, *Atmos. Meas. Tech.*, 13, 4885–4898, <https://doi.org/10.5194/amt-13-4885-2020>, 2020.
- Chen, X., Virkkula, A., Kerminen, V.-M., Manninen, H. E., Busetto, M., Lanconelli, C., Lupi, A., Vitale, V., Del Guasta, M., Grigioni, P., Väänänen, R., Duplissy, E.-M., Petäjä, T., and Kulmala, M.: Features in air ions measured by an air ion spectrometer (AIS) at Dome C, *Atmos. Chem. Phys.*, 17, 13783–13800, <https://doi.org/10.5194/acp-17-13783-2017>, 2017.
- Chu, B., Kerminen, V.-M., Bianchi, F., Yan, C., Petäjä, T., and Kulmala, M.: Atmospheric new particle formation in China, *Atmos. Chem. Phys.*, 19, 115–138, <https://doi.org/10.5194/acp-19-115-2019>, 2019.
- Cusack, M., Alastuey, A., and Querol, X.: Case studies of new particle formation and evaporation processes in the western Mediterranean regional background, *Atmos. Environ.*, 81, 651–659, <https://doi.org/10.1016/j.atmosenv.2013.09.025>, 2013.
- Dada, L., Paasonen, P., Nieminen, T., Buenrostro Mazon, S., Kontkanen, J., Peräkylä, O., Lehtipalo, K., Hussein, T., Petäjä, T., Kerminen, V.-M., Bäck, J., and Kulmala, M.: Long-term analysis of clear-sky new particle formation events and non-events in Hyytiälä, *Atmos. Chem. Phys.*, 17, 6227–6241, <https://doi.org/10.5194/acp-17-6227-2017>, 2017.
- Dada, L., Chellapermal, R., Buenrostro Mazon, S., Paasonen, P., Lampilahti, J., Manninen, H. E., Junninen, H., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Refined classification and characterization of atmospheric new-particle formation

- events using air ions, *Atmos. Chem. Phys.*, 18, 17883–17893, <https://doi.org/10.5194/acp-18-17883-2018>, 2018.
- Dada, L., Ylivinkka, I., Baalbaki, R., Li, C., Guo, Y., Yan, C., Yao, L., Sarnela, N., Jokinen, T., Daellenbach, K. R., Yin, R., Deng, C., Chu, B., Nieminen, T., Wang, Y., Lin, Z., Thakur, R. C., Kontkanen, J., Stolzenburg, D., Sipilä, M., Hussein, T., Paasonen, P., Bianchi, F., Salma, I., Weidinger, T., Pikridas, M., Sciare, J., Jiang, J., Liu, Y., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Sources and sinks driving sulfuric acid concentrations in contrasting environments: implications on proxy calculations, *Atmos. Chem. Phys.*, 20, 11747–11766, <https://doi.org/10.5194/acp-20-11747-2020>, 2020.
- Dai, L., Wang, H., Zhou, L., An, J., Tang, L., Lu, C., Yan, W., Liu, R., Kong, S., Chen, M., Lee, S., and Yu, H.: Regional and local new particle formation events observed in the Yangtze River Delta region, China, *J. Geophys. Res.-Atmos.*, 122, 2389–2402, <https://doi.org/10.1002/2016jd026030>, 2017.
- Dal Maso, M., Kulmala, M., Lehtinen, K. E. J., Mäkelä, J. M., Aalto, P., and O'Dowd, C. D.: Condensation and coagulation sinks and formation of nucleation mode particles in coastal and boreal forest boundary layers, *J. Geophys. Res.*, 107, PAR 2-1–PAR 2-10, <https://doi.org/10.1029/2001jd001053>, 2002.
- Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P., and Lehtinen, K. E.: Formation and growth of fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II, Hyytiälä, Finland, *Boreal Environ. Res.*, 10, 323–336, 2005.
- Dal Maso, M., Sogacheva, L., Aalto, P. P., Riipinen, I., Kompula, M., Tunved, P., Korhonen, L., Suur-Uski, V., Hirsikko, A., Kurtén, T., Kerminen, V.-M., Lihavainen, H., Viisanen, Y., Hansson, H.-C., and Kulmala, M.: Aerosol size distribution measurements at four Nordic field stations: identification, analysis and trajectory analysis of new particle formation bursts, *Tellus B*, 59, 350–361, <https://doi.org/10.1111/j.1600-0889.2007.00267.x>, 2007.
- Dal Maso, M., Sogacheva, L., Anisimov, M. P., Arshinov, M., Baklanov, A., Belan, B., Khodzher, T. V., Obolkin, V. A., Staroverova, A., and Vlasov, A.: Aerosol particle formation events at two Siberian stations inside the boreal forest, *Boreal Environ. Res.*, 13, 81–92, 2008.
- Dall'Osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Ström, J., Hansson, H. C., Yoon, Y. J., Park, K.-T., Becagli, S., Udristi, R., Onasch, T., O'Dowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to atmospheric new particle formation, *Sci. Rep.-UK*, 7, 3318, <https://doi.org/10.1038/s41598-017-03328-1>, 2017.
- Dall'Osto, M., Beddows, D. C. S., Asmi, A., Poulain, L., Hao, L., Freney, E., Allan, J. D., Canagaratna, M., Crippa, M., Bianchi, F., de Leeuw, G., Eriksson, A., Swietlicki, E., Hansson, H. C., Henzing, J. S., Granier, C., Zemanekova, K., Laj, P., Onasch, T., Prevot, A., Putaud, J. P., Sellegri, K., Vidal, M., Virtanen, A., Simo, R., Worsnop, D., O'Dowd, C., Kulmala, M., and Harrison, R. M.: Novel insights on new particle formation derived from a pan-european observing system, *Scientific Reports*, 8, 1482, <https://doi.org/10.1038/s41598-017-17343-9>, 2018.
- Davies, L., and Gather, U.: The Identification of Multiple Outliers, *J. Am. Stat. Assoc.*, 88, 782–792, <https://doi.org/10.1080/01621459.1993.10476339>, 1993.
- De Reus, M., Dentener, F., Thomas, A., Borrmann, S., Ström, J., and Lelieveld, J.: Airborne observations of dust aerosol over the North Atlantic Ocean during ACE 2: Indications for heterogeneous ozone destruction, *J. Geophys. Res.*, 105, 15263–15275, <https://doi.org/10.1029/2000JD900164>, 2000.
- Debevec, C., Sauvage, S., Gros, V., Sellegri, K., Sciare, J., Pikridas, M., Stavroulas, I., Leonardis, T., Gaudion, V., Depelchin, L., Fronval, I., Sarda-Esteve, R., Baisnée, D., Bonsang, B., Savvides, C., Vrekoussis, M., and Locoge, N.: Driving parameters of biogenic volatile organic compounds and consequences on new particle formation observed at an eastern Mediterranean background site, *Atmos. Chem. Phys.*, 18, 14297–14325, <https://doi.org/10.5194/acp-18-14297-2018>, 2018.
- Drinovec, L., Sciare, J., Stavroulas, I., Bezantakos, S., Pikridas, M., Unga, F., Savvides, C., Višić, B., Remškar, M., and Močnik, G.: A new optical-based technique for real-time measurements of mineral dust concentration in PM10 using a virtual impactor, *Atmos. Meas. Tech.*, 13, 3799–3813, <https://doi.org/10.5194/amt-13-3799-2020>, 2020.
- Dupart, Y., King, S. M., Nekat, B., Nowak, A., Wiedensohler, A., Herrmann, H., David, G., Thomas, B., Miffre, A., Rairoux, P., D'Anna, B., and George, C.: Mineral dust photochemistry induces nucleation events in the presence of SO₂, *P. Natl. Acad. Sci. USA*, 109, 20842–20847, <https://doi.org/10.1073/pnas.1212297109>, 2012.
- Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, <https://doi.org/10.1038/nature13032>, 2014.
- Flanders Marine Institute: Maritime Boundaries Geodatabase: Maritime Boundaries and Exclusive Economic Zones (200NM), version 11, <https://doi.org/10.14284/386>, 2019.
- Fu, Y., Xue, M., Cai, R., Kangasluoma, J., and Jiang, J.: Theoretical and experimental analysis of the core sampling method: Reducing diffusional losses in aerosol sampling line, *Aerosol Sci. Tech.*, 53, 793–801, <https://doi.org/10.1080/02786826.2019.1608354>, 2019.
- Fuchs, N. A.: The mechanics of aerosols, translated by: Daisley, R. E. and Fuchs, M., edited by: Davies, C. N., Pergamon Press, London, 1964.
- Fuchs, N. A. and Sutugin, A. G.: High-dispersed aerosols, in: *Topics in Current Aerosol Research*, edited by: Hidy, G. M. and Brock, J. R., Pergamon, 1, 1971.
- Fuller, E. N., Schettler, P. D., and Giddings, J. C.: New method for prediction of binary gas-phase diffusion coefficients, *Ind. Eng. Chem.*, 58, 18–27, <https://doi.org/10.1021/ie50677a007>, 1966.
- Gagné, S., Lehtipalo, K., Manninen, H. E., Nieminen, T., Schobesberger, S., Franchin, A., Yli-Juuti, T., Boulon, J., Sonntag, A., Mirme, S., Mirme, A., Hörrak, U., Petäjä, T., Asmi, E., and Kulmala, M.: Intercomparison of air ion spectrometers: an evaluation of results in varying conditions, *Atmos. Meas. Tech.*, 4, 805–822, <https://doi.org/10.5194/amt-4-805-2011>, 2011.

- Giorgi, F. and Lionello, P.: Climate change projections for the Mediterranean region, *Global Planet. Change*, 63, 90–104, <https://doi.org/10.1016/j.gloplacha.2007.09.005>, 2008.
- Gong, X., Wex, H., Müller, T., Wiedensohler, A., Höhler, K., Kandler, K., Ma, N., Dietel, B., Schiebel, T., Möhler, O., and Stratmann, F.: Characterization of aerosol properties at Cyprus, focusing on cloud condensation nuclei and ice-nucleating particles, *Atmos. Chem. Phys.*, 19, 10883–10900, <https://doi.org/10.5194/acp-19-10883-2019>, 2019.
- Gonser, S. G., Klein, F., Birmili, W., Größ, J., Kulmala, M., Manninen, H. E., Wiedensohler, A., and Held, A.: Ion – particle interactions during particle formation and growth at a coniferous forest site in central Europe, *Atmos. Chem. Phys.*, 14, 10547–10563, <https://doi.org/10.5194/acp-14-10547-2014>, 2014.
- Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozhkov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R., Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *J. Geophys. Res.-Atmos.*, 122, 8739–8760, <https://doi.org/10.1002/2017jd026844>, 2017.
- Gormley, P. G. and Kennedy, M.: Diffusion from a Stream Flowing through a Cylindrical Tube, *P. Roy. Irish Acad. A*, 52, 163–169, 1948.
- Greene, C. A., Thirumalai, K., Kearney, K. A., Delgado, J. M., Schwanghart, W., Wolfenbarger, N. S., Thyng, K. M., Gwyther, D. E., Gardner, A. S., and Blankenship, D. D.: The Climate Data Toolbox for MATLAB, *Geochem. Geophys. Geos.*, 20, 3774–3781, <https://doi.org/10.1029/2019GC008392>, 2019.
- Größ, J., Hamed, A., Sonntag, A., Spindler, G., Manninen, H. E., Nieminen, T., Kulmala, M., Hörrak, U., Plass-Dülmer, C., Wiedensohler, A., and Birmili, W.: Atmospheric new particle formation at the research station Melpitz, Germany: connection with gaseous precursors and meteorological parameters, *Atmos. Chem. Phys.*, 18, 1835–1861, <https://doi.org/10.5194/acp-18-1835-2018>, 2018.
- Hakala, S., Alghamdi, M. A., Paasonen, P., Vakkari, V., Khoder, M. I., Neitola, K., Dada, L., Abdelmaksoud, A. S., Al-Jeelani, H., Shabbaj, I. I., Almeahadi, F. M., Sundström, A.-M., Lihavainen, H., Kerminen, V.-M., Kontkanen, J., Kulmala, M., Hussein, T., and Hyvärinen, A.-P.: New particle formation, growth and apparent shrinkage at a rural background site in western Saudi Arabia, *Atmos. Chem. Phys.*, 19, 10537–10555, <https://doi.org/10.5194/acp-19-10537-2019>, 2019.
- Hamed, A., Joutsensaari, J., Mikkonen, S., Sogacheva, L., Dal Maso, M., Kulmala, M., Cavalli, F., Fuzzi, S., Facchini, M. C., Decesari, S., Mircea, M., Lehtinen, K. E. J., and Laaksonen, A.: Nucleation and growth of new particles in Po Valley, Italy, *Atmos. Chem. Phys.*, 7, 355–376, <https://doi.org/10.5194/acp-7-355-2007>, 2007.
- Hirsikko, A., Bergman, T., Laakso, L., Dal Maso, M., Riipinen, I., Hörrak, U., and Kulmala, M.: Identification and classification of the formation of intermediate ions measured in boreal forest, *Atmos. Chem. Phys.*, 7, 201–210, <https://doi.org/10.5194/acp-7-201-2007>, 2007.
- Hirsikko, A., Vakkari, V., Tiitta, P., Manninen, H. E., Gagné, S., Laakso, H., Kulmala, M., Mirme, A., Mirme, S., Mabaso, D., Beukes, J. P., and Laakso, L.: Characterisation of sub-micron particle number concentrations and formation events in the western Bushveld Igneous Complex, South Africa, *Atmos. Chem. Phys.*, 12, 3951–3967, <https://doi.org/10.5194/acp-12-3951-2012>, 2012.
- Hirsikko, A., Vakkari, V., Tiitta, P., Hatakka, J., Kerminen, V.-M., Sundström, A.-M., Beukes, J. P., Manninen, H. E., Kulmala, M., and Laakso, L.: Multiple daytime nucleation events in semi-clean savannah and industrial environments in South Africa: analysis based on observations, *Atmos. Chem. Phys.*, 13, 5523–5532, <https://doi.org/10.5194/acp-13-5523-2013>, 2013.
- Hussein, T., Martikainen, J., Junninen, H., Sogacheva, L., Wagner, R., Maso, M. D., Riipinen, I., Aalto, P. P., and Kulmala, M.: Observation of regional new particle formation in the urban atmosphere, *Tellus B*, 60, 509–521, <https://doi.org/10.1111/j.1600-0889.2008.00365.x>, 2008.
- Hussein, T., Atashi, N., Sogacheva, L., Hakala, S., Dada, L., Petäjä, T., and Kulmala, M.: Characterization of Urban New Particle Formation in Amman–Jordan, *Atmosphere*, 11, 79, <https://doi.org/10.3390/atmos11010079>, 2020.
- Jokinen, T., Kontkanen, J., Lehtipalo, K., Manninen, H. E., Aalto, J., Porcar-Castell, A., Garmash, O., Nieminen, T., Ehn, M., Kangasluoma, J., Junninen, H., Levula, J., Duplissy, J., Ahonen, L. R., Rantala, P., Heikkinen, L., Yan, C., Sipilä, M., Worsnop, D. R., Bäck, J., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Solar eclipse demonstrating the importance of photochemistry in new particle formation, *Sci. Rep.-UK*, 7, 45707, <https://doi.org/10.1038/srep45707>, 2017.
- Jun, Y.-S., Jeong, C.-H., Sabaliauskas, K., Richard Leitch, W., and Evans, G. J.: A year-long comparison of particle formation events at paired urban and rural locations, *Atmos. Pollut. Res.*, 5, 447–454, <https://doi.org/10.5094/APR.2014.052>, 2014.
- Kalivitis, N., Stavroulas, I., Bougiatioti, A., Kouvarakis, G., Gagné, S., Manninen, H. E., Kulmala, M., and Mihalopoulos, N.: Night-time enhanced atmospheric ion concentrations in the marine boundary layer, *Atmos. Chem. Phys.*, 12, 3627–3638, <https://doi.org/10.5194/acp-12-3627-2012>, 2012.
- Kalivitis, N., Kerminen, V.-M., Kouvarakis, G., Stavroulas, I., Bougiatioti, A., Nenes, A., Manninen, H. E., Petäjä, T., Kulmala, M., and Mihalopoulos, N.: Atmospheric new particle formation as a source of CCN in the eastern Mediterranean marine boundary layer, *Atmos. Chem. Phys.*, 15, 9203–9215, <https://doi.org/10.5194/acp-15-9203-2015>, 2015.
- Kalivitis, N., Kerminen, V.-M., Kouvarakis, G., Stavroulas, I., Tzitzikalaki, E., Kalkavouras, P., Daskalakis, N., Myriokefalitakis, S., Bougiatioti, A., Manninen, H. E., Roldin, P., Petäjä, T., Boy, M., Kulmala, M., Kanakidou, M., and Mihalopoulos, N.: Formation and growth of atmospheric nanoparticles in the eastern Mediterranean: results from long-term measurements and process simulations, *Atmos. Chem. Phys.*, 19, 2671–2686, <https://doi.org/10.5194/acp-19-2671-2019>, 2019.
- Kalkavouras, P., Bossioli, E., Bezantakos, S., Bougiatioti, A., Kalivitis, N., Stavroulas, I., Kouvarakis, G., Protonotariou, A. P., Dandou, A., Biskos, G., Mihalopoulos, N., Nenes, A., and Tombrou, M.: New particle formation in the southern Aegean Sea during the Etesians: importance for CCN production and

- cloud droplet number, *Atmos. Chem. Phys.*, 17, 175–192, <https://doi.org/10.5194/acp-17-175-2017>, 2017.
- Kalkavouras, P., Bougiatioti, A., Kalivitis, N., Stavroulas, I., Tombrou, M., Nenes, A., and Mihalopoulos, N.: Regional new particle formation as modulators of cloud condensation nuclei and cloud droplet number in the eastern Mediterranean, *Atmos. Chem. Phys.*, 19, 6185–6203, <https://doi.org/10.5194/acp-19-6185-2019>, 2019.
- Kalkavouras, P., Bougiatioti, A., Grivas, G., Stavroulas, I., Kalivitis, N., Liakakou, E., Gerasopoulos, E., Pilinis, C., and Mihalopoulos, N.: On the regional aspects of new particle formation in the Eastern Mediterranean: A comparative study between a background and an urban site based on long term observations, *Atmos. Res.*, 239, 104911, <https://doi.org/10.1016/j.atmosres.2020.104911>, 2020.
- Kammer, J., Perraudin, E., Flaud, P. M., Lamaud, E., Bonnefond, J. M., and Villenave, E.: Observation of nighttime new particle formation over the French Landes forest, *Sci. Total Environ.*, 621, 1084–1092, <https://doi.org/10.1016/j.scitotenv.2017.10.118>, 2018.
- Kangasluoma, J., Cai, R., Jiang, J., Deng, C., Stolzenburg, D., Ahonen, L. R., Chan, T., Fu, Y., Kim, C., Laurila, T. M., Zhou, Y., Dada, L., Sulo, J., Flagan, R. C., Kulmala, M., Petäjä, T., and Lehtipalo, K.: Overview of measurements and current instrumentation for 1–10 nm aerosol particle number size distributions, *J. Aerosol Sci.*, 105584, <https://doi.org/10.1016/j.jaerosci.2020.105584>, 2020.
- Kerminen, V.-M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi, F.: Atmospheric new particle formation and growth: review of field observations, *Environ. Res. Lett.*, 13, 103003, <https://doi.org/10.1088/1748-9326/aadf3c>, 2018.
- Kerminen, V.-M., Paramonov, M., Anttila, T., Riipinen, I., Fountoukis, C., Korhonen, H., Asmi, E., Laakso, L., Lihavainen, H., Swietlicki, E., Svenningsson, B., Asmi, A., Pandis, S. N., Kulmala, M., and Petäjä, T.: Cloud condensation nuclei production associated with atmospheric nucleation: a synthesis based on existing literature and new results, *Atmos. Chem. Phys.*, 12, 12037–12059, <https://doi.org/10.5194/acp-12-12037-2012>, 2012.
- Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagko-georgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid, ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429–433, <https://doi.org/10.1038/nature10343>, 2011.
- Kivekäs, N., Carpmann, J., Roldin, P., Leppä, J., O'Connor, E., Kristensson, A., and Asmi, E.: Coupling an aerosol box model with one-dimensional flow: a tool for understanding observations of new particle formation events, *Tellus B*, 68, 29706, <https://doi.org/10.3402/tellusb.v68.29706>, 2016.
- Kleanthous, S., Vrekoussis, M., Mihalopoulos, N., Kalabokas, P., and Lelieveld, J.: On the temporal and spatial variation of ozone in Cyprus, *Sci. Total Environ.*, 476–477, 677–687, <https://doi.org/10.1016/j.scitotenv.2013.12.101>, 2014.
- Kontkanen, J., Lehtinen, K. E. J., Nieminen, T., Manninen, H. E., Lehtipalo, K., Kerminen, V.-M., and Kulmala, M.: Estimating the contribution of ion–ion recombination to sub-2 nm cluster concentrations from atmospheric measurements, *Atmos. Chem. Phys.*, 13, 11391–11401, <https://doi.org/10.5194/acp-13-11391-2013>, 2013.
- Kopaniakis, I., Chatoutsidou, S. E., Torseth, K., Glytsos, T., and Lazaridis, M.: Particle number size distribution in the eastern Mediterranean: Formation and growth rates of ultrafine airborne atmospheric particles, *Atmos. Environ.*, 77, 790–802, <https://doi.org/10.1016/j.atmosenv.2013.05.066>, 2013.
- Kostopoulou, E. and Jones, P. D.: Comprehensive analysis of the climate variability in the eastern Mediterranean. Part I: map-pattern classification, *Int. J. Climatol.*, 27, 1189–1214, <https://doi.org/10.1002/joc.1467>, 2007a.
- Kostopoulou, E. and Jones, P. D.: Comprehensive analysis of the climate variability in the eastern Mediterranean. Part II: relationships between atmospheric circulation patterns and surface climatic elements, *Int. J. Climatol.*, 27, 1351–1371, <https://doi.org/10.1002/joc.1466>, 2007b.
- Kristensson, A., Dal Maso, M., Swietlicki, E., Hussein, T., Zhou, J., Kerminen, V. M., and Kulmala, M.: Characterization of new particle formation events at a background site in Southern Sweden: relation to air mass history, *Tellus B*, 60, 330–344, <https://doi.org/10.1111/j.1600-0889.2008.00345.x>, 2008.
- Kulmala, M., Maso, M. D., Mäkelä, J. M., Pirjola, L., Väkevä, M., Aalto, P., Miiikkulainen, P., Hämeri, K., and O'dowd, C. D.: On the formation, growth and composition of nucleation mode particles, *Tellus B*, 53, 479–490, <https://doi.org/10.3402/tellusb.v53i4.16622>, 2001.
- Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V.-M.: Measurement of the nucleation of atmospheric aerosol particles, *Nat. Protoc.*, 7, 1651–1667, <https://doi.org/10.1038/nprot.2012.091>, 2012.
- Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger, S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P., Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., Duplissy, J., Vehkamäki, H., Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M. V., Smith, J. N., Ehn, M., Mentel, T. F., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., and Worsnop, D. R.: Direct Observations of Atmospheric Aerosol Nucleation, *Science*, 339, 943–946, <https://doi.org/10.1126/science.1227385>, 2013.
- Kulmala, M., Petäjä, T., Ehn, M., Thornton, J., Sipilä, M., Worsnop, D. R., and Kerminen, V.-M.: Chemistry of Atmospheric Nucleation: On the Recent Advances on Precursor Characterization and Atmospheric Cluster Composition in Connection with Atmospheric New Particle Formation, *Annu. Rev. Phys. Chem.*, 65, 21–37, <https://doi.org/10.1146/annurev-physchem-040412-110014>, 2014.

- Kulmala, M., Kerminen, V. M., Petäjä, T., Ding, A. J., and Wang, L.: Atmospheric gas-to-particle conversion: why NPF events are observed in megacities?, *Faraday Discuss.*, 200, 271–288, <https://doi.org/10.1039/C6FD00257A>, 2017.
- Kürten, A., Li, C., Bianchi, F., Curtius, J., Dias, A., Donahue, N. M., Duplissy, J., Flagan, R. C., Hakala, J., Jokinen, T., Kirkby, J., Kulmala, M., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Onnela, A., Rissanen, M. P., Simon, M., Sipilä, M., Stozhkov, Y., Tröstl, J., Ye, P., and McMurry, P. H.: New particle formation in the sulfuric acid–dimethylamine–water system: reevaluation of CLOUD chamber measurements and comparison to an aerosol nucleation and growth model, *Atmos. Chem. Phys.*, 18, 845–863, <https://doi.org/10.5194/acp-18-845-2018>, 2018.
- Kurtén, T., Loukonen, V., Vehkamäki, H., and Kulmala, M.: Amines are likely to enhance neutral and ion-induced sulfuric acid-water nucleation in the atmosphere more effectively than ammonia, *Atmos. Chem. Phys.*, 8, 4095–4103, <https://doi.org/10.5194/acp-8-4095-2008>, 2008.
- Laaksonen, A., Hamed, A., Joutsensaari, J., Hiltunen, L., Cavalli, F., Junkermann, W., Asmi, A., Fuzzi, S., and Facchini, M. C.: Cloud condensation nucleus production from nucleation events at a highly polluted region, *Geophys. Res. Lett.*, 32, L06812, <https://doi.org/10.1029/2004GL022092>, 2005.
- Lee, H., Lee, K., Lunder, C. R., Krejci, R., Aas, W., Park, J., Park, K.-T., Lee, B. Y., Yoon, Y. J., and Park, K.: Atmospheric new particle formation characteristics in the Arctic as measured at Mount Zeppelin, Svalbard, from 2016 to 2018, *Atmos. Chem. Phys.*, 20, 13425–13441, <https://doi.org/10.5194/acp-20-13425-2020>, 2020.
- Lee, S.-H., Gordon, H., Yu, H., Lehtipalo, K., Haley, R., Li, Y., and Zhang, R.: New Particle Formation in the Atmosphere: From Molecular Clusters to Global Climate, *J. Geophys. Res.-Atmos.*, 124, 7098–7146, <https://doi.org/10.1029/2018jd029356>, 2019.
- Lehtipalo, K., Sipilä, M., Junninen, H., Ehn, M., Berndt, T., Kajos, M. K., Worsnop, D. R., Petäjä, T., and Kulmala, M.: Observations of Nano-CN in the Nocturnal Boreal Forest, *Aerosol Sci. Tech.* 45, 499–509, <https://doi.org/10.1080/02786826.2010.547537>, 2011.
- Lehtipalo, K., Leppä, J., Kontkanen, J., Kangasluoma, J., Franchin, A., Wimmer, D., Schobesberger, S., Junninen, H., Petäjä, T., and Sipilä, M.: Methods for determining particle size distribution and growth rates between 1 and 3 nm using the Particle Size Magnifier, *Boreal Environ. Res.*, 19, 215–236, 2014.
- Lehtipalo, K., Rondo, L., Kontkanen, J., Schobesberger, S., Jokinen, T., Sarnela, N., Kürten, A., Ehrhart, S., Franchin, A., Nieminen, T., Riccobono, F., Sipilä, M., Yli-Juuti, T., Duplissy, J., Adamov, A., Ahlm, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., Dommen, J., Downard, A. J., Dunne, E. M., Flagan, R. C., Guida, R., Hakala, J., Hansel, A., Jud, W., Kangasluoma, J., Kerminen, V.-M., Keskinen, H., Kim, J., Kirkby, J., Kupc, A., Kupiainen-Määttä, O., Laaksonen, A., Lawler, M. J., Leiminger, M., Mathot, S., Olenius, T., Ortega, I. K., Onnela, A., Petäjä, T., Praplan, A., Rissanen, M. P., Ruuskanen, T., Santos, F. D., Schallhart, S., Schnitzhofer, R., Simon, M., Smith, J. N., Tröstl, J., Tsagkogeorgas, G., Tomé, A., Vaattovaara, P., Vehkamäki, H., Vrtala, A. E., Wagner, P. E., Williamson, C., Wimmer, D., Winkler, P. M., Virtanen, A., Donahue, N. M., Carslaw, K. S., Baltensperger, U., Riipinen, I., Curtius, J., Worsnop, D. R., and Kulmala, M.: The effect of acid–base clustering and ions on the growth of atmospheric nano-particles, *Nat. Commun.*, 7, 11594, <https://doi.org/10.1038/ncomms11594>, 2016.
- Leino, K., Nieminen, T., Manninen, H. E., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Intermediate ions as a strong indicator of new particle formation bursts in a boreal forest, *Boreal Environ. Res.*, 21, 274–286, 2016.
- Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Korrman, R., Lawrence, M. G., Levin, Z., Markowicz, K. M., Mihalopoulos, N., Minikin, A., Ramanathan, V., de Reus, M., Roelofs, G. J., Scheeren, H. A., Sciare, J., Schlager, H., Schultz, M., Siegmund, P., Steil, B., Stephanou, E. G., Stier, P., Traub, M., Warneke, C., Williams, J., and Ziereis, H.: Global Air Pollution Crossroads over the Mediterranean, *Science*, 298, 794–799, <https://doi.org/10.1126/science.1075457>, 2002.
- Lelieveld, J., Proestos, Y., Hadjinicolaou, P., Tanarhte, M., Tyrllis, E., and Zittis, G.: Strongly increasing heat extremes in the Middle East and North Africa (MENA) in the 21st century, *Climatic Change*, 137, 245–260, <https://doi.org/10.1007/s10584-016-1665-6>, 2016.
- Lu, Y., Yan, C., Fu, Y., Chen, Y., Liu, Y., Yang, G., Wang, Y., Bianchi, F., Chu, B., Zhou, Y., Yin, R., Baalbaki, R., Garmash, O., Deng, C., Wang, W., Liu, Y., Petäjä, T., Kerminen, V.-M., Jiang, J., Kulmala, M., and Wang, L.: A proxy for atmospheric daytime gaseous sulfuric acid concentration in urban Beijing, *Atmos. Chem. Phys.*, 19, 1971–1983, <https://doi.org/10.5194/acp-19-1971-2019>, 2019.
- Manninen, H. E., Nieminen, T., Riipinen, I., Yli-Juuti, T., Gagné, S., Asmi, E., Aalto, P. P., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Charged and total particle formation and growth rates during EUCAARI 2007 campaign in Hyytiälä, *Atmos. Chem. Phys.*, 9, 4077–4089, <https://doi.org/10.5194/acp-9-4077-2009>, 2009.
- Manninen, H. E., Nieminen, T., Asmi, E., Gagné, S., Häkkinen, S., Lehtipalo, K., Aalto, P., Vana, M., Mirme, A., Mirme, S., Hörrak, U., Plass-Dülmer, C., Stange, G., Kiss, G., Hoffer, A., Törö, N., Moerman, M., Henzing, B., de Leeuw, G., Brinkenberg, M., Kouvarakis, G. N., Bougiatioti, A., Mihalopoulos, N., O’Dowd, C., Ceburnis, D., Arneth, A., Svenningsson, B., Swietlicki, E., Tarozzi, L., Decesari, S., Facchini, M. C., Birmili, W., Sonntag, A., Wiedensohler, A., Boulon, J., Sellegri, K., Laj, P., Gysel, M., Bukowiecki, N., Weingartner, E., Wehrle, G., Laaksonen, A., Hamed, A., Joutsensaari, J., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: EUCAARI ion spectrometer measurements at 12 European sites – analysis of new particle formation events, *Atmos. Chem. Phys.*, 10, 7907–7927, <https://doi.org/10.5194/acp-10-7907-2010>, 2010.
- Manninen, H. E., Mirme, S., Mirme, A., Petäjä, T., and Kulmala, M.: How to reliably detect molecular clusters and nucleation mode particles with Neutral cluster and Air Ion Spectrometer (NAIS), *Atmos. Meas. Tech.*, 9, 3577–3605, <https://doi.org/10.5194/amt-9-3577-2016>, 2016.
- Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, *Atmos. Chem. Phys.*, 9, 8601–8616, <https://doi.org/10.5194/acp-9-8601-2009>, 2009.
- Mikkonen, S., Romakkaniemi, S., Smith, J. N., Korhonen, H., Petäjä, T., Plass-Duelmer, C., Boy, M., McMurry, P. H., Lehtinen, K. E. J., Joutsensaari, J., Hamed, A., Mauldin III, R. L., Bir-

- mili, W., Spindler, G., Arnold, F., Kulmala, M., and Laaksonen, A.: A statistical proxy for sulphuric acid concentration, *Atmos. Chem. Phys.*, 11, 11319–11334, <https://doi.org/10.5194/acp-11-11319-2011>, 2011.
- Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS – a spectrometer for the measurement of cluster ion and nanometer aerosol size distributions, *Atmos. Meas. Tech.*, 6, 1061–1071, <https://doi.org/10.5194/amt-6-1061-2013>, 2013.
- Ndour, M., Conchon, P., D’Anna, B., Ka, O., and George, C.: Photochemistry of mineral dust surface as a potential atmospheric renoxification process, *Geophys. Res. Lett.*, 36, L05816, <https://doi.org/10.1029/2008GL036662>, 2009.
- Nie, W., Ding, A., Wang, T., Kerminen, V.-M., George, C., Xue, L., Wang, W., Zhang, Q., Petäjä, T., Qi, X., Gao, X., Wang, X., Yang, X., Fu, C., and Kulmala, M.: Polluted dust promotes new particle formation and growth, *Sci. Rep.-UK*, 4, 6634, <https://doi.org/10.1038/srep06634>, 2014.
- Nieminen, T., Kerminen, V.-M., Petäjä, T., Aalto, P. P., Arshinov, M., Asmi, E., Baltensperger, U., Beddows, D. C. S., Beukes, J. P., Collins, D., Ding, A., Harrison, R. M., Henzing, B., Hooda, R., Hu, M., Hörrak, U., Kivekäs, N., Komsaare, K., Krejci, R., Kristensson, A., Laakso, L., Laaksonen, A., Leaitch, W. R., Lihavainen, H., Mihalopoulos, N., Németh, Z., Nie, W., O’Dowd, C., Salma, I., Sellegri, K., Svenningsson, B., Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler, A., Wu, Z., Virtanen, A., and Kulmala, M.: Global analysis of continental boundary layer new particle formation based on long-term measurements, *Atmos. Chem. Phys.*, 18, 14737–14756, <https://doi.org/10.5194/acp-18-14737-2018>, 2018.
- Olenius, T. and Riipinen, I.: Molecular-resolution simulations of new particle formation: Evaluation of common assumptions made in describing nucleation in aerosol dynamics models, *Aerosol Sci. Tech.*, 51, 397–408, <https://doi.org/10.1080/02786826.2016.1262530>, 2017.
- Pearson, R. K., Neuvo, Y., Astola, J., and Gabbouj, M.: Generalized Hampel Filters, *EURASIP J. Adv. Sig. Pr.*, 2016, 87, <https://doi.org/10.1186/s13634-016-0383-6>, 2016.
- Petäjä, T., Kerminen, V.-M., Dal Maso, M., Junninen, H., Koponen, I. K., Hussein, T., Aalto, P. P., Andronopoulos, S., Robin, D., Hämeri, K., Bartzis, J. G., and Kulmala, M.: Sub-micron atmospheric aerosols in the surroundings of Marseille and Athens: physical characterization and new particle formation, *Atmos. Chem. Phys.*, 7, 2705–2720, <https://doi.org/10.5194/acp-7-2705-2007>, 2007.
- Petäjä, T., Mauldin, III, R. L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, *Atmos. Chem. Phys.*, 9, 7435–7448, <https://doi.org/10.5194/acp-9-7435-2009>, 2009.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmos. Chem. Phys.*, 7, 1961–1971, <https://doi.org/10.5194/acp-7-1961-2007>, 2007.
- Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation and primary emission rates, *Atmos. Chem. Phys.*, 9, 1339–1356, <https://doi.org/10.5194/acp-9-1339-2009>, 2009.
- Pikridas, M., Bougiatioti, A., Hildebrandt, L., Engelhart, G. J., Kostenidou, E., Mohr, C., Prévôt, A. S. H., Kouvarakis, G., Zarmas, P., Burkhart, J. F., Lee, B.-H., Psichoudaki, M., Mihalopoulos, N., Pilinis, C., Stohl, A., Baltensperger, U., Kulmala, M., and Pandis, S. N.: The Finokalia Aerosol Measurement Experiment – 2008 (FAME-08): an overview, *Atmos. Chem. Phys.*, 10, 6793–6806, <https://doi.org/10.5194/acp-10-6793-2010>, 2010.
- Pikridas, M., Riipinen, I., Hildebrandt, L., Kostenidou, E., Manninen, H., Mihalopoulos, N., Kalivitis, N., Burkhart, J. F., Stohl, A., Kulmala, M., and Pandis, S. N.: New particle formation at a remote site in the eastern Mediterranean, *J. Geophys. Res.-Atmos.*, 117, D12205, <https://doi.org/10.1029/2012JD017570>, 2012.
- Pikridas, M., Vrekoussis, M., Sciare, J., Kleanthous, S., Vasiladou, E., Kizas, C., Savvides, C., and Mihalopoulos, N.: Spatial and temporal (short and long-term) variability of submicron, fine and sub-10 μ m particulate matter (PM₁, PM_{2.5}, PM₁₀) in Cyprus, *Atmos. Environ.*, 191, 79–93, <https://doi.org/10.1016/j.atmosenv.2018.07.048>, 2018.
- Qian, S., Sakurai, H., and McMurry, P. H.: Characteristics of regional nucleation events in urban East St. Louis, *Atmos. Environ.*, 41, 4119–4127, <https://doi.org/10.1016/j.atmosenv.2007.01.011>, 2007.
- Riipinen, I., Pierce, J. R., Yli-Juuti, T., Nieminen, T., Häkkinen, S., Ehn, M., Junninen, H., Lehtipalo, K., Petäjä, T., Slowik, J., Chang, R., Shantz, N. C., Abbatt, J., Leaitch, W. R., Kerminen, V.-M., Worsnop, D. R., Pandis, S. N., Donahue, N. M., and Kulmala, M.: Organic condensation: a vital link connecting aerosol formation to cloud condensation nuclei (CCN) concentrations, *Atmos. Chem. Phys.*, 11, 3865–3878, <https://doi.org/10.5194/acp-11-3865-2011>, 2011.
- Rohan Jayaratne, E., Pushpawela, B., and Morawska, L.: Temporal evolution of charged and neutral nanoparticle concentrations during atmospheric new particle formation events and its implications for ion-induced nucleation, *Front. Env. Sci. Eng.*, 10, 13, <https://doi.org/10.1007/s11783-016-0862-x>, 2016.
- Rose, C., Sellegri, K., Freney, E., Dupuy, R., Colomb, A., Pichon, J.-M., Ribeiro, M., Bourianne, T., Burnet, F., and Schwarzenboeck, A.: Airborne measurements of new particle formation in the free troposphere above the Mediterranean Sea during the HYMEX campaign, *Atmos. Chem. Phys.*, 15, 10203–10218, <https://doi.org/10.5194/acp-15-10203-2015>, 2015.
- Rose, C., Zha, Q., Dada, L., Yan, C., Lehtipalo, K., Junninen, H., Mazon, S. B., Jokinen, T., Sarnela, N., Sipilä, M., Petäjä, T., Kerminen, V.-M., Bianchi, F., and Kulmala, M.: Observations of biogenic ion-induced cluster formation in the atmosphere, *Science Advances*, 4, eaar5218, <https://doi.org/10.1126/sciadv.aar5218>, 2018.
- Salma, I. and Németh, Z.: Dynamic and timing properties of new aerosol particle formation and consecutive growth events, *Atmos. Chem. Phys.*, 19, 5835–5852, <https://doi.org/10.5194/acp-19-5835-2019>, 2019.
- Salma, I., Borsós, T., Weidinger, T., Aalto, P., Hussein, T., Dal Maso, M., and Kulmala, M.: Production, growth and properties of ultrafine atmospheric aerosol particles in an urban environment, *Atmos. Chem. Phys.*, 11, 1339–1353, <https://doi.org/10.5194/acp-11-1339-2011>, 2011.
- Salma, I., Németh, Z., Kerminen, V.-M., Aalto, P., Nieminen, T., Weidinger, T., Molnár, Á., Imre, K., and Kulmala, M.: Regional effect on urban atmospheric nucleation, *Atmos. Chem. Phys.*, 16, 8715–8728, <https://doi.org/10.5194/acp-16-8715-2016>, 2016a.

- Salma, I., Németh, Z., Weidinger, T., Kovács, B., and Kristóf, G.: Measurement, growth types and shrinkage of newly formed aerosol particles at an urban research platform, *Atmos. Chem. Phys.*, 16, 7837–7851, <https://doi.org/10.5194/acp-16-7837-2016>, 2016b.
- Sciare, J.: The Agia Marina Xyliatou Observatory: A remote supersite in Cyprus to monitor changes in the atmospheric composition of the Eastern Mediterranean and the Middle East, EGU General Assembly Conference Abstracts, 2016.
- Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics: from air pollution to climate change, John Wiley & Sons, Hoboken, USA, 2012.
- Sellegri, K., Rose, C., Marinoni, A., Lupi, A., Wiedensohler, A., Andrade, M., Bonasoni, P., and Laj, P.: New Particle Formation: A Review of Ground-Based Observations at Mountain Research Stations, *Atmosphere*, 10, 493, <https://doi.org/10.3390/atmos10090493>, 2019.
- Siakavaras, D., Samara, C., Petrakakis, M., and Biskos, G.: Nucleation events at a coastal city during the warm period: Kerbside versus urban background measurements, *Atmos. Environ.*, 140, 60–68, <https://doi.org/10.1016/j.atmosenv.2016.05.054>, 2016.
- Simon, M., Dada, L., Heinritzi, M., Scholz, W., Stolzenburg, D., Fischer, L., Wagner, A. C., Kürten, A., Rörup, B., He, X.-C., Almeida, J., Baalbaki, R., Baccarini, A., Bauer, P. S., Beck, L., Bergen, A., Bianchi, F., Bräkling, S., Brilke, S., Caudillo, L., Chen, D., Chu, B., Dias, A., Draper, D. C., Duplissy, J., El-Haddad, I., Finkenzeller, H., Frege, C., Gonzalez-Carracedo, L., Gordon, H., Granzin, M., Hakala, J., Hofbauer, V., Hoyle, C. R., Kim, C., Kong, W., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Mai, H., Manninen, H. E., Marie, G., Marten, R., Mentler, B., Molteni, U., Nichman, L., Nie, W., Ojdanic, A., Onnela, A., Partoll, E., Petäjä, T., Pfeifer, J., Philippov, M., Quéléver, L. L. J., Ranjithkumar, A., Rissanen, M. P., Schallhart, S., Schobesberger, S., Schuchmann, S., Shen, J., Sipilä, M., Steiner, G., Stozhkov, Y., Tauber, C., Tham, Y. J., Tomé, A. R., Vazquez-Pufleau, M., Vogel, A. L., Wagner, R., Wang, M., Wang, D. S., Wang, Y., Weber, S. K., Wu, Y., Xiao, M., Yan, C., Ye, P., Ye, Q., Zauner-Wieczorek, M., Zhou, X., Baltensperger, U., Dommen, J., Flagan, R. C., Hansel, A., Kulmala, M., Volkamer, R., Winkler, P. M., Worsnop, D. R., Donahue, N. M., Kirkby, J., and Curtius, J.: Molecular understanding of new-particle formation from α -pinene between -50 and $+25$ °C, *Atmos. Chem. Phys.*, 20, 9183–9207, <https://doi.org/10.5194/acp-20-9183-2020>, 2020.
- Sipilä, M., Berndt, T., Petäjä, T., Brus, D., Vanhanen, J., Stratmann, F., Patokoski, J., Mauldin, R. L., Hyvärinen, A.-P., Lihavainen, H., and Kulmala, M.: The Role of Sulfuric Acid in Atmospheric Nucleation, *Science*, 327, 1243–1246, <https://doi.org/10.1126/science.1180315>, 2010.
- Skrabalova, L., Zikova, N., and Zdimal, V.: Shrinkage of Newly Formed Particles in an Urban Environment, *Aerosol Air Qual. Res.*, 15, 1313–1324, <https://doi.org/10.4209/aaqr.2015.01.0015>, 2015.
- Solomou, E., Poupkou, A., Bolis, S., Zanis, P., Lazaridis, M., and Melas, D.: Evaluating near-surface ozone levels simulated from MACC global and regional modelling systems in Eastern Mediterranean under the influence of Etesian winds, *Atmos. Res.*, 208, 191–200, <https://doi.org/10.1016/j.atmosres.2017.09.010>, 2018.
- Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Mann, G. W., and Sihto, S.-L.: The contribution of boundary layer nucleation events to total particle concentrations on regional and global scales, *Atmos. Chem. Phys.*, 6, 5631–5648, <https://doi.org/10.5194/acp-6-5631-2006>, 2006.
- Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V.-M., Sihto, S.-L., Riipinen, I., Merikanto, J., Mann, G. W., Chipperfield, M. P., Wiedensohler, A., Birmili, W., and Lihavainen, H.: Contribution of particle formation to global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 35, L06808, <https://doi.org/10.1029/2007gl033038>, 2008.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, <https://doi.org/10.5194/acp-5-2461-2005>, 2005.
- Stolzenburg, D., Fischer, L., Vogel, A. L., Heinritzi, M., Schervish, M., Simon, M., Wagner, A. C., Dada, L., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bianchi, F., Breitenlechner, M., Brilke, S., Buenrostro Mazon, S., Chen, D., Dias, A., Draper, D. C., Duplissy, J., El Haddad, I., Finkenzeller, H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., He, X., Helm, J., Hofbauer, V., Hoyle, C. R., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lampilahti, J., Lawler, M., Lehtipalo, K., Leiminger, M., Mai, H., Mathot, S., Mentler, B., Molteni, U., Nie, W., Nieminen, T., Nowak, J. B., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Quéléver, L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S., Tauber, C., Tomé, A., Wagner, R., Wang, M., Weitz, L., Wimmer, D., Xiao, M., Yan, C., Ye, P., Zha, Q., Baltensperger, U., Curtius, J., Dommen, J., Flagan, R. C., Kulmala, M., Smith, J. N., Worsnop, D. R., Hansel, A., Donahue, N. M., and Winkler, P. M.: Rapid growth of organic aerosol nanoparticles over a wide tropospheric temperature range, *P. Natl. Acad. Sci. USA*, 115, 9122, <https://doi.org/10.1073/pnas.1807604115>, 2018.
- Tröstl, J., Chuang, W. K., Gordon, H., Heinritzi, M., Yan, C., Molteni, U., Ahlm, L., Frege, C., Bianchi, F., Wagner, R., Simon, M., Lehtipalo, K., Williamson, C., Craven, J. S., Duplissy, J., Adamov, A., Almeida, J., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Gysel, M., Hansel, A., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Keskinen, H., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lawler, M., Leiminger, M., Mathot, S., Möhler, O., Nieminen, T., Onnela, A., Petäjä, T., Piel, F. M., Miettinen, P., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Sipilä, M., Smith, J. N., Steiner, G., Tomé, A., Virtanen, A., Wagner, A. C., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Carslaw, K. S., Curtius, J., Dommen, J., Kirkby, J., Kulmala, M., Riipinen, I., Worsnop, D. R., Donahue, N. M., and Baltensperger, U.: The role of low-volatility organic compounds in initial particle growth in the atmosphere, *Nature*, 533, 527–531, <https://doi.org/10.1038/nature18271>, 2016.
- Tsagkogeorgas, G., Roldin, P., Duplissy, J., Rondo, L., Tröstl, J., Slowik, J. G., Ehrhart, S., Franchin, A., Kürten, A., Amorim, A., Bianchi, F., Kirkby, J., Petäjä, T., Baltensperger, U., Boy, M., Curtius, J., Flagan, R. C., Kulmala, M., Donahue, N. M., and Stratmann, F.: Evaporation of sulfate aerosols at low relative humidity, *Atmos. Chem. Phys.*, 17, 8923–8938, <https://doi.org/10.5194/acp-17-8923-2017>, 2017.

- Tyrlis, E. and Lelieveld, J.: Climatology and Dynamics of the Summer Etesian Winds over the Eastern Mediterranean, *J. Atmos. Sci.*, 70, 3374–3396, <https://doi.org/10.1175/JAS-D-13-035.1>, 2013.
- Ulbrich, U., Lionello, P., Belušić, D., Jacobeit, J., Knippertz, P., Kuglitsch, F. G., Leckebusch, G. C., Luterbacher, J., Maugeri, M., Maheras, P., Nissen, K. M., Pavan, V., Pinto, J. G., Saaroni, H., Seibert, S., Toretì, A., Xoplaki, E., and Ziv, B.: Climate of the Mediterranean: Synoptic Patterns, Temperature, Precipitation, Winds, and Their Extremes, chap. 5, in: *The Climate of the Mediterranean Region*, edited by: Lionello, P., Elsevier, Oxford, 301–346, 2012.
- Vakkari, V., Laakso, H., Kulmala, M., Laaksonen, A., Mabaso, D., Molefe, M., Kgabi, N., and Laakso, L.: New particle formation events in semi-clean South African savannah, *Atmos. Chem. Phys.*, 11, 3333–3346, <https://doi.org/10.5194/acp-11-3333-2011>, 2011.
- Vanhanen, J., Mikkilä, J., Lehtipalo, K., Sipilä, M., Manninen, H. E., Siivola, E., Petäjä, T., and Kulmala, M.: Particle Size Magnifier for Nano-CN Detection, *Aerosol Sci. Tech.*, 45, 533–542, <https://doi.org/10.1080/02786826.2010.547889>, 2011.
- Wagner, R., Yan, C., Lehtipalo, K., Duplissy, J., Nieminen, T., Kangasluoma, J., Ahonen, L. R., Dada, L., Kontkanen, J., Manninen, H. E., Dias, A., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A.-K., Bianchi, F., Brilke, S., Mazon, S. B., Chen, X., Draper, D. C., Fischer, L., Frege, C., Fuchs, C., Garmash, O., Gordon, H., Hakala, J., Heikkinen, L., Heinritzi, M., Hofbauer, V., Hoyle, C. R., Kirkby, J., Kürten, A., Kvashnin, A. N., Laurila, T., Lawler, M. J., Mai, H., Makhmutov, V., Mauldin III, R. L., Molteni, U., Nichman, L., Nie, W., Ojdanic, A., Onnela, A., Piel, F., Quéléver, L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S., Sengupta, K., Simon, M., Stolzenburg, D., Stozhkov, Y., Tröstl, J., Viisanen, Y., Vogel, A. L., Wagner, A. C., Xiao, M., Ye, P., Baltensperger, U., Curtius, J., Donahue, N. M., Flagan, R. C., Gallagher, M., Hansel, A., Smith, J. N., Tomé, A., Winkler, P. M., Worsnop, D., Ehn, M., Sipilä, M., Kerminen, V.-M., Petäjä, T., and Kulmala, M.: The role of ions in new particle formation in the CLOUD chamber, *Atmos. Chem. Phys.*, 17, 15181–15197, <https://doi.org/10.5194/acp-17-15181-2017>, 2017.
- Wang, M. and Penner, J. E.: Aerosol indirect forcing in a global model with particle nucleation, *Atmos. Chem. Phys.*, 9, 239–260, <https://doi.org/10.5194/acp-9-239-2009>, 2009.
- Wang, M., Kong, W., Marten, R., He, X.-C., Chen, D., Pfeifer, J., Heitto, A., Kontkanen, J., Dada, L., Kürten, A., Yli-Juuti, T., Manninen, H. E., Amanatidis, S., Amorim, A., Baalbaki, R., Baccarini, A., Bell, D. M., Bertozzi, B., Bräkling, S., Brilke, S., Murillo, L. C., Chiu, R., Chu, B., De Menezes, L.-P., Duplissy, J., Finkenzeller, H., Carracedo, L. G., Granzin, M., Guida, R., Hansel, A., Hofbauer, V., Krechmer, J., Lehtipalo, K., Lamkadam, H., Lampimäki, M., Lee, C. P., Makhmutov, V., Marie, G., Mathot, S., Mauldin, R. L., Mentler, B., Müller, T., Onnela, A., Partoll, E., Petäjä, T., Philippov, M., Pospisilova, V., Ranjithkumar, A., Rissanen, M., Rörup, B., Scholz, W., Shen, J., Simon, M., Sipilä, M., Steiner, G., Stolzenburg, D., Tham, Y. J., Tomé, A., Wagner, A. C., Wang, D. S., Wang, Y., Weber, S. K., Winkler, P. M., Wlasits, P. J., Wu, Y., Xiao, M., Ye, Q., Zauner-Wieczorek, M., Zhou, X., Volkamer, R., Riipinen, I., Dommen, J., Curtius, J., Baltensperger, U., Kulmala, M., Worsnop, D. R., Kirkby, J., Seinfeld, J. H., El-Haddad, I., Flagan, R. C., and Donahue, N. M.: Rapid growth of new atmospheric particles by nitric acid and ammonia condensation, *Nature*, 581, 184–189, <https://doi.org/10.1038/s41586-020-2270-4>, 2020.
- Wang, S. C. and Flagan, R. C.: Scanning Electrical Mobility Spectrometer, *Aerosol Sci. Tech.*, 13, 230–240, <https://doi.org/10.1080/02786829008959441>, 1990.
- Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measured atmospheric new particle formation rates: implication for nucleation mechanisms, *Chem. Eng. Commun.*, 151, 53–64, <https://doi.org/10.1080/00986449608936541>, 1996.
- Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measurements of new particle formation and ultrafine particle growth rates at a clean continental site, *J. Geophys. Res.*, 102, 4375–4385, <https://doi.org/10.1029/96jd03656>, 1997.
- Wiedensohler, A., Birmili, W., Nowak, A., Sonntag, A., Weinhold, K., Merkel, M., Wehner, B., Tuch, T., Pfeifer, S., Fiebig, M., Fjåraa, A. M., Asmi, E., Sellegri, K., Depuy, R., Venzac, H., Villani, P., Laj, P., Aalto, P., Ogren, J. A., Swietlicki, E., Williams, P., Roldin, P., Quincey, P., Hüglin, C., Fierz-Schmidhauser, R., Gysel, M., Weingartner, E., Riccobono, F., Santos, S., Grünig, C., Faloon, K., Beddows, D., Harrison, R., Monahan, C., Jennings, S. G., O’Dowd, C. D., Marinoni, A., Horn, H.-G., Keck, L., Jiang, J., Scheckman, J., McMurry, P. H., Deng, Z., Zhao, C. S., Moerman, M., Henzing, B., de Leeuw, G., Lösschauer, G., and Bastian, S.: Mobility particle size spectrometers: harmonization of technical standards and data structure to facilitate high quality long-term observations of atmospheric particle number size distributions, *Atmos. Meas. Tech.*, 5, 657–685, <https://doi.org/10.5194/amt-5-657-2012>, 2012.
- Wildt, J., Mentel, T. F., Kiendler-Scharr, A., Hoffmann, T., Andres, S., Ehn, M., Kleist, E., Müssgen, P., Rohrer, F., Rudich, Y., Springer, M., Tillmann, R., and Wahner, A.: Suppression of new particle formation from monoterpene oxidation by NO_x, *Atmos. Chem. Phys.*, 14, 2789–2804, <https://doi.org/10.5194/acp-14-2789-2014>, 2014.
- Williamson, C. J., Kupc, A., Axisa, D., Bilsback, K. R., Bui, T., Campuzano-Jost, P., Dollner, M., Froyd, K. D., Hodshire, A. L., Jimenez, J. L., Kodros, J. K., Luo, G., Murphy, D. M., Nault, B. A., Ray, E. A., Weinzierl, B., Wilson, J. C., Yu, F., Yu, P., Pierce, J. R., and Brock, C. A.: A large source of cloud condensation nuclei from new particle formation in the tropics, *Nature*, 574, 399–403, <https://doi.org/10.1038/s41586-019-1638-9>, 2019.
- Wu, Z., Hu, M., Liu, S., Wehner, B., Bauer, S., Maßling, A., Wiedensohler, A., Petäjä, T., Dal Maso, M., and Kulmala, M.: New particle formation in Beijing, China: Statistical analysis of a 1-year data set, *J. Geophys. Res.*, 112, D09209, <https://doi.org/10.1029/2006jd007406>, 2007.
- Wyslouzil, B. E., Seinfeld, J. H., Flagan, R. C., and Okuyama, K.: Binary nucleation in acid–water systems. II. Sulfuric acid–water and a comparison with methanesulfonic acid–water, *J. Chem. Phys.*, 94, 6842–6850, <https://doi.org/10.1063/1.460262>, 1991.
- Xie, Y., Ding, A., Nie, W., Mao, H., Qi, X., Huang, X., Xu, Z., Kerminen, V.-M., Petäjä, T., Chi, X., Virkkula, A., Boy, M., Xue, L., Guo, J., Sun, J., Yang, X., Kulmala, M., and Fu, C.: Enhanced sulfate formation by nitrogen dioxide: Implications from in situ observations at the SOR-

- PES station, *J. Geophys. Res.-Atmos.*, 120, 12679–12694, <https://doi.org/10.1002/2015JD023607>, 2015.
- Yan, C., Nie, W., Vogel, A. L., Dada, L., Lehtipalo, K., Stolzenburg, D., Wagner, R., Rissanen, M. P., Xiao, M., Ahonen, L., Fischer, L., Rose, C., Bianchi, F., Gordon, H., Simon, M., Heinritzi, M., Garmash, O., Roldin, P., Dias, A., Ye, P., Hofbauer, V., Amorim, A., Bauer, P. S., Bergen, A., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S. B., Canagaratna, M. R., Chen, X., Ding, A., Dommen, J., Draper, D. C., Duplissy, J., Frege, C., Heyn, C., Guida, R., Hakala, J., Heikkinen, L., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kirkby, J., Kontkanen, J., Kürten, A., Lawler, M. J., Mai, H., Mathot, S., Mauldin, R. L., Molteni, U., Nichman, L., Nieminen, T., Nowak, J., Ojdanic, A., Onnela, A., Pajunoja, A., Petäjä, T., Piel, F., Quéléver, L. L. J., Sarnela, N., Schallhart, S., Sengupta, K., Sipilä, M., Tomé, A., Tröstl, J., Väisänen, O., Wagner, A. C., Ylisirniö, A., Zha, Q., Baltensperger, U., Carslaw, K. S., Curtius, J., Flagan, R. C., Hansel, A., Riipinen, I., Smith, J. N., Virtanen, A., Winkler, P. M., Donahue, N. M., Kerminen, V.-M., Kulmala, M., Ehn, M., and Worsnop, D. R.: Size-dependent influence of NO_x on the growth rates of organic aerosol particles, *Science Advances*, 6, eaay4945, <https://doi.org/10.1126/sciadv.aay4945>, 2020.
- Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S. B., Ehn, M., Paasonen, P., Sipilä, M., Wang, M., Wang, X., Xiao, S., Chen, H., Lu, Y., Zhang, B., Wang, D., Fu, Q., Geng, F., Li, L., Wang, H., Qiao, L., Yang, X., Chen, J., Kerminen, V.-M., Petäjä, T., Worsnop, D. R., Kulmala, M., and Wang, L.: Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278–281, <https://doi.org/10.1126/science.aao4839>, 2018.
- Yao, X., Choi, M. Y., Lau, N. T., Lau, A. P. S., Chan, C. K., and Fang, M.: Growth and Shrinkage of New Particles in the Atmosphere in Hong Kong, *Aerosol Sci. Tech.*, 44, 639–650, <https://doi.org/10.1080/02786826.2010.482576>, 2010.
- Ye, Q., Wang, M., Hofbauer, V., Stolzenburg, D., Chen, D., Schervish, M., Vogel, A., Mauldin, R. L., Baalbaki, R., Brilke, S., Dada, L., Dias, A., Duplissy, J., El Haddad, I., Finkenzeller, H., Fischer, L., He, X., Kim, C., Kürten, A., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Manninen, H. E., Marten, R., Mentler, B., Partoll, E., Petäjä, T., Rissanen, M., Schobesberger, S., Schuchmann, S., Simon, M., Tham, Y. J., Vazquez-Pufleau, M., Wagner, A. C., Wang, Y., Wu, Y., Xiao, M., Baltensperger, U., Curtius, J., Flagan, R., Kirkby, J., Kulmala, M., Volkamer, R., Winkler, P. M., Worsnop, D., and Donahue, N. M.: Molecular Composition and Volatility of Nucleated Particles from α -Pinene Oxidation between -50°C and $+25^\circ\text{C}$, *Environ. Sci. Technol.*, 53, 12357–12365, <https://doi.org/10.1021/acs.est.9b03265>, 2019.
- Yli-Juuti, T., Riipinen, I., Aalto, P. P., Nieminen, T., Maenhaut, W., Janssens, I. A., Claeys, M., Salma, I., Ocskay, R., and Hoffer, A.: Characteristics of new particle formation events and cluster ions at K-pusztá, Hungary, *Boreal Environ. Res.*, 14, 683–698, 2009.
- Yli-Juuti, T., Nieminen, T., Hirsikko, A., Aalto, P. P., Asmi, E., Hörrak, U., Manninen, H. E., Patokoski, J., Dal Maso, M., Petäjä, T., Rinne, J., Kulmala, M., and Riipinen, I.: Growth rates of nucleation mode particles in Hyytiälä during 2003–2009: variation with particle size, season, data analysis method and ambient conditions, *Atmos. Chem. Phys.*, 11, 12865–12886, <https://doi.org/10.5194/acp-11-12865-2011>, 2011.
- Young, L.-H., Lee, S.-H., Kanawade, V. P., Hsiao, T.-C., Lee, Y. L., Hwang, B.-F., Liou, Y.-J., Hsu, H.-T., and Tsai, P.-J.: New particle growth and shrinkage observed in subtropical environments, *Atmos. Chem. Phys.*, 13, 547–564, <https://doi.org/10.5194/acp-13-547-2013>, 2013.
- Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691–7710, <https://doi.org/10.5194/acp-9-7691-2009>, 2009.
- Zhang, J., Chen, Z., Lu, Y., Gui, H., Liu, J., Wang, J., Yu, T., and Cheng, Y.: Observations of New Particle Formation, Subsequent Growth and Shrinkage during Summer-time in Beijing, *Aerosol Air Qual. Res.*, 16, 1591–1602, <https://doi.org/10.4209/aaqr.2015.07.0480>, 2016.
- Zhao, D., Schmitt, S. H., Wang, M., Acir, I.-H., Tillmann, R., Tan, Z., Novelli, A., Fuchs, H., Pullinen, I., Wegener, R., Rohrer, F., Wildt, J., Kiendler-Scharr, A., Wahner, A., and Mentel, T. F.: Effects of NO_x and SO_2 on the secondary organic aerosol formation from photooxidation of α -pinene and limonene, *Atmos. Chem. Phys.*, 18, 1611–1628, <https://doi.org/10.5194/acp-18-1611-2018>, 2018.