


# Parametrizing drifting snow sublimation in the saltation layer

Armin Sigmund<sup>1</sup>, Varun Sharma<sup>1,2</sup>, Daniela Brito Melo<sup>1</sup>, Francesco Comola<sup>1</sup>, Jérôme Dujardin<sup>1,2</sup>, Franziska Gerber<sup>1,2</sup>, Hendrik Huwald<sup>1,2</sup>, and Michael Lehning<sup>1,2</sup>

<sup>1</sup>Laboratory of Cryospheric Sciences, EPFL, Switzerland

<sup>2</sup>SLF, Davos, Switzerland




EPFL



# Goals

- Propose and validate a parametrization for sublimation of saltating snow particles using Large-eddy simulations (LES) as reference
  - Evaluate the performance of the Thorpe-Mason (TM) formula and an alternative
  - Assess the importance of accurate near-surface humidity and temperature profiles



# Thorpe-Mason formula and alternative




- LES domain: 38 x 19 x 18 m<sup>3</sup>
- Reproduce conditions ( $u$ ,  $q$ ,  $T$ ) measured at S17, Antarctica
- Steady state profiles (400-s average)
- Lagrangian particles:

- Sublimation: 
$$\frac{dm_p}{dt} = \pi D d_p (\rho_{w,\infty} - \rho_{w,p}) Sh$$

- Heat balance: 
$$c_i m_p \frac{dT_p}{dt} = L_s \frac{dm_p}{dt} + \pi k d_p (T_{a,\infty} - T_p) Nu$$

TM:  $0 = \underbrace{\Delta \text{storage}}_{c_i m_p \frac{dT_p}{dt}} + \underbrace{\text{Latent heat}}_{L_s \frac{dm_p}{dt}} + \underbrace{\text{Sensible heat}}_{\pi k d_p (T_{a,\infty} - T_p) Nu}$

(Details on method: Sigmund et al., 2021)



Case	$u_{3m}$ (m s <sup>-1</sup> )	RH <sub>1m</sub> (%)	$T_{1m} - T_0$ (K)	$\frac{S_{TM}}{S_{Trans}}$ (%)
A	12	98.7	-0.7	25
B	16	99.9	-0.2	6

# Thorpe-Mason formula and alternative



- LES domain: 38 x 19 x 18 m<sup>3</sup>
- Reproduce conditions ( $u$ ,  $q$ ,  $T$ ) measured at S17, Antarctica
- **Steady state** profiles (400-s average)
- Lagrangian particles:

– Sublimation: 
$$\frac{dm_p}{dt} = \pi D d_p (\rho_{w,\infty} - \rho_{w,p}) Sh$$

– Heat balance: 
$$c_i m_p \frac{dT_p}{dt} = L_s \frac{dm_p}{dt} + \pi k d_p (T_{a,\infty} - T_p) Nu$$


? =  $\underbrace{c_i m_p \frac{dT_p}{dt}}_{\Delta \text{storage}} = \underbrace{L_s \frac{dm_p}{dt}}_{\text{Latent heat}} + \underbrace{\pi k d_p (T_{a,\infty} - T_p) Nu}_{\text{Sensible heat}}$

(Details on method: Sigmund et al., 2021)

- Alternative to TM:

– Estimate empirically 
$$\frac{dT_p}{dt} = f(T_a - T_0, d_p, z, u_*)$$



based on LES output averaged per  $d_p$  class and height



Case	$u_{3m}$ (m s <sup>-1</sup> )	RH <sub>1m</sub> (%)	$T_{1m} - T_0$ (K)	$\frac{S_{TM}}{S_{Trans}}$ (%)
A	12	98.7	-0.7	25
B	16	99.9	-0.2	6

# 1D model for humidity and temperature






- Particle concentration
  - Assume known profile (from LES)
- $T$  and  $q$  boundary conditions (0 m, 9 m)
  - Dirichlet type
- Fluxes at surface
  - Monin-Obukhov (MOST) bulk method
- $T$  and  $q$  initial conditions:
  - $z \geq 2.25$  m: From LES
  - $z < 2.25$  m: MOST profiles
- Simulate until steady state, so far with **TM formula**
  - $z \geq 2.25$  m: Prognostically
  - $z < 2.25$  m: MOST profiles
- **All  $z$ : Prognostically**



# 1D model for humidity and temperature



- Particle concentration
  - Assume known profile (from LES)
- $T$  and  $q$  boundary conditions (0 m, 9 m)
  - Dirichlet type
- Fluxes at surface
  - Monin-Obukhov (MOST) bulk method
- $T$  and  $q$  initial conditions:
  - $z \geq 2.25$  m: From LES
  - $z < 2.25$  m: MOST profiles
- Simulate until steady state, so far with **TM formula**
  - $z \geq 2.25$  m: Prognostically
  - $z < 2.25$  m: MOST profiles
- **All  $z$ : Prognostically**



# Conclusions



A parametrization of sublimation in the saltation layer should ...

... account for an (empirical) expression for  $\frac{dT_p}{dt}$  (to be derived from LES)

... solve prognostically for  $T$  and  $q$  at a few levels

# Thank you!



## References:

- Sharma, V., Comola, F., and Lehning, M., *On the suitability of the Thorpe-Mason model for Calculating Sublimation of Saltating Snow*, *The Cryosphere*, 12, 3499–3509, 2018.
- Sigmund, A., Dujardin, J., Comola, F., Sharma, V., Huwald, H., Melo, D.B., Hirasawa, N., Nishimura, K., Lehning, M., *Evidence of Strong Flux Underestimation by Bulk Parametrizations During Drifting and Blowing Snow*, *Boundary-Layer Meteorol*, 2021.
- Thorpe, A.D. and Mason, B.J., *The evaporation of ice spheres and ice crystals*, *British Journal of Applied Physics* 17, 541–548, 1966.