



Discovery of a Binary-origin Classical Cepheid in a Binary System with a 59 day Orbital Period^{*†}

Bogumił Pilecki¹, Ian B. Thompson², Felipe Espinoza-Arancibia¹, Richard I. Anderson³, Wolfgang Gieren⁴, Weronika Narloch⁴, Javier Minniti¹, Grzegorz Pietrzyński¹, Mónica Taormina¹, Giuseppe Bono⁵, and Gergely Hajdu¹

¹Centrum Astronomiczne im. Mikołaja Kopernika, PAN, Bartycza 18, 00-716 Warsaw, Poland; pilecki@camk.edu.pl

²Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101-1292, USA

³Institute of Physics, Laboratory of Astrophysics, EPFL, Observatoire de Sauverny, 1290 Versoix, Switzerland

⁴Universidad de Concepción, Departamento de Astronomía, Casilla 160-C, Concepción, Chile

⁵Dipartimento di Fisica Università di Roma Tor Vergata, viadella Ricerca Scientifica 1, I-00133 Rome, Italy

Received 2022 September 24; revised 2022 October 31; accepted 2022 November 1; published 2022 November 30

Abstract

We report the discovery of a surprising binary configuration of the double-mode Cepheid OGLE-LMC-CEP-1347 pulsating in the first ($P_1 = 0.690$ days) and second-overtone ($P_2 = 0.556$ days) modes. The orbital period ($P_{\text{orb}} = 59$ days) of the system is five times shorter than the shortest known to date (310 days) for a binary Cepheid. The Cepheid itself is also the shortest-period one ever found in a binary system and the first double-mode Cepheid in a spectroscopically double-lined binary. OGLE-LMC-CEP-1347 is most probably on its first crossing through the instability strip, as inferred from both its short period and fast period increase, consistent with evolutionary models, and from the short orbital period (not expected for binary Cepheids whose components have passed through the red giant phase). Our evolutionary analysis yielded a first-crossing Cepheid with a mass in a range of $2.9\text{--}3.4 M_{\odot}$ (lower than any measured Cepheid mass), consistent with observations. The companion is a stable star, at least two times fainter and less massive than the Cepheid (preliminary mass ratio $q = 0.55$), while also redder and thus at the subgiant or more advanced evolutionary stage. To match these characteristics, the Cepheid has to be a product of binary interaction, most likely a merger of two less massive stars, which makes it the second known classical Cepheid of binary origin. Moreover, further evolution of the components may lead to another binary interaction.

Unified Astronomy Thesaurus concepts: Cepheid variable stars (218); Double-mode Cepheid variable stars (402); Spectroscopic binary stars (1557); Late-type stars (909)

Supporting material: data behind figure

1. Introduction

Classical Cepheids (hereafter also Cepheids) are crucial for various fields of astronomy including stellar oscillations and the evolution of intermediate and massive stars, and with enormous influence on modern cosmology. Since the discovery of the relationship between their pulsation period and luminosity (the Leavitt Law, Leavitt & Pickering 1912), the Cepheids have been extensively used to measure distances in the universe. They are radially pulsating evolved intermediate and high-mass giants and supergiants, mostly located in a well-defined position on the helium-burning loop (called the *blue loop*). Theory predicts masses of Cepheids in the range $3\text{--}11 M_{\odot}$ (Cox 1980; Bono et al. 1999, 2001; Anderson et al. 2016), indicating that unless a binary interaction has occurred in their past, all progenitors are B-type stars (Pecaut & Mamajek 2013; Evans et al. 2020). As such, they are mostly members of binary and multiple systems (Moe & Di Stefano 2017).

For low-mass ($\lesssim 3.5 M_{\odot}$) stars the blue loops predicted by evolution theory are too short to reach the instability strip and

explain the existence of short-period Cepheids (Anderson et al. 2016; De Somma et al. 2021). Such low-mass stars cross the instability strip only during the earlier rapid subgiant phase of evolution (in the Hertzsprung gap), but only a few percent of Cepheids are expected to be found in that stage (De Somma et al. 2021). Nevertheless, these first-crossing Cepheids are currently the only possible explanation for the existence of short-period Cepheids (see, e.g., Ripepi et al. 2022). The difficulty in proving this scenario arises mainly from the low number of accurate dynamical mass measurements of Cepheids. Only seven have been measured thus far (Pilecki et al. 2018b; Gallenne et al. 2018), and all of them occur in a very narrow mass range around $4 M_{\odot}$. The lowest mass, $3.61 M_{\odot}$, was measured for the binary Cepheid OGLE-LMC-CEP-4506 (Gieren et al. 2015; Pilecki et al. 2018b) in the Large Magellanic Cloud (LMC).

Binary systems with Cepheid stars are expected to have orbital periods longer than 200 days. These long periods are required so that the binaries survive the evolution of the components on the red giant branch (Neilson et al. 2015b; Moe & Di Stefano 2017). However, the first-crossing Cepheids can have shorter periods, as they have not yet passed through this stage. At the moment the shortest period measured for a genuine binary Cepheid is 310 days for OGLE-LMC-CEP-0227 (Pietrzyński et al. 2010; Pilecki et al. 2013), while in general periods of about 1 yr or longer are being found.

We have recently started a project to identify and characterize a large sample of Cepheids in spectroscopically double-lined binary (SB2) systems, suitable for dynamical mass measurements. The survey is currently focused on the

* Based on observations collected at the European Southern Observatory, Chile.

† This Letter includes data gathered with the 6.5 m Magellan Clay Telescope at Las Campanas Observatory, Chile.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.




Figure 1. *I*-band light curves showing variability for the first (1O) and second-overtone (2O) modes with the same y-axis span. Data come from the OGLE project (Soszyński et al. 2017) and were disentangled to show variability in each mode separately.

LMC galaxy but is being extended to other galaxies as well. In the first step of the project we have confirmed that virtually all Cepheids that lie significantly above the period–luminosity relation have a red giant companion easily detectable in the spectra (Pilecki et al. 2021). The next step is to confirm the orbital motion of these Cepheids and to determine the orbital periods of the systems.

In this Letter we present the discovery of a surprising binary configuration of the Cepheid OGLE-LMC-CEP-1347⁶ (henceforth also LMC-CEP-1347), one of the 41 SB2 Cepheid candidates presented in Pilecki et al. (2021). In Section 2 we present the spectroscopic data that we have acquired together with the preliminary orbital solution. In Section 3 we discuss the origin and the current state of the Cepheid and draw a few direct conclusions.

2. Spectroscopic Observations

LMC-CEP-1347 ($I = 16.4$ mag, $V = 17.1$ mag; $\alpha_{J2000} = 05^{\text{h}}15^{\text{m}}06^{\text{s}}.42$, $\delta_{J2000} = -69^{\circ}39'52''.8$; Soszyński et al. 2017) is a double-mode Cepheid with a first-overtone (1O) period of $P_1 = 0.690$ day and second-overtone (2O) period of $P_2 = 0.556$ day ($P_2/P_1 = 0.805$). The *I*-band amplitude of this Cepheid is almost six times higher in the 1O mode than in the 2O mode—see Figure 1. We observed this target spectroscopically between 2021 October and 2022 January using two instruments, UVES at the 8.2-m Very Large Telescope UT2 (Paranal, European Southern Observatory, ESO) and MIKE at the 6.5 m Clay Magellan Telescope (Las Campanas, Carnegie) in Chile. Four spectra were taken with the UVES spectrograph (wavelength range 3300–6600 Å) in service mode. With an exposure time of 1400 s, we obtained a typical signal-to-noise ratio (S/N) ~ 10 at resolution $R \sim 60,000$. The spectra were reduced using the *eso-reflex* software available at the ESO software repository.⁷ Five spectra were taken with the MIKE spectrograph (3300–9600 Å) in remote mode with an exposure time of 1800 s (typical S/N ~ 15 at $R \sim 40,000$) and were reduced using Daniel Kelson’s pipeline available at the Carnegie Observatories Software Repository.⁸

We used the broadening function (BF) technique (Rucinski 1992, 1999) implemented in the RaveSpan code (Pilecki et al. 2017) to measure the radial velocities (RVs) of the components of the binary. This technique provides narrower velocity




Figure 2. Broadening function profiles of the Cepheid (blue) and the companion (red) for one of the UVES spectra. The signal from the Cepheid is ~ 4 times stronger indicating the companion is significantly fainter (see text).

profiles than the cross-correlation function method, helping in the separation of components, and increasing the accuracy of the RV measurements. This is important as the companion is significantly fainter than the Cepheid. To measure the BF profiles we used a set of wide wavelength bands between 4100 Å and 6600 Å, omitting broad and atmospheric absorption lines. In eight spectra the profiles of the Cepheid and its companion were clearly separated providing accurate measurements of the RVs of the individual components. For one MIKE spectrum the BF profiles overlap, but velocities of both components could still be measured (they differ by 16 km s^{-1} only). The BF profiles obtained for one of the UVES spectra are shown in Figure 2. The ratio of the integrated profiles is about 4. For stars of the same spectral type this would mean that the companion is roughly four times fainter over the wavelength range used for the BF measurement. The ratio of the integrated BF profiles is ~ 5 for a wavelength range of 4100 Å–5300 Å, while the ratio is ~ 3 for a wavelength range of 5300 Å–6600 Å, indicating that the companion is redder than the Cepheid.

2.1. Orbital Solution

To obtain the orbital solution we again used the RaveSpan code, adjusting the following orbital parameters: the orbital period (P), reference time (T_0), systemic velocity (γ), semiamplitudes (K_1 and K_2), eccentricity (e), and the argument

⁶ in SIMBAD: OGLE LMC-CEP-1347.

⁷ <https://www.eso.org/sci/software/pipelines/>

⁸ <http://code.obs.carnegiescience.edu>

Table 1
Preliminary Orbital Solution for OGLE-LMC-CEP-1347

Parameter	Value	Unit
P	58.85 ± 0.08	days
T_0 (HJD)	2459050.0 ± 0.7	days
$a \sin i$	93.2 ± 1.0	R_\odot
$m_1 \sin^3 i$	2.02 ± 0.04	M_\odot
$m_2 \sin^3 i$	1.12 ± 0.06	M_\odot
$q = m_2/m_1$	0.553 ± 0.016	...
e	0.0	...
γ	239.97 ± 0.13	km s^{-1}
K_1	28.5 ± 0.8	km s^{-1}
K_2	51.56 ± 0.17	km s^{-1}
rms_1	2.61	km s^{-1}
rms_2	0.36	km s^{-1}

Note. Only the statistical error is included. Second-overtone pulsation is not taken into account (note the high rms_1 value).

of periastron (ω). For the current set of data the analysis yielded a circular orbit ($e = 0$). During the fitting the main pulsation variability with the period 0.690 day was described by a second-order Fourier series. The solution is presented in Table 1 and shown in Figure 3. In the table a means semimajor axis, i means orbital inclination, m_i means mass of the i th component, and q means the mass ratio. Because of the low number of observational points we did not take the $2O$ mode into account but its amplitude seems to be significantly lower than for the $1O$ mode with an rms value of 2.6 km s^{-1} for residual RVs compared to 9.8 km s^{-1} if neither mode is included in the fit.

The most striking feature of this system is the unexpectedly short orbital period, much shorter than for any other known binary Cepheid. A total of 1.5 orbital cycles are covered with a uniform distribution, so this detection is firm. The high scatter for the Cepheid RVs has a negligible effect here as the orbital period is constrained mainly by the precise RVs of the companion. The orbital solution we obtained suggests a circular orbit and a high mass ratio of the components with the Cepheid being almost two times more massive than its companion. The minimum mass ($m \sin^3 i$) of the Cepheid is $\sim 2 M_\odot$. Although the solution is still preliminary and the eccentricity and mass ratio may change slightly, these measurements already indicate some interesting aspects of the system.

3. Discussion

Population synthesis done by Neilson et al. (2015b) predicts that binary systems with Cepheids, with components having passed through the red giant phase, cannot have periods shorter than about 200 days. The reason is that binaries with shorter periods would have interacted. This is consistent with observations, as all binary Cepheids confirmed either spectroscopically or astrometrically, have periods of about 1 yr or longer (Evans et al. 2013; Pilecki et al. 2018b). Although some candidates have apparent orbital periods shorter than 300 days (Soszyński et al. 2010), the shortest period measured for a confirmed binary Cepheid is 310 days. That is why the very short period of only 59 days measured for LMC-CEP-1347 is so extraordinary. The most plausible explanation is that the Cepheid is on the first crossing through the instability strip and has not yet evolved onto the red giant branch. This scenario

is supported by the observed rapid period increase $\log \dot{P}_{1O} = -1.00 \pm 0.03 \text{ s yr}^{-1}$ (Pilecki et al. 2021) of the Cepheid. For the fundamentalized period ($P_F = 0.966$ day) we obtain $\log \dot{P}_F = -0.85 \text{ s yr}^{-1}$, which matches theoretical predictions, e.g., $\log \dot{P}_F$ between -0.7 and -0.95 s yr^{-1} in Turner et al. (2006). Moreover, at the LMC metallicity, modern evolutionary models usually do not predict the existence of Cepheids with such short pulsation periods after the red giant branch (Anderson et al. 2016; De Somma et al. 2021).

We calculated evolutionary models using the MESA code (Paxton et al. 2019) for a set of masses and metallicities ($[\text{Fe}/\text{H}] = -0.54$ to -0.27) consistent with that of the LMC (Romaniello et al. 2022), and found the expected mass of a first-crossing Cepheid with a $1O$ period equal to that of LMC-CEP-1347 to be within $2.9\text{--}3.4 M_\odot$. If observationally confirmed, this would be the lowest-mass Cepheid known. We also obtained expected period change rates $\log \dot{P}_{1O}$ between -0.74 and -1.00 s yr^{-1} . Using the information from the orbital solution, the mass of the companion would be $\sim 1.7 M_\odot$ and the orbital inclination around 60° . Assuming the Cepheid is on a higher crossing, we would obtain an even lower mass at which, however, our models do not predict blue loops that enter the instability strip. The evolutionary tracks were calculated taking into account diffusion, semiconvection, thermohaline mixing, and mass loss. We used a mixing length of 1.88 and overshooting mixing efficiency of 0.019, both solar calibrated. The boundaries of the instability strip were obtained applying the MESA Radial Stellar Pulsations package (Smolec & Moskalik 2008) to the evolutionary models.

The preliminary mass ratio of the components is far from unity, while the stars are apparently in a similar evolutionary stage as indicated by their spectral features and comparable brightness. Indeed, subtracting the typical Cepheid brightness obtained from the period–luminosity relations (Soszynski et al. 2008) from the observed brightness in V and I bands, we find that the companion is fainter ($\Delta V = 0.64$ mag) and redder ($\Delta(V - I) = 0.17$ mag) than the Cepheid, which supports the conclusion from the analysis of BF profiles in Section 2. This means that assuming single star evolution the companion would have to be at a similar or even later evolutionary stage than the Cepheid, while having the same age as the Cepheid it would be still on the main sequence as seen in Figure 4. It would be also much bluer (hotter) than the Cepheid. All the observational facts (including the mass ratio and the color difference) can be explained if we assume that LMC-CEP-1347 has been a triple system, with the components of the inner binary having merged creating the current Cepheid with a mass about twice that of the companion. A similar scenario was proposed for the binary Cepheid OGLE-LMC-CEP-1812 (Neilson et al. 2015a) with the mass ratio of about 0.7 and components also at a similar evolutionary stage. Another possibility is mass transfer from the current secondary as in Pilecki et al. (2018a). However in that case we would expect either the current rejuvenated primary (more massive component) to still be on the main sequence or the evolutionarily advanced secondary (less massive component), stripped of the envelope, to be hardly detectable in the spectra.

The short orbital period and low component separation means the Cepheid may in the future again interact, this time with its current companion, either by merging or mass transfer. This is a hint that for some Cepheids the first crossing may also be the last one. As there are many B-type binary systems with similarly short orbital periods (Duchêne & Kraus 2013;




Figure 3. (Left) Orbital radial velocity curves with a preliminary orbital solution. The $1O$ mode variation for the Cepheid was subtracted. The orbit is circular and different semiamplitudes indicate the companion is about two times less massive. (Right) Pulsational RV curve for $1O$ mode with orbital variation removed. High scatter of the Cepheid RVs around the fits comes from the unaccounted $2O$ pulsation. Measurement errors are smaller than the plotting points.

(The data used to create this figure are available.)




Figure 4. Example evolutionary tracks for stars with masses expected for components of OGLE-LMC-CEP-1347 assuming single star evolution. The position of the Cepheid with a $1O$ period of 0.690 day and the corresponding instability strip are shown. The hypothetical position of the companion with the same age as the Cepheid clearly contradicts the observational data (it should be cooler than the Cepheid).

Moe & Di Stefano 2017), the number of first-crossing Cepheids may be higher than typically assumed from a comparison of evolutionary time spent within the instability strip at different crossings (e.g., Ripepi et al. 2022).

Apart from being an example of the complexity of interactions in multiple systems of intermediate and high-mass stars, LMC-CEP-1347 has some other impressive features. It does not only belong to the binary system with about a five times shorter orbital period than previously known for binary Cepheids, but is also the shortest-period Cepheid found in a binary system so far. Although other double-mode Cepheids in binary systems have been identified (e.g., Y Car; Evans 1992), LMC-CEP-1347 is also the first one to be found in an SB2 system. Double-mode Cepheids provide a very robust, purely photometric method to constrain Cepheid masses and luminosities through their period ratios (Petersen 1973; Bono et al. 1996), i.e., independently of the distance and reddening. Unfortunately, this method has never been calibrated

empirically because of a lack of dynamical mass measurements for double-mode Cepheids. LMC-CEP-1347 may thus be the first to be used for that purpose once more data is acquired.

The research leading to these results has received funding from the Polish National Science Center grant SONATA BIS 2020/38/E/ST9/00486. We also acknowledge support from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No. 951549). W.G. gratefully acknowledges support from the ANID BASAL project ACE210002. This work is based on observations collected at the European Southern Observatory under ESO program 108.22BS.001. We also thank Carnegie, and the CNTAC for the generous allocation of observing time for this project. We would like to thank the support staff at the Las Campanas Observatory for their help in remote observations.

This research has made use of NASA’s Astrophysics Data System Service.

Facilities: VLT:Kueyen (UVES), Magellan:Clay (MIKE).

Software: RaveSpan (Pilecki et al. 2017; <https://users.camk.edu.pl/pilecki/ravespan/>), MESA (Paxton et al. 2019; <https://docs.mesastar.org/>).

ORCID iDs

Bogumił Pilecki <https://orcid.org/0000-0003-3861-8124>
 Ian B. Thompson <https://orcid.org/0000-0001-5130-6040>
 Richard I. Anderson <https://orcid.org/0000-0001-8089-4419>
 Wolfgang Gieren <https://orcid.org/0000-0003-1405-9954>
 Mónica Taormina <https://orcid.org/0000-0002-1560-8620>
 Giuseppe Bono <https://orcid.org/0000-0002-4896-8841>
 Gergely Hajdu <https://orcid.org/0000-0003-0594-9138>

References

- Anderson, R. I., Saio, H., Ekström, S., Georgy, C., & Meynet, G. 2016, *A&A*, **591**, A8
- Bono, G., Caputo, F., Castellani, V., & Marconi, M. 1996, *ApJL*, **471**, L33
- Bono, G., Gieren, W. P., Marconi, M., & Fouqué, P. 2001, *ApJL*, **552**, L141
- Bono, G., Marconi, M., & Stellingwerf, R. F. 1999, *ApJS*, **122**, 167
- Cox, A. N. 1980, *ARA&A*, **18**, 15
- De Somma, G., Marconi, M., Cassisi, S., et al. 2021, *MNRAS*, **508**, 1473

- Duchêne, G., & Kraus, A. 2013, [ARA&A](#), **51**, 269
- Evans, N. E., Bond, H. E., Schaefer, G. H., et al. 2013, [AJ](#), **146**, 93
- Evans, N. R. 1992, [ApJ](#), **385**, 680
- Evans, N. R., Günther, H. M., Bond, H. E., et al. 2020, [ApJ](#), **905**, 81
- Gallenne, A., Kervella, P., Evans, N. R., et al. 2018, [ApJ](#), **867**, 121
- Gieren, W., Pilecki, B., Pietrzyński, G., et al. 2015, [ApJ](#), **815**, 28
- Leavitt, H. S., & Pickering, E. C. 1912, *HarCi*, **173**, 1
- Moe, M., & Di Stefano, R. 2017, [ApJS](#), **230**, 15
- Neilson, H. R., Izzard, R. G., Langer, N., & Ignace, R. 2015a, [A&A](#), **581**, L1
- Neilson, H. R., Schneider, F. R. N., Izzard, R. G., Evans, N. R., & Langer, N. 2015b, [A&A](#), **574**, A2
- Paxton, B., Smolec, R., Schwab, J., et al. 2019, [ApJS](#), **243**, 10
- Pecaut, M. J., & Mamajek, E. E. 2013, [ApJS](#), **208**, 9
- Petersen, J. O. 1973, *A&A*, **27**, 89
- Pietrzyński, G., Thompson, I. B., Gieren, W., et al. 2010, *Natur*, **468**, 542
- Pilecki, B., Dervisoğlu, A., Gieren, W., et al. 2018a, [ApJ](#), **868**, 30
- Pilecki, B., Gieren, W., Pietrzyński, G., et al. 2018b, [ApJ](#), **862**, 43
- Pilecki, B., Gieren, W., Smolec, R., et al. 2017, [ApJ](#), **842**, 110
- Pilecki, B., Graczyk, D., Pietrzyński, G., et al. 2013, *MNRAS*, **436**, 953
- Pilecki, B., Pietrzyński, G., Anderson, R. I., et al. 2021, [ApJ](#), **910**, 118
- Ripepi, V., Chemin, L., Molinaro, R., et al. 2022, *MNRAS*, **512**, 563
- Romaniello, M., Riess, A., Mancino, S., et al. 2022, [A&A](#), **658**, A29
- Rucinski, S. 1999, in ASP Conf. Ser. 185, IAU Colloq. 170: Precise Stellar Radial Velocities, ed. J. B. Hearnshaw & C. D. Scarfe (San Francisco, CA: ASP), 82
- Rucinski, S. M. 1992, [AJ](#), **104**, 1968
- Smolec, R., & Moskalik, P. 2008, *AcA*, **58**, 193
- Soszynski, I., Poleski, R., Udalski, A., et al. 2008, *AcA*, **58**, 163
- Soszynski, I., Poleski, R., Udalski, A., et al. 2010, *AcA*, **60**, 17
- Soszynski, I., Udalski, A., Szymański, M. K., et al. 2017, [AcA](#), **67**, 103
- Turner, D. G., Abdel-Sabour Abdel-Latif, M., & Berdnikov, L. N. 2006, *PASP*, **118**, 410