

First diastereoselective formation of lanthanide triple helical complexes with a terdentate chiral C_2 symmetric ligand†

Gilles Muller,^{a‡} Jean-Claude G. Bünzli,^{a§} James P. Riehl,^b Dominique Suhr,^c Alex von Zelewsky^c and Hansruedi Mürner^{*a¶}

^a Institute of Inorganic and Analytical Chemistry, BCH, University of Lausanne, CH-1015 Lausanne, Switzerland. E-mail: hansruedi_murner@hotmail.com

^b Department of Chemistry, University of Minnesota Duluth, Duluth, MN 55812-2496, USA

^c Institute of Inorganic and Analytical Chemistry, University of Fribourg, Pérolles, CH-1700 Fribourg, Switzerland

Received (in Cambridge, UK) 18th April 2002, Accepted 28th May 2002

First published as an Advance Article on the web 13th June 2002


Steric interactions between three enantiopure terdentate ligands leads to the diastereoselective formation of luminescent triple helical lanthanide complexes.

Enantiopure compounds are of fundamental interest in coordination chemistry. In addition to the classical separation of racemates, several approaches for the selective synthesis of desired stereoisomers have emerged in the last decade for transition metal complexes.¹ With lanthanide centers the situation is more delicate since only slight preferences for coordination number and coordination polyhedra exist along the series. Therefore even small changes in reaction conditions can lead to addition or loss of coordinated solvent molecules and/or counter ions, opening pathways for easy racemization.² The design of ligands predisposed to form only one of many possible stereoisomers is therefore a challenging task. Chiral luminescent lanthanide complexes can be envisioned as discriminating probes of local environments. For practical reasons, they should be highly emissive and be conformationally rigid on the timescale of the luminescence emission lifetime. We recently reported on complexes with chiral ligands derived from 2,6-pyridinedicarboxylic acid^{3a} and bis(benzimidazolyl)pyridine^{3b} displaying only slight diastereomeric excesses in solution. The first examples of enantiopure lanthanide complexes involving macrocyclic ligands were reported by Parker and coworkers⁴ formed by chiral octadentate C_4 symmetric ligands of the DOTA family. Here we report on the first selective formation of a single stereoisomer in a mononuclear lanthanide coordination compound resulting from the interplay of three independent terdentate chiral C_2 -symmetrical ligands.

The approach to introduce well defined carbon stereocenters derived from terpene sources into classical coordination ligands was pioneered by the group of von Zelewsky.⁵ Most widely used is the pinene fragment which can be conveniently introduced in various positions in oligopyridine ligands by Kröhnke-type strategy. The steric bulk in the terpene fragments leads to the preferential formation of a limited number of stereoisomers.

We recently reported on the strong luminescence enhancement in lanthanide complexes obtained by alkylation of terpy ligands in the positions 4, 4' and 4".⁶ Starting points for the present investigations were the questions if such an enhancing effect can also be produced by pinene fragments and whether a chiral discrimination in lanthanide complexes of pinene-substituted terpy ligands can be achieved. To test these

hypotheses, we have used the known enantiopure ligand $L(-)$.⁷



Preliminary studies with a chiral terpy containing the pinene group attached in positions 9 and 10 showed formation of only LnL_2 complexes ($Ln = La, Eu, Lu$).⁸ Steric interactions most probably interfere with the coordination of a third ligand in mononuclear complexes as observed in related examples.⁹ No such problems were encountered when the pinene moieties are facing away from the metal center (positions 10 and 11) and complexes $[Ln(L(-))_3](ClO_4)_3$ ($Ln = La, Eu, Tb$) have been isolated in good yields (80–90%, see ESI† for details). The ¹H NMR spectra of solutions of 5×10^{-3} M $Ln(ClO_4)_3 \cdot xH_2O$ ($Ln = La$ or Eu and $x = 0.3-0.6$) and three equivalents of $L(-)$ in anhydrous CD_3CN display only one set of signals at 298 K. This indicates that on the NMR time scale the complexes adopt helical, nine-coordinate C_3 symmetric structures and the 1:3 species is strongly predominant. No additional peaks were detectable in temperature dependent studies in the range 243–298 K with $[Ln(L(-))_3](ClO_4)_3$ ($Ln = La, Eu$) in anhydrous CD_3CN . Fig. 1 displays the results of the experiment with the La complex. Further evidence for the stereoselective helical arrangement of the ligand strands is provided by the specific molar rotatory angle of solutions of 1×10^{-3} M $[Ln(L(-))_3](ClO_4)_3$ ($Ln = La, Eu, Tb$). In all cases the value for the complexes ($[\phi] = -280$ deg $dm^2 mol^{-1}$, La; -600 , Eu; -935 , Tb) is larger than the sum of three ligand molecules ($[\phi] = -68.9$ deg $dm^2 mol^{-1}$)⁷ pointing to the importance of the structural contribution by the chiral helicate.

In contrast to alkylation in the pyridine 4-position⁶ the pinene moiety leads to no strong enhancement of the metal-centered luminescence upon ligand excitation, as demonstrated by the determined quantum yields $Q^{Eu} = 4\%$ and $Q^{Tb} = 0.022\%$. Measurements at 77 K with $[La(L(-))_3](ClO_4)_3$ indicate that an inefficient intersystem crossing in the ligand strands ($I(\text{triplet})/I(\text{singlet}) = 0.033$) is probably one of the main factors leading to such low values. Other processes such as photo induced electron transfers may also be involved.¹⁰ The excitation spectrum of $[Eu(L(-))_3]^{3+}$ in acetonitrile recorded at 295 K in the spectral range of the $^5D_0 \leftarrow ^7F_0$ transition shows the presence of a symmetrical band at $17\,228\text{ cm}^{-1}$ with a small shoulder at lower wavelength. The emission was monitored at wavelengths corresponding to the $^5D_0 \rightarrow ^7F_1$ or 7F_2 transition, and identical spectra were obtained. This result is consistent with the presence of a dominant species in solution. A broader asymmetrical band is observed at $17\,238\text{ cm}^{-1}$ at 10 K for a

† Electronic supplementary information (ESI) available: synthesis and characterisation. Fig. S1: excitation spectra. See <http://www.rsc.org/suppdata/cc/b2/b203691f/>

‡ Present address: Department of Chemistry, University of Minnesota Duluth, Duluth, MN, 55812-2496, USA.

§ Present address: Swiss Federal Institute of Technology, Institute of Molecular and Biological Chemistry, Laboratory of Lanthanide Supramolecular Chemistry, BCH-1402, CH-1015 Lausanne, Switzerland.

¶ Present address: Waldweg 21, CH-1717 St. Ursen, Switzerland.

powdered sample. The solid state result is indicative of a range of distorted Eu(III) environments which may be due to packing effects, or other structural changes occurring at low temperature (Fig. S1, ESI). More details on the photophysical properties will be described elsewhere.

Total luminescence and circularly polarized luminescence spectra for a solution of 5×10^{-3} M $[\text{Eu}(\text{L}(-))_3]^{3+}$ in CH_3CN are shown in Fig. 2. The Eu(III) luminescence is fairly intense when excited at 361 nm into the ligand states. The observation of only two components for the ${}^5\text{D}_0 \rightarrow {}^7\text{F}_1$ transition is compatible with a trigonal symmetry of the complex.

The latter transition is particularly well suited for the measurement of circularly polarized luminescence (CPL), since it satisfies the magnetic-dipole selection rule, $\Delta J = 0, \pm 1$.¹¹ In the spectra shown, however, the magnitude of the luminescence dissymmetry ratio, g_{lum} , defined as follows: $g_{\text{lum}} = 2\Delta I/I = 2(I_L - I_R)/(I_L + I_R)$, where I_L and I_R refer, respectively, to the intensity of left and right circularly polarized light, is approximately the same (0.028 and 0.020 at the intensity maxima) for both transitions. To date there is no reliable quantitative relationship between the sign or magnitude of g_{lum} for Eu(III) transitions and molecular structure, and, for this reason, CPL measurements are most useful in studies aimed at simply verifying the existence of pure enantiomers or racemic mixtures.

A useful experiment to determine whether or not the solution contains a mixture of species is to excite directly the Eu(III) ion with circularly polarized laser light. We have used laser excitation at 557 nm, corresponding to the ${}^5\text{D}_1 \leftarrow {}^7\text{F}_2$ transition. Even though the initial state is only 1% populated at room temperature, this transition satisfies the selection rule given above and may lead to a large chiral discrimination. At this excitation wavelength, g_{lum} is independent of the polarization of the excitation light. Since it is known that the primary source of chirality for Ln(III) ions is the Δ or Λ helical configuration of the chelating ligands, this result is consistent with the presence of only one diastereomer in solution¹² during the lifetime of the excited state which for Eu(III) is in the ms regime. If the solution contained a mixture of diastereomers or enantiomers, the CPL should have been dependent on excitation polarization, as previously observed for 1:3 complexes with related terdentate ligands.³ In these cases, the CPL results were interpreted in terms of the coexistence of 1:2 and 1:3 species in solution.

Finally, it should be mentioned that the corresponding Tb(III) complex is much less luminescent than the Eu(III) analogue, presumably due to deactivation through energy back transfer to




Fig. 2 Total luminescence (bottom) and CPL (top) spectra of the transitions ${}^5\text{D}_0 \rightarrow {}^7\text{F}_1$ (left) and ${}^5\text{D}_0 \rightarrow {}^7\text{F}_2$ (right) of 5×10^{-3} M $[\text{Eu}(\text{L}(-))_3]^{3+}$ in anhydrous MeCN at 295 K, upon excitation at 361 nm.

the ligand, consistent with the temperature dependence of the Tb(${}^5\text{D}_4$) lifetime (from 1.34 to 0.22 ms between 20 and 295 K). Although it was not possible to record a complete CPL spectrum due to very low emission intensity, we were able to measure the luminescence dissymmetry ratio, g_{lum} , for the ${}^5\text{D}_4 \rightarrow {}^7\text{F}_5$ transition of Tb(III) upon indirect excitation of the ligand. The result at the peak wavelength obtained was 0.06, a value three times higher than that observed for a corresponding complex with a derivative of 2,6-pyridinedicarboxylic acid.^{3a}

In conclusion, we have shown the strength of the pinene approach in controlling the stereochemistry of luminescent lanthanide coordination compounds. The resultant emitted light is polarized in a direction determined by the helicity of the metal complex, which in turn is controlled by the absolute configuration at the remote carbon centers in the pinene moieties. Taken into account the easy synthetic availability of both pinene enantiomers, and well-known strategies to generate water robust complexes with a terpy framework,¹³ this opens further opportunities for the synthesis of chiral luminescent probes.

This work is supported through grants by the Swiss National Science Foundation and the University of Minnesota Duluth. We thank M. Düggeli for recording temperature dependent NMR experiments.

Notes and references

- U. Knof and A. von Zelewsky, *Angew. Chem., Int. Ed.*, 1999, **38**, 302.
- D. L. Kepert, *Inorganic Stereochemistry*, Springer-Verlag, Berlin, 1982.
- (a) G. Muller, B. Schmidt, J. Jiricek, G. Hopfgartner, J. P. Riehl, J.-C. G. Bünzli and C. Piguet, *J. Chem. Soc., Dalton Trans.*, 2001, 2655; (b) G. Muller, J. P. Riehl, K. J. Schenk, G. Hopfgartner, C. Piguet and J.-C. G. Bünzli, submitted for publication.
- (a) R. S. Dickins, J. A. K. Howard, C. L. Maupin, J. M. Moloney, D. Parker, J. P. Riehl, G. Siligardi and J. A. G. Williams, *Chem. Eur. J.*, 1999, **5**, 1095; (b) C. L. Maupin, R. S. Dickins, L. G. Govenlock, C. E. Mathieu, D. Parker, J. A. G. Williams and J. P. Riehl, *J. Phys. Chem. A*, 2000, **104**, 6709.
- P. Hayoz, A. von Zelewsky and H. Stoeckli-Evans, *J. Am. Chem. Soc.*, 1993, **115**, 5111.
- H. Mürner, E. Chassat, R. P. Thummel and J.-C. G. Bünzli, *J. Chem. Soc., Dalton Trans.*, 2000, 2809.
- M. Ziegler, V. Monney, H. Stoeckli-Evans, A. von Zelewsky, I. Sakai, G. Dupic, J.-C. Daran and G. G. A. Balavoine, *J. Chem. Soc., Dalton Trans.*, 1999, 667.
- H. Mürner, G. Muller and J.-C. G. Bünzli, unpublished results.
- G. Muller, J.-C. G. Bünzli, K. J. Schenk, C. Piguet and G. Hopfgartner, *Inorg. Chem.*, 2001, **40**, 2642.
- F. R. Gonçalves e Silva, R. Longo, O. L. Malta, C. Piguet and J.-C. G. Bünzli, *Phys. Chem. Chem. Phys.*, 2000, **2**, 5400.
- F. S. Richardson, *Inorg. Chem.*, 1980, **19**, 2806.
- J. P. Riehl and F. S. Richardson, *Chem. Rev.*, 1986, **86**, 1.
- V.-M. Mikkala, C. Sund, M. Kwiatkowski, P. Pasanen, M. Högborg, J. Kankare and H. Takalo, *Helv. Chim. Acta*, 1992, **75**, 1621.




Fig. 1 ${}^1\text{H}$ NMR spectra of 5×10^{-3} M $[\text{La}(\text{L}(-))_3]^{3+}$ in anhydrous CD_3CN , measured as a function of temperature; asterisks correspond to the signal of the solvent.