

# SCIENTIFIC REPORTS

**OPEN**

## Microbial metabolism mediates interactions between dissolved organic matter and clay minerals in streamwater

Received: 09 February 2016

Accepted: 12 July 2016


Published: 02 August 2016

W. R. Hunter<sup>1</sup> & T. J. Battin<sup>2</sup>


Sorption of organic molecules to mineral surfaces is an important control upon the aquatic carbon (C) cycle. Organo-mineral interactions are known to regulate the transport and burial of C within inland waters, yet the mechanisms that underlie these processes are poorly constrained. Streamwater contains a complex and dynamic mix of dissolved organic compounds that coexists with a range of organic and inorganic particles and microorganisms. To test how microbial metabolism and organo-mineral complexation alter amino acid and organic carbon fluxes we experimented with <sup>13</sup>C-labelled amino acids and two common clay minerals (kaolinite and montmorillonite). The addition of <sup>13</sup>C-labelled amino acids stimulated increased microbial activity. Amino acids were preferentially mineralized by the microbial community, concomitant with the leaching of other (non-labelled) dissolved organic molecules that were removed from solution by clay-mediated processes. We propose that microbial processes mediate the formation of organo-mineral particles in streamwater, with potential implications for the biochemical composition of organic matter transported through and buried within fluvial environments.

In fluvial environments, amino acids and other dissolved organic compounds coexist with a range of organic and inorganic particles, onto which they can adsorb and desorb<sup>1,2</sup>. Amino acids are important constituents of the pool of dissolved organic matter (DOM) in aquatic ecosystems. They are prime sources of organic carbon and nitrogen for the microbial metabolism<sup>3,4</sup> and they may serve as olfactory cues mediating animal behaviour in streams and rivers<sup>3-5</sup>. The fine-scale mechanisms that control the persistence of amino acids in streams and rivers, however, remains poorly understood. Organo-mineral complexation is potentially an important control upon the bioavailability of amino acids and other dissolved organic compounds. Organo-mineral complexes form via ion exchange reactions between dissolved organic molecules and the mineral surfaces<sup>1,2,6,7</sup>. Within these reactions, mineral structure is critical, such that the potential for stabilisation of organic molecules against microbial degradation is believed to be an important property of clay minerals<sup>1,7-10</sup>. Kaolinite and montmorillonite are common clay species in soils and sediments<sup>11</sup>, and they highlight the structural variability of these minerals. Kaolinite is composed of a single tetrahedral sheet and a single octahedral sheet held together by hydrogen bonding of aluminol (Al–OH) and siloxane (Si–O) groups. These bonding forces mean that kaolinite is a non-swelling clay with limited cation exchange capacity<sup>11-13</sup>. Montmorillonite is composed of an octahedral sheet of alumina sandwiched between two tetrahedral silicate sheets. These sheets are only weakly bonded and so can become separated by interactions with water (as a polar liquid), providing a greater surface area to which dissolved organic molecules may bind<sup>6,12,13</sup>. Mineral structure and the ambient chemical environment are important properties determining the sorption dynamics of organic molecules to minerals. For instance, it has been shown that the more complex structure of montmorillonite binds more amino acids under abiotic conditions<sup>1,7-9</sup>. The role of microbial activity for organo-mineral interactions and resulting biogeochemical fluxes in aquatic ecosystems are currently poorly

<sup>1</sup>Queen's University Marine Laboratory, School of Biological Sciences, The Queen's University of Belfast, Portaferry, BT22 1PF, United Kingdom. <sup>2</sup>Stream Biofilm and Ecosystem Research Laboratory, School of Architecture, Civil and Environmental Engineering, Ecole Polytechnique Fédérale Lausanne (EPFL), CH-1015 Lausanne, Switzerland. Correspondence and requests for materials should be addressed to H.W.R. (email: w.hunter@qub.ac.uk) or T.J.B. (email: tom.battin@epfl.ch)



**Figure 1. Experimental design matrix.** Illustration of the experimental design, with each treatment was replicated in triplicate independent systems. As the experimental design was not fully factorial, statistical analyses were done by one-way analysis of variance.




**Figure 2. Microbial respiration in the experimental systems.** Boxplots of oxygen consumption over the 72-hour incubations. Letters indicate treatments that are statistically indistinguishable based on post-hoc Tukey's HSD tests ( $p > 0.05$ ). Significant differences between treatments are identified at  $p < 0.05$  (\*);  $p < 0.01$  (\*\*) or  $p < 0.001$  (\*\*\*).

understood<sup>10</sup>. Fine-scale mechanistic understanding of these processes is critical though to predict the dynamics of minerals and DOM, and their link to the carbon cycle in streams and rivers<sup>14</sup>.

In this study we test how the competing processes of microbial metabolism and organo-mineral complexation affect the fate of free amino acids and other dissolved organic molecules in streamwater. We amended natural streamwater microbial communities with <sup>13</sup>C-labelled amino acids and exposed them to different concentrations of initially organic-free kaolinite and montmorillonite particles as outlined in Fig. 1. By tracing the <sup>13</sup>C label in the organic and inorganic carbon pools allowed us to assess the mineralization and retention of the amino acids. Loss of amino acids and DOM was then estimated via mass balance, from which we infer the relative importance of organo-mineral interactions for carbon fluxes in our experimental systems. Organo-mineral sorption is a surface-area limited reaction and we hypothesize that clay type and concentration influence the sorption dynamics of free amino acids and their mineralization by microorganisms.

## Results

**Microbial respiration.** The experimental addition of the <sup>13</sup>C-labelled amino acids and clay particles to natural microbial communities in streamwater had significant effects upon microbial oxygen consumption ( $p < 0.001$ ,  $F = 214.5$ ;  $df = 5$ ) (Fig. 2). Addition of the <sup>13</sup>C-labelled amino acids significantly increased oxygen consumption between the two controls ( $p < 0.001$ ), and the presence of clay particles significantly increased respiration relative to the amino acid control ( $p < 0.001$ ). There were no significant differences in oxygen consumption observed between the four clay treatments (Fig. 2).



**Figure 3. Fluxes of the  $^{13}\text{C}$ -labelled amino acids and non-labelled dissolved organic carbon.** (A) Mineralisation of the amino acids and other DOM, (B) their concentrations remaining in the DOC pool and (C) the residual amino acid and non-labelled DOM concentrations (not detected in the DIC or DOC) after 72 hour incubations. Letters indicate treatments that are statistically indistinguishable based on post-hoc Tukey's HSD tests ( $p > 0.05$ ). Significant differences are between treatments are identified at  $p < 0.05$  (\*);  $p < 0.01$  (\*\*);  $p < 0.001$  (\*\*\*)

**Fluxes of the  $^{13}\text{C}$ -labelled amino acids and non-labelled DOM.** Upon addition of the clay particles we observed a significant increase in the mineralisation of both the  $^{13}\text{C}$ -labelled amino acids ( $p < 0.001$ ,  $F = 38.63$ ;  $df = 4$ ) and non-labelled DOM within streamwater ( $p = 0.001$ ,  $F = 10.96$ ;  $df = 5$ ) (Fig. 3A). This reflects the increased oxygen demand observed in Fig. 2, with no significant differences between clay treatments.


The presence of clay minerals significantly reduced the concentrations of dissolved  $^{13}\text{C}$ -labelled amino acids ( $p < 0.001$ ,  $F = 19.85$ ;  $df = 4$ ) and non-labelled DOM ( $p < 0.001$ ,  $F = 234.40$ ;  $df = 5$ ) (Fig. 3B). There were no significant differences in dissolved  $^{13}\text{C}$ -labelled amino acids between clay treatments. However, non-labelled DOM concentrations showed a treatment effect, with significantly higher DOC concentrations recorded in the  $200 \text{ mg l}^{-1}$  montmorillonite treatment, compared with the other clay treatments (Fig. 3B).

The residual pool of  $^{13}\text{C}$ -labelled amino acids was calculated as the proportion of the  $^{13}\text{C}$ -amino acid not detected in either the DIC or DOC (Fig. 3C). The residual  $^{13}\text{C}$ -labelled amino acids exhibited no difference between the clay treatments and the control ( $p = 0.12$ ,  $F = 2.39$ ;  $df = 4$ ). A residual non-labelled carbon pool was calculated as the proportion non-labelled DOM not detected in either the DIC or DOC of the clay treatments, relative to the amino acid control. In the presence of clay particles there was a significant loss of non-labelled DOM to the residual pool ( $p < 0.001$ ,  $F = 62.61$ ,  $df = 4$ ), with no significant differences between the clay treatments (Fig. 3C).

**Changes in non-labelled DOM concentrations.** The change in the non-labelled DOM concentrations ( $\Delta\text{DOM}$ ) was calculated relative to background streamwater DOC concentration. These data were corrected against the final concentration of dissolved  $^{13}\text{C}$ -labelled amino acids and reveal significant DOM production in the amino acid controls ( $p < 0.001$ ,  $F = 234.4$ ,  $df = 5$ ) (Fig. 4). The  $\Delta\text{DOM}$  values of the clay treatments were comparable with the procedural control.

## Discussion

In this study we tested how changes in clay type (kaolinite or montmorillonite) and concentration ( $0$ ,  $200$  or  $2000 \text{ mg l}^{-1}$ ) influence the sorption dynamics of dissolved amino acids and their mineralization by microorganisms, in streamwater. The study suggests that microbial metabolism was the dominant process determining the fate of dissolved amino acids, with  $55.7 \pm 2.3\%$  of the amino acids mineralised in the clay treatments and  $45.7 \pm 0.9\%$  mineralised in the amino acid control (Fig. 3A). The addition of  $^{13}\text{C}$ -labelled amino acids appears to stimulate microbial activity, with a concomitant increase in the production of non-labelled DOM. This is observable in the amino acid control, however, there was no increase in non-labelled DOM concentrations. This suggests that the presence of clay minerals within the streamwater facilitated the removal of non-labelled



**Figure 4.** Changes in non-labelled DOM concentrations relative to background DOC concentrations ( $120.55 \pm 0.17 \mu\text{mol l}^{-1}$ ), over the 72-hour incubations. Letters indicate treatments that are statistically indistinguishable based on post-hoc Tukey's HSD tests ( $p > 0.05$ ). Significant differences are between treatments are identified at  $p < 0.05$  (\*);  $p < 0.01$  (\*\*) or  $p < 0.001$  (\*\*\*)


DOM. The incubations were limited to 72 hours, over which time dissolved oxygen concentrations in the clay treatments decreased by ~50%. This reflects the time-scales over which previous studies report amino acid sorption to occur<sup>1,2</sup> and mitigates against potentially confounding effects associated with the switch between aerobic and anaerobic metabolism by the streamwater microbial community. However, we acknowledge that the short duration of these experiments limit our capacity to discuss the longer term preservation of carbon within organo-mineral complexes.

Previous studies have focussed on the organo-mineral interactions, both in artificial<sup>1,6</sup> and natural systems<sup>2</sup>, yet microbial contributions to these processes remain understudied. Amino acids readily adsorb to mineral particles, but often this does not offer complete protection from microbial degradation. Instead a portion of the sorbed amino acids remains available for the microbial metabolism<sup>9,10,15</sup>. In aquatic ecosystems, the interplay between clay particles and microorganisms may facilitate the flocculation of dissolved organic matter<sup>16-18</sup>. This process entails the concentration of active microbial cells and dissolved organic matter around a mineral core, which can be sites of elevated microbial activity in stream ecosystems<sup>19</sup>. DOM molecules adsorbed to minerals may thus be brought into close proximity to microorganisms, and the metabolic activity of microorganisms will impact both flocculation and adhesion of DOM. In turn, this may change the chemical composition of organo-mineral particles and their dynamics as they move through the fluvial continuum<sup>19,20</sup>.

Clay minerals often enter fluvial networks via headwater streams where steep hill slopes and loose parent material facilitate erosion<sup>21</sup>. Concentrations of clay minerals used in this study were high, but reflect the sediment loads in streams during storm events<sup>22-24</sup>. During these events, the adsorption of DOC molecules to clay minerals metabolism may compete with microbial heterotrophs during phases of reduced DOC availability. In our experimental systems, microbial activity was the primary control on the fate of the <sup>13</sup>C-labelled amino acids. This underscores the role of amino acids as a source of carbon and nitrogen for microbial metabolism<sup>3,4</sup>, and that they are preferentially metabolised relative to other DOC sources<sup>25</sup>.

We experimentally demonstrate that the addition of amino acids stimulated DOC production by the microbial community (Fig. 4). This supports the notion that microorganisms simultaneously consume and produce dissolved organic matter<sup>26</sup>. However, in the presence of clay particles, there was comparatively little change in the non-labelled DOM pool within the experiment. Given that microbial respiration and amino acid mineralisation both increased in response to clay addition, we would predict non-labelled DOM production to likewise increase. Within the clay treatments, there was little increase of non-labelled DOM (Figs 3B and 4), suggesting the removal of the molecules via organo-mineral sorption (Figs 3C and 4). We propose that microorganisms mediate organo-mineral sorption in aquatic systems. This occurs via preferential mineralisation of labile organic molecules, such as amino acids, leading to the release of non-characterised dissolved organic molecules. These are in turn removed from solution via organo-mineral sorption (Fig. 5). The amino acid doses used within this study were high, reflecting the maximal DOC concentrations detected within our study system (the Oberer Seebach stream). This represents a standard practise in stable-isotope labelling experiments, to ensure the <sup>13</sup>C-label is detectable above the natural variations in the  $\delta^{13}\text{C}$ -signature of each biochemical pool<sup>10,27</sup>. Further study is necessary to fully elucidate how microbial activity influences organo-mineral sorption at natural streamwater DOM concentrations. Here we provide a data-driven hypothesis and conceptual model (Fig. 5) that will support future work on this important biogeochemical question.

Finally, there were no clear trends in the fate of either <sup>13</sup>C-labelled amino acids or non-labelled DOM associated with clay type (kaolinite or montmorillonite) or concentration (200 or 2000 mg l<sup>-1</sup>). The clay concentrations



**Figure 5. Conceptual diagram of the proposed pathways for microbially mediated organo-mineral sorption within the flask experiments.** Here we show the proportion of  $^{13}\text{C}$ -labelled amino acids (black) and non-labelled DOM (white) within the DOC, DIC and Residual C pools. Intermediate Leached DOC proportions are derived from the proportions of  $^{13}\text{C}$ -labelled amino acids and non-labelled DOM in the DOC of Amino Acid Control treatment (no clay).

used in the present study were high, reflecting point source measurements of suspended sediments associated with storm events and mining impacts in high-order streams<sup>22–24</sup>. However, it is likely that the effects of microbial metabolism combined with these high clay concentrations may have obscured the more subtle effects of clay structure upon organo-mineral interactions<sup>1</sup>. Based upon the structural differences between the two clays, we suggest montmorillonite to exhibit a higher carrying capacity for, and stronger sorption of organic molecules<sup>6,12</sup>. However, this is not reflected within our results (Fig. 3). Based on the data available it is not possible to draw any sound conclusion upon these fine-scale processes governing organo-mineral sorption. To fully elucidate these mechanisms would require further experimentation, investigating organo-mineral interactions across a range of clay doses under both biotic and abiotic conditions. Whilst this represents a clear avenue for future study, it does not invalidate our central observation that organo-mineral interactions were governed by microbial metabolic processes, in this study.

## Material and Methods

Experiments were conducted over 72 hours (10 °C, darkness) in pre-combusted 100 mL Schott bottles with no headspace. We used raw streamwater from Oberer Seebach (OSB, Lunz am See, Austria) free of coarse particles (larger than 63  $\mu\text{m}$ ) but that contained the natural microbial community (up to  $6 \times 10^9$  cells liter<sup>-1</sup>). Experiments were conducted in triplicate bottles amended with a 167  $\mu\text{mol C l}^{-1}$  of 10 atom %  $^{13}\text{C}$ -labelled “cell-free” amino acid mix (Euriso-Top GmbH, Germany) and either 0, 200 or 2000  $\text{mg l}^{-1}$  of kaolinite or montmorillonite (Sigma-Aldrich, UK). Clay concentrations were 10- and 100- times the maximal suspended sediment loads of OSB. Triplicate procedural (no clay, no  $^{13}\text{C}$ -labelled amino acids) and amino acid (no clay) controls were established alongside the experimental treatments. In each bottle, oxygen consumption by microorganisms was determined from measurements of dissolved oxygen concentration using a PreSens Oxy-4 optode system (PreSens Precision Sensing GmbH, Germany).

Concentrations and isotopic ratios of DOC and DIC were analysed in the streamwater (as background) and in the assay bottles. DOC concentrations and  $^{12}\text{C}/^{13}\text{C}$  ratios were analysed by liquid chromatography isotope ratio mass spectroscopy (LC-IRMS) using a Dionex High Precision Liquid Chromatograph (Dionex Corporation, Sunnyvale, CA, USA) coupled to a Finnigan Delta V Advantage isotope ratio mass spectrometer via a Finnigan LC Isolink Interface (Thermo Fisher, Bremen, Germany). Analyses were performed on 0.5 ml filtered (Whatman GF/F, 0.7  $\mu\text{m}$ ) water samples acidified with 10 ml of 85% orthophosphoric acid<sup>28</sup>. Concentrations and  $^{12}\text{C}/^{13}\text{C}$  ratios of DIC were determined from 1 ml water samples, injected into  $\text{N}_2$ -flushed 12 ml vials containing 1 ml 85% orthophosphoric acid to liberate gaseous  $\text{CO}_2$  into the headspace. Headspace gas was then analysed via purge-and-trap isotope ratio mass spectrometry (PT-IRMS) using a GasBench II gas preparation unit coupled to a Delta V Advantage IRMS (Thermo Fisher, Bremen, Germany)<sup>27</sup>. All samples were processed within 10 days of sampling. Background DOC and DIC concentrations in the streamwater averaged  $120.55 \pm 0.17 \mu\text{mol l}^{-1}$  and  $2509.9 \pm 52.81 \mu\text{mol l}^{-1}$  at the beginning of the experimental incubations.

The  $^{13}\text{C}$  enrichment of DOC and DIC samples was determined from the  $^{12}\text{C}/^{13}\text{C}$  ratios by

$$X = \left[ \frac{R_{\text{sample}}}{R_{\text{sample}} + 1} \right] \times 100 \quad (1)$$

where  $X$  is the  $^{13}\text{C}$  enrichment (in atom %) and  $R_{\text{sample}}$  is the  $^{12}\text{C}/^{13}\text{C}$  ratio of the sample. We then calculated the concentration of  $^{13}\text{C}$  of each sample ( $E$ ) as

$$E = \frac{(X_{\text{sample}} - X_{\text{control}})}{100} \times S_{\text{conc.}} \quad (2)$$

where  $X_{\text{sample}}$  is the  $^{13}\text{C}$  enrichment of the sample (atom %),  $X_{\text{control}}$  is the background  $^{13}\text{C}$  value (atom %) and  $S_{\text{conc.}}$  is the concentration. Contributions of  $^{13}\text{C}$ -labelled amino acids to the DIC and DOC pools were determined by normalising the  $^{13}\text{C}$  concentrations to the specific labelling of the  $^{13}\text{C}$ -labelled amino acids (10 atom %). The contributions of non-labelled DOM to the DOC pools were calculated as the differences between total DOC and dissolved  $^{13}\text{C}$ -labelled amino acids. The contribution of respired non-labelled DOM to the DIC pool was calculated from the measured changes in oxygen concentration and the  $^{13}\text{C}$ -labelled amino acids contributions to the DIC pool as

$$\text{DIC}_{\text{DOM}} = \left( \Delta[\text{O}_2] - \left( \frac{\text{DIC}_{\text{AA}}}{0.91} \right) \right) \times 1.20 \quad (3)$$

where  $\Delta[\text{O}_2]$  is the measured change in oxygen concentration,  $\text{DIC}_{\text{AA}}$  is the contribution of the  $^{13}\text{C}$ -labelled amino acids to the DIC pool, 0.91 is the respiratory quotient calculated for the  $^{13}\text{C}$ -labelled amino acids (supplementary Table 1) and 1.20 is the Berggren *et al.*'s<sup>29</sup> general respiratory quotient for aquatic bacterioplankton utilising an unknown DOC source. The residual pools of  $^{13}\text{C}$ -labelled amino acids and non-labelled DOM were calculated as the proportion of each not detected in either the DIC or DOC. All data are summarised in Supplementary Tables 2 and 3.

We used a one-way factorial design to test for significant differences in oxygen consumption, and the contributions of  $^{13}\text{C}$ -labelled amino acids and non-labelled DOM to the DIC, DOC and residual C pools. All data were visually explored prior to analysis to ensure their conformity to assumptions of normality and homoscedacity<sup>30</sup>. Differences between treatments were tested using a one-way analysis of variance (ANOVA). Pairwise comparisons were done using the Tukey's Honest Significant Difference. Data analyses were carried out using the *base* and *lattice* packages in R<sup>31,32</sup>.

## References

- Hedges, J. I. & Hare, P. E. Amino acid adsorption by clay minerals in distilled water. *Geochim. Cosmochim. Acta* **51**, 255–259 (1987).
- Aufdenkampe, A. K., Hedges, J. I., Richey, J. E., Krusche, A. V. & Llerena, C. A. Sorptive fractionation of dissolved organic nitrogen and amino acids onto fine sediments within the Amazon Basin. *Limnol. Oceanogr.* **46**, 1921–1935 (2001).
- Hedges, J. I. *et al.* Origins and processing of organic matter in the Amazon River indicated by carbohydrates and amino acids. *Limnol. Oceanogr.* **39**, 743–761 (1994).
- Berggren, M., Hjalmar, L., Haei, M., Ström, L. & Jansson, M. Efficient aquatic bacterial metabolism of dissolved low-molecular-weight compounds from terrestrial sources. *ISME J.* **4**, 408–416 (2010).
- Hassenköhler, T., Pallesen, L. P., Schild, D. & Manzini, I. Amino Acid- vs. Peptide-Odorants: Responses of Individual Olfactory Receptor Neurons in an Aquatic Species. *PLoS One.* **7**, e53097 (2012).
- Benetoli, L. O. B. *et al.* Amino Acid Interaction with and Adsorption on Clays: FT-IR and Mössbauer Spectroscopy and X-ray Diffractometry Investigations. *Orig. Life Evol. Biosph.* **37**, 479–493 (2007).
- Lambert, J. F. Adsorption and Polymerization of Amino Acids on Mineral Surfaces: A Review. *Orig. Life Evol. Biosph.* **38**, 211–242 (2008).
- Theng, B. K. G. Complexes of clay minerals with amino acids and peptides. *Chem. Erd.* **33**, 125–144 (1974).
- Dashman, T. & Stotzky, G. Physical properties of homoionic montmorillonite and kaolinite complexed with amino acids and peptides. *Soil Biol. Biochem.* **17**, 189–195 (1985).
- Hunter, W. R. *et al.* Metabolism of mineral-sorbed organic matter depends upon microbial lifestyle in fluvial ecosystems. *Geophysical Research Letters*, doi: 10.1002/2016GL067719 (2016).
- Mitchell, J. K. *Fundamentals of Soil Behaviour* (3rd edition) John Wiley & Sons, Inc, 592pp (2005).
- Fernandez, R., Fernando Martirena, F. & Scrivener, K. L. The origin of the pozzolanic activity of calcined clay minerals: A comparison between kaolinite, illite and montmorillonite *Cem. Concr. Res.* **41**, 113–122 (2011).
- Deer, W. A., Howie, R. A. & Zussman, J. *An Introduction to the Rock-forming Minerals* (3rd edition). Mineralogical Society of Great Britain and Ireland, London, 498pp (2013).
- Settele, J. *et al.* Terrestrial and inland water systems. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field, C. B. *et al.* (ed.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 271–359 (2014).
- Dippold, M., Biryukova, M. & Kuzyakov, Y. Sorption affects amino acid pathways in soil: Implications from position-specific labeling of alanine. *Soil Biol. Biochem.* **72**, 180–192 (2014).
- Wotton, R. S. *The Biology of particles in aquatic systems*. CRC Press, 336pp (1994).
- Larsen, L. G. *et al.* Predicting organic floc transport dynamics in shallow aquatic ecosystems: Insights from the field, the laboratory, and numerical modeling. *Wat. Resour. Res.* **45**, W01411 (2009).
- Tranvik, L. J. *et al.* 2009. Lakes and reservoirs as regulators of carbon cycling and climate. *Limnol. Oceanogr.* **54**, 2298–314 (2013).
- Richardson, D. C., Newbold, J. D., Aufdenkampe, A. K., Taylor, P. G. & Kaplan, L. A. A method for measuring bacterial mineralization rates of suspended particulate organic carbon in stream ecosystems. *Limnology and Oceanography Methods* **11**, 257–261 (2013).
- Aufdenkampe, A. K. *et al.* Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere. *Front. Ecol. Environ.* **9**, 53–60 (2011).



21. Gomi, T., Sidle, R. C. & Richardson, J. S. Understanding processes and downstream linkages of headwater systems. *BioScience* **52**, 905–916 (2002).
22. Walling, D. E. & Webb, B. W. Sediment availability and the prediction of storm-period sediment yields. Recent Developments in the Explanation and Prediction of Erosion and Sediment Yield (Proceedings of the Exeter Symposium, July 1982). AHS Publ. No. **I 137** (1982).
23. Lenzi, M. A. & Marchi, L. Suspended sediment load during floods in a small stream of the Dolomites in northeastern Italy. *Catena* **39**, 267–282 (2000).
24. López-Tarazón, J. A., Batalla, R. J., Vericat, D. & Balasch, J. C. Rainfall, runoff and sediment transport relations in a mesoscale mountainous catchment: The River Isábena (Ebro basin). *Catena* **82**, 23–34 (2010).
25. Vallino, J., Hopkinson, C. S. & Hobbie, J. E. Modeling bacterial utilization of dissolved organic matter: Optimization replaces Monod growth kinetics *Limnol. Oceanogr.* **41**, 1591–1609 (1996).
26. Guillemette, F. & del Giorgio, P. A. Simultaneous consumption and production of fluorescent dissolved organic matter by lake bacterioplankton. *Env. Microbiol.* **14**, 1432–1443 (2012).
27. Bengtsson, M. M. *et al.* No evidence of aquatic priming effects in hyporheic zone microcosms. *Sci. Rep.* **4**, 5187 (2014).
28. Wild, B., Wanek, W., Postl, W. & Richter, A. Contribution of carbon fixed by Rubisco and PEPC to phloem export in the Crassulacean acid metabolism plant *Kalanchoëdaigremontiana*. *J. Exp. Bot.* **61**, 1375–1383 (2010).
29. Berggren, M., Lapiere, J.-F. & del Giorgio, P. A. Magnitude and regulation of bacterioplankton respiratory quotient across freshwater environmental gradients. *ISME J.* **6**, 984–993 (2012).
30. Zuur, A. F., Ieno, E. N. & Elphick, C. S. A protocol for data exploration to avoid common statistical problems. *Meth. Ecol. Evol.* **1**, 3–14 (2010).
31. R. Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/> (2013).
32. Sarkar, D. *Lattice: Multivariate Data Visualization with R*. Springer-Verlag, New York, USA. URL <http://lmdvr.r-forge.r-project.org/> (2008).

## Acknowledgements

Our thanks to Birgit Wild, Margarete Watzka, Hubert Kraill and Markus Gruber for assistance with the processing and analysis of experimental samples, and to Stephan Krämer for fruitful discussions about experimental design. This study was funded by a Marie Curie Intra-European Fellowship to WRH (FP7-PEOPLE-2011-IEF-302297) and the Austrian Science Fund to TB (START Y420-B17).

## Author Contributions

The study was designed by W.R.H. and T.B. Experimental work and analyses were completed by W.R.H. Both authors contributed to the writing of the manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Hunter, W. R. and Battin, T. J. Microbial metabolism mediates interactions between dissolved organic matter and clay minerals in streamwater. *Sci. Rep.* **6**, 30971; doi: 10.1038/srep30971 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016